

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07C 237/42, 237/44, A61K 31/165, 31/33, C07D 295/18, 295/20, 211/58, 211/46, 213/80, 209/48, C07C 271/16</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 96/41795</b> (43) International Publication Date: 27 December 1996 (27.12.96)</p>
<p>(21) International Application Number: PCT/JP96/01533 (22) International Filing Date: 6 June 1996 (06.06.96) (30) Priority Data: 9511694.3 9 June 1995 (09.06.95) GB (71) Applicant (for all designated States except US): FUJISAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): SETOI, Hiroyuki [JP/JP]; 13-1, Namiki 4-chome, Tsukuba-shi, Ibaraki 305 (JP). OHKAWA, Takehiko [JP/JP]; 944-2, Mukoishige, Ishigemachi, Yuki-gun, Ibaraki 300-27 (JP). ZENKOH, Tatsuya [JP/JP]; 6-3-3-A101, Keyakidai, Moriyamachi, Kitasouma-gun, Ibaraki 302-01 (JP). SAWADA, Hitoshi [JP/JP]; 2-25-10, Matsushiro, Tsukuba-shi, Ibaraki 305 (JP). SATO, Kentaro [JP/JP]; 2-25-10, Matsushiro, Tsukuba-shi, Ibaraki 305 (JP). TANAKA, Hirokazu [JP/JP]; 3-10-21, Hanayashiki Souen, Takarazuka-shi, Hyogo 665 (JP).</p>	<p>(74) Agent: SEKI, Hideo; Fujisawa Pharmaceutical Co., Ltd., Osaka Factory, 1-6, Kashima 2-chome, Yodogawa-ku, Osaka-shi, Osaka 532 (JP). (81) Designated States: AU, CA, CN, HU, IL, JP, KR, MX, NZ, SG, US, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published With international search report.</p>	
<p>(54) Title: BENZAMIDE DERIVATIVES AND THEIR USE AS VASOPRESSIN ANTAGONISTS</p> <div style="text-align: center;"> <p>(I)</p> </div> <div style="text-align: center;"> <p>(a)</p> </div> <p>(57) Abstract</p> <p>This invention relates to new benzamide derivatives having a vasopressin antagonistic activity, etc., and represented by general formula (I), wherein R<sup>1</sup> is aryl optionally substituted with lower alkoxy, etc., R<sup>2</sup> is lower alkyl, etc., R<sup>3</sup> is hydrogen, etc., R<sup>4</sup> is lower alkoxy, etc., R<sup>5</sup> is hydrogen, etc., A is NH, etc., E is (a), etc., X is -CH=CH-, -CH=N-, or S, and Y is CH or N, and pharmaceutically acceptable salts thereof, to processes for preparation thereof and to a pharmaceutical composition comprising the same.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

## DESCRIPTION

## BENZAMIDE DERIVATIVES AND THEIR USE AS VASOPRESSIN ANTAGONISTS

## 5 TECHNICAL FIELD

This invention relates to new benzamide derivatives and pharmaceutically acceptable salts thereof which are useful as a medicament.

## 10 BACKGROUND ART

Some benzamide derivatives have been known as vasopressin antagonist, for example, in PCT International Publication Nos. WO 91/05549 and WO 95/29152, and EP Application Publication No. 0620216.

15

## DISCLOSURE OF INVENTION

This invention relates to new benzamide derivatives and pharmaceutically acceptable salts thereof.

More particularly, it relates to new benzamide  
20 derivatives and pharmaceutically acceptable salts thereof which possess activities as vasopressin antagonistic activity, vasodilating activity, hypotensive activity, activity for inhibiting saccharide release in liver, activity for inhibiting growth of mesangium cells, water diuretic  
25 activity, platelet agglutination inhibitory activity, oxytocin antagonistic activity and the like, to a pharmaceutical composition comprising the same and to a method for the treatment and/or prevention of hypertension, heart failure, renal insufficiency, edema, ascites,  
30 vasopressin parasecretion syndrome, hepatocirrhosis, hyponatremia, hypokalemia, diabetic, circulation disorder, cerebrovascular disease (e.g. cerebral edema, cerebral infarction, etc.), Meniere's syndrome (e.g. Meniere's disease, etc.), motion sickness and the like in human beings  
35 or animals.

- 2 -

One object of this invention is to provide new and useful benzamide derivatives which possess aforesaid activities.

Another object of this invention is to provide processes for the preparation of said benzamide derivatives and salts thereof.

A further object of this invention is to provide a pharmaceutical composition comprising, as an active ingredient, said benzamide derivatives and pharmaceutically acceptable salts thereof.

Still further object of this invention is to provide a therapeutical method for the treatment and/or prevention of aforesaid diseases in human beings or animals, using said benzamide derivatives and pharmaceutically acceptable salts thereof.


The object benzamide derivatives of this invention are new and can be represented by the following general formula (I) :

20

25

30

35



(I)



- 3 -

wherein

- $R^1$  is aryl, cyclo(lower)alkyl or a heterocyclic group,  
each of which may be substituted with substituent(s)  
selected from the group consisting of halogen;  
5 hydroxy; nitro; amino; acyl; substituted acyl;  
acyl(lower)alkylsulfinyl; acyl(lower)alkylsulfonyl;  
acyloxy; lower alkylamino(lower)alkylcarbamoxyloxy;  
aryl; cyano; a heterocyclic group;  
lower alkenyl optionally substituted with acyl,  
10 substituted acyl, aryl or acyl-substituted aryl;  
lower alkynyl optionally substituted with amino,  
acylamino or substituted acylamino;  
lower alkyl optionally substituted with halogen,  
amino, lower alkylamino, acylamino, substituted  
15 acylamino, hydroxy, acyloxy, acyl(lower)alkanoyloxy,  
acyl, substituted acyl, acyl(lower)alkoxyimino, aryl  
or acyl-substituted aryl;  
lower alkylthio optionally substituted with acyl or  
substituted acyl;  
20 alkoxy optionally substituted with aryl, substituted  
aryl, hydroxy, acyloxy, amino, lower alkylamino,  
protected amino, a heterocyclic group, acyl-  
substituted pyridyl, substituted acyl-substituted  
pyridyl, halogen, acyl(lower)alkylamino, N-protected-  
25 acyl(lower)alkylamino, N-acyl(lower)alkyl-N-lower  
alkylamino, acyl, substituted acyl, acylamino,  
substituted acylamino, lower  
alkylhydrazinocarbonylamino, hydroxyimino,  
acyl(lower)alkoxyimino, substituted  
30 acyl(lower)alkoxyimino, acyl(lower)alkoxy, guanidino  
or N-protected guanidino; and  
lower alkenyloxy optionally substituted with acyl or  
substituted acyl;  
 $R^2$  is hydrogen; lower alkyl optionally substituted with  
35 hydroxy, aryl or acyl; or cyclo(lower)alkyl;

- 4 -

- $R^3$  is hydrogen; halogen; hydroxy; acyloxy; substituted  
 acyloxy; lower alkyl optionally substituted with  
 hydroxy or lower alkoxy; lower alkoxy optionally  
 substituted with aryl, amino, protected amino, acyl,  
 5 hydroxy, cyano or lower alkylthio; nitro; amino;  
 acyl; substituted acyl; or cyclo(lower)alkyloxy;
- $R^4$  is hydroxy; halogen; nitro; amino; protected amino;  
 lower alkylamino; acyloxy; amino(lower)alkylamino;  
 N-protected amino(lower)alkylamino;  
 10 lower alkoxy optionally substituted with hydroxy,  
 aryl, substituted aryl, acyl, substituted acyl,  
 amino, lower alkylamino, acylamino, substituted  
 acylamino, protected amino, a heterocyclic group or  
 guanidino; lower alkylthio optionally substituted  
 15 with acyl, substituted acyl, amino, lower alkylamino,  
 acylamino, substituted acylamino, protected amino, a  
 heterocyclic group, hydroxy, lower alkylsulfonyloxy,  
 arylsulfonyloxy, ar(lower)alkoxy or substituted  
 ar(lower)alkoxy; lower alkyl substituted with acyl,  
 20 substituted acyl, amino, lower alkylamino, acylamino,  
 substituted acylamino, protected amino, a  
 heterocyclic group, hydroxy, lower alkylsulfonyloxy  
 or arylsulfonyloxy; lower alkenyl optionally  
 substituted with acyl; lower alkynyl optionally  
 25 substituted with hydroxy, amino, protected amino,  
 lower alkylsulfonyloxy or arylsulfonyloxy;  
 amino(lower)alkylsulfonyl; N-protected  
 amino(lower)alkylsulfonyl; lower alkylaminosulfonyl;  
 a heterocyclicsulfonyl; amino(lower)alkylsulfinyl;  
 30 N-protected amino(lower)alkylsulfinyl; piperidyloxy;  
 or N-protected piperidyloxy;
- $R^5$  is hydrogen, lower alkyl, lower alkoxy or halogen;  
 A is a single bond, O or NH;
- 35 E is lower alkylene, lower alkenylene,  $-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ ,  $-\overset{\text{O}}{\underset{\text{O}}{\text{S}}}-$ , or

- 5 -

a group of the formula :

-G-J-

5 in which G is lower alkylene and J is O or  $\begin{array}{c} \text{R}^6 \\ | \\ \text{-N-} \end{array}$

(wherein  $\text{R}^6$  is hydrogen or N-protective group);

X is  $-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{N}-$  or S; and

Y is CH or N;


10 and pharmaceutically acceptable salts thereof.

The object compound (I) for its salt can be prepared by the processes as illustrated in the following reaction schemes.

15

Process 1

20




25

(II)

or its salt

+



(III)


or its reactive derivative  
at the carboxy group  
or the sulfo group,  
or a salt thereof

30

35

- 6 -


5



10

Process 2


15



20

(IV)  
or its salt


+



(V)  
or its reactive derivative  
at the carboxy group  
or a salt thereof


25

30





35

- 7 -




Process 3

10


Process 4Process 5

35


- 8 -

Process 6Process 7Process 8

- 9 -


Process 9

10

Process 10

20

25


Process 11

35

- 10 -

Process 12


5



10

Process 13

15




20

25

Process 14


30




35




- 11 -

Process 15

10


15 Process 16

25


Process 17

35

- 12 -


Process 18

- 13 -


Process 21

10


15

Process 22


25

Process 23

- 14 -


Process 24

10

Process 25


20

25


Process 26

35

- 15 -


Process 27

10

Process 28

20

25


Process 29

35

- 16 -

Process 30

5




10

Process 31

15

20




25

Process 32

30


35



- 17 -

Process 33


5



10

Process 34

15




20

25


Process 35

30




35


- 18 -

Process 36

10


15 Process 37

25

Process 38




- 19 -

Process 39

15

wherein

$R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,



25  $R_a^1$  is aryl, haloaryl, cyclo(lower)alkyl or a heterocyclic group, each of which is substituted with esterified carboxy; lower alkenyl substituted with esterified carboxy or esterified carboxy-substituted aryl;

30 lower alkyl substituted with esterified carboxy, esterified carboxy(lower)alkanoyloxy or esterified carboxy(lower)alkoxyimino;

lower alkylthio substituted with esterified carboxy; alkoxy substituted with esterified carboxy-substituted aryl, esterified carboxy-substituted pyridyl, esterified carboxy(lower)alkylamino, N-protected-esterified carboxy(lower)alkylamino, N-esterified carboxy(lower)alkyl-N-lower alkylamino, esterified carboxy or esterified carboxy(lower)alkoxyimino; or

35 lower alkenyloxy substituted with esterified carboxy;

- 20 -

- $R_D^1$  is aryl, haloaryl, cyclo(lower)alkyl or  
a heterocyclic group, each of which is substituted  
with carboxy; lower alkenyl substituted with carboxy  
or carboxy-substituted aryl;  
5 lower alkyl substituted with carboxy, carboxy(lower)-  
alkanoyloxy or carboxy(lower)alkoxyimino;  
lower alkylthio substituted with carboxy;  
alkoxy substituted with carboxy-substituted aryl,  
carboxy-substituted pyridyl, carboxy(lower)-  
10 alkylamino, N-protected-carboxy(lower)alkylamino,  
N-carboxy(lower)alkyl-N-lower alkylamino, carboxy or  
carboxy(lower)alkoxyimino; or lower alkenyloxy  
substituted with carboxy;
- $R_a^4$  is lower alkoxy substituted with esterified carboxy;  
15 lower alkylthio substituted with esterified carboxy;  
lower alkyl substituted with esterified carboxy; or  
lower alkenyl substituted with esterified carboxy;
- $R_D^4$  is lower alkoxy substituted with carboxy; lower  
alkylthio substituted with carboxy; lower alkyl  
20 substituted with carboxy; or lower alkenyl  
substituted with carboxy;
- $R_C^4$  is protected amino; N-protected piperidyloxy;  
N-protected amino(lower)alkylamino;  
lower alkoxy substituted with protected amino;  
25 lower alkylthio substituted with protected amino;  
lower alkyl substituted with protected amino;  
lower alkynyl substituted with protected amino; or  
N-protected amino(lower)alkylsulfonyl;
- $R_D^4$  is amino; piperidyloxy; amino(lower)alkylamino;  
30 lower alkoxy substituted with amino; lower alkylthio  
substituted with amino; lower alkyl substituted with  
amino; lower alkynyl substituted with amino;  
or amino(lower)alkylsulfonyl;
- $R_C^1$  is aryl, haloaryl, cyclo(lower)alkyl or a heterocyclic  
35 group, each of which is substituted with substituted

- 21 -

or unsubstituted N-containing heterocycliccarbonyl;  
carbamoyl; substituted or unsubstituted lower  
alkylcarbamoyl; lower alkenyl substituted with  
substituted or unsubstituted N-containing  
5 heterocycliccarbonyl, carbamoyl, substituted or  
unsubstituted lower alkylcarbamoyl or N-containing  
heterocycliccarbonyl-substituted aryl;  
lower alkyl substituted with substituted or  
unsubstituted N-containing heterocycliccarbonyl,  
10 carbamoyl, substituted or unsubstituted lower  
alkylcarbamoyl, substituted or unsubstituted  
N-containing heterocycliccarbonyl(lower)alkanoyloxy,  
carbamoyl(lower)alkanoyloxy, substituted or  
unsubstituted lower alkylcarbamoyl(lower)alkanoyloxy,  
15 substituted or unsubstituted N-containing  
heterocycliccarbonyl(lower)alkoxyimino, carbamoyl-  
(lower)alkoxyimino or substituted or unsubstituted  
lower alkylcarbamoyl(lower)alkoxyimino;  
lower alkylthio substituted with substituted or  
20 unsubstituted N-containing heterocycliccarbonyl,  
carbamoyl or substituted or unsubstituted lower  
alkylcarbamoyl; alkoxy substituted with substituted  
or unsubstituted N-containing heterocycliccarbonyl-  
substituted aryl, carbamoyl-substituted aryl,  
25 substituted or unsubstituted lower alkylcarbamoyl-  
substituted aryl, substituted or unsubstituted  
N-containing heterocycliccarbonyl-substituted  
pyridyl, carbamoyl-substituted pyridyl, substituted  
or unsubstituted lower alkylcarbamoyl-substituted  
30 pyridyl, substituted or unsubstituted  
N-containing heterocycliccarbonyl(lower)alkylamino,  
carbamoyl(lower)alkylamino, substituted or  
unsubstituted lower alkylcarbamoyl(lower)alkylamino,  
N-protected-(substituted or unsubstituted  
35 N-containing heterocyclic)carbonyl(lower)alkylamino,

- 22 -

- N-protected-carbamoyl(lower)alkylamino, N-protected  
 substituted or unsubstituted lower alkylcarbamoyl-  
 (lower)alkylamino, N-(substituted or unsubstituted  
 N-containing heterocyclic)carbonyl(lower)alkyl-N-  
 5 lower alkylamino, N-carbamoyl(lower)alkyl-N-lower  
 alkylamino, substituted or unsubstituted N-lower  
 alkylcarbamoyl-N-lower alkylamino, substituted or  
 unsubstituted N-containing heterocycliccarbonyl,  
 carbamoyl, aminocarbamoyl, pyridylcarbamoyl,  
 10 N-(lower alkyl)piperazinylcarbonyl,  
 substituted or unsubstituted lower alkylcarbamoyl,  
 substituted or unsubstituted N-containing  
 heterocycliccarbonyl(lower)alkoxyimino,  
 carbamoyl(lower)alkoxyimino or substituted or  
 15 unsubstituted lower alkylcarbamoyl(lower)alkoxyimino;  
 or lower alkenyloxy substituted with substituted or  
 unsubstituted N-containing heterocycliccarbonyl,  
 carbamoyl or substituted or unsubstituted lower  
 alkylcarbamoyl;
- 20  $R_e^4$  is lower alkoxy, lower alkylthio, lower alkyl or lower  
 alkenyl, each of which is substituted with  
 substituted or unsubstituted N-containing  
 heterocycliccarbonyl, carbamoyl, or substituted or  
 unsubstituted lower alkylcarbamoyl;
- 25  $R_f^4$  is methoxy substituted with aryl or substituted aryl;  
 or lower alkylthio which is substituted with methoxy  
 substituted with aryl or substituted aryl;
- $R_g^4$  is hydroxy; or lower alkylthio substituted with hydroxy;  
 $R_{ga}^4$  is hydroxy;
- 30  $R^7$  is lower alkyl substituted with hydroxy, aryl,  
 substituted aryl, acyl, amino, lower alkylamino,  
 acylamino, protected amino or a heterocyclic group;  
 or N-protected piperidyl;
- $Z^1$  is hydroxy; or acid residue;
- 35  $R_h^4$  is lower alkoxy substituted with hydroxy, aryl,

- 23 -

- substituted aryl, acyl, amino, lower alkylamino, acylamino, protected amino or a heterocyclic group; or N-protected piperidyloxy;
- 5  $R_{da}^4$  is lower alkoxy substituted with amino; lower alkylthio substituted with amino; or lower alkyl substituted with amino;
- 10  $R_i^4$  is lower alkoxy substituted with acylamino or substituted acylamino; lower alkylthio substituted with acylamino or substituted acylamino; or lower alkyl substituted with acylamino or substituted acylamino;
- 15  $R_{db}^4$  is amino; lower alkoxy substituted with amino; lower alkylthio substituted with amino; or lower alkyl substituted with amino;
- 20  $R_j^4$  is lower alkoxy substituted with lower alkylamino; lower alkylthio substituted with lower alkylamino; lower alkyl substituted with lower alkylamino; lower alkylamino; or N-protected amino(lower)alkylamino;
- $R_k^4$  is acyloxy;
- $Z^2$  is acid residue;
- $E_b$  is lower alkylene;
- $R_l^4$  is lower alkylthio substituted with amino or protected amino;
- 25  $R_m^4$  is lower alkylsulfinyl substituted with amino or protected amino, or lower alkylsulfonyl substituted with amino or protected amino;
- $E_c$  is lower alkenylene;
- 30  $R_d^1$  is aryl which is substituted with methoxy substituted with aryl or substituted aryl;
- $R_e^1$  is aryl which is substituted with hydroxy;
- $Z^3$  is hydroxy; or acid residue;
- 35  $R^8$  is lower alkyl optionally substituted with acyl, acylamino, protected amino, aryl, substituted aryl, acyl-substituted pyridyl or N-protected guanidino;

- 24 -

- $R_f^1$  is aryl which is substituted with lower alkoxy  
optionally substituted with acyl, acylamino,  
protected amino, aryl, substituted aryl, acyl-  
substituted pyridyl or N-protected guanidino;
- 5  $R_a^3$  is methoxy substituted with aryl; acyloxy; or  
substituted acyloxy;
- $Z^4$  is acid residue;
- $R^9$  is lower alkyl optionally substituted with esterified  
carboxy;
- 10  $R_b^3$  is lower alkoxy optionally substituted with esterified  
carboxy;
- $R_c^3$  is lower alkoxy substituted with esterified carboxy;
- $R_d^3$  is lower alkoxy substituted with carboxy;
- $R_n^4$  is halogen;
- 15  $R_o^4$  is lower alkynyl optionally substituted with hydroxy,  
amino, protected amino, lower alkylsulfonyloxy or  
arylsulfonyloxy;
- $R_p^4$  is lower alkylthio, lower alkyl or lower alkynyl, each  
of which is substituted with hydroxy;
- 20  $Z^5$  is halogen;
- $R^{10}$  is lower alkylsulfonyl or arylsulfonyl;
- $R_q^4$  is lower alkylthio, lower alkyl or lower alkynyl, each  
of which is substituted with lower alkylsulfonyloxy  
or arylsulfonyloxy;
- 25  $R_r^4$  is lower alkylthio, lower alkyl or lower alkynyl, each  
of which is substituted with phthalimido;
- $R_s^4$  is lower alkyl optionally substituted with hydroxy,  
amino, protected amino, lower alkylsulfonyloxy or  
arylsulfonyloxy;
- 30  $E_d$  is a single bond or lower alkylene;
- $Z^6$  is acid residue;
- $R_a^2$  is lower alkyl optionally substituted with aryl or  
acyl;
- 35  $R_g^1$  is aryl which is substituted with lower alkoxy  
substituted with amino;

- 25 -

- $R_n^1$  is aryl which is substituted with lower alkoxy  
substituted with acylamino or substituted acylamino;
- $R_l^1$  is aryl which is substituted with lower alkoxy  
substituted with oxopiperidylcarbonyl;
- 5  $R_j^1$  is aryl which is substituted with lower alkoxy  
substituted with hydroxypiperidylcarbonyl;
- $R_k^1$  is aryl which is substituted with lower alkoxy  
substituted with formyl or oxopiperidylcarbonyl;
- 10  $R_q^1$  is aryl which is substituted with lower alkoxy  
substituted with aminopiperidylcarbonyl or N-lower  
alkylpiperazinyl;
- $R_m^1$  is aryl which is substituted with lower alkoxy  
substituted with carboxy;
- 15  $R_n^1$  is aryl which is substituted with lower alkoxy  
substituted with lower alkylamino(lower)-  
alkoxycarbonyl;
- $R_o^1$  is aryl which is substituted with lower alkoxy  
substituted with esterified carboxy;
- 20  $R_p^1$  is aryl which is substituted with lower alkoxy  
substituted with hydroxy;
- $R_q^1$  is aryl which is substituted with lower alkoxy  
substituted with formyl;
- $R_r^1$  is aryl which is substituted with lower alkoxy  
substituted with cyano-substituted aryl;
- 25  $R_s^1$  is aryl which is substituted with lower alkoxy  
substituted with tetrazolyl-substituted aryl;
- $R_t^4$  is lower alkoxy substituted with amino;
- $R_u^4$  is lower alkoxy substituted with guanidino;
- 30  $R_t^1$  is aryl which is substituted with lower alkoxy  
substituted with protected amino, N-protected  
amino(lower)alkanoylamino, N-protected  
piperazinylcarbonyl or N-protected guanidino;
- 35  $R_u^1$  is aryl which is substituted with lower alkoxy  
substituted with amino, amino(lower)alkanoylamino,  
piperazinylcarbonyl or guanidino;

- 26 -

$R_V^1$  is aryl which is substituted with lower alkoxy substituted with phenoxycarbonylamino;

$R_W^1$  is aryl which is substituted with lower alkoxy substituted with N-lower  
5 alkympiperazinylcarbonylamino, dimethylaminopiperidylcarbonylamino, carbamoylamino or dimethylcarbamoylamino; and

$R_E^3$  is lower alkoxy which is substituted with carbamoyl optionally substituted with lower alkyl.

10 In the above and subsequent description of the present specification, suitable examples of the various definitions to be included within the scope of the invention are explained in detail in the following.

15 The term "lower" is intended to mean a group having 1 to 6 carbon atom(s), unless otherwise provided.

The "higher" is intended to mean 7 to 20 carbon atoms, unless otherwise provided.

20 The lower moiety in the terms "cyclo(lower)alkyl" and "cyclo(lower)alkyloxy" is intended to mean a group having 3 to 6 carbon atoms.

The lower moiety in the terms "lower alkenyl", "lower alkenyloxy" and "lower alkynyl" is intended to mean a  
25 group having 2 to 6 carbon atoms.

The term "alkoxy" may included lower alkoxy and higher alkoxy.

Suitable "lower alkoxy" and lower alkoxy moiety in the terms "acyl(lower)alkoxy", "acyl(lower)alkoxyimino",  
30 "esterified carboxy(lower)alkoxyimino", "carboxy(lower)alkoxyimino", "N-containing heterocycliccarbonyl(lower)alkoxyimino", "carbamoyl(lower)alkoxyimino", "lower alkylcarbamoyl-(lower)alkoxyimino", "lower alkoxycarbonyl" and  
35 "ar(lower)alkoxy" may be straight or branched  $C_1-C_6$



- 27 -

alkoxy such as methoxy, ethoxy, propoxy, isopropoxy, methylpropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, hexyloxy or the like.

Suitable "higher alkoxy" may be straight or branched  
 5 C<sub>7</sub>-C<sub>20</sub> alkoxy such as heptyloxy, octyloxy, nonyloxy, decyloxy, undecyloxy, dodecyloxy, tridecyloxy, tetradecyloxy, pentadecyloxy, hexadecyloxy, heptadecyloxy, octadecyloxy, nonadecyloxy, eicosyloxy, methylheptyloxy, methyloctyloxy, methylnonyloxy, methyldecyloxy,  
 10 ethylheptyloxy, ethyloctyloxy, ethylnonyloxy, ethyldecyloxy or the like, in which preferable one is heptyloxy.

Suitable "lower alkyl" and lower alkyl moiety in the terms "acyl(lower)alkylsulfinyl", "acyl(lower)alkyl-sulfonyl", "lower alkylamino(lower)alkylcarbamoxyloxy",  
 15 "acyl(lower)alkylamino", "N-protected-acyl(lower)-alkylamino", "N-acyl(lower)alkyl-N-lower alkylamino", "lower alkylhydrazinocarbonylamino", "esterified carboxy(lower)alkylamino", "N-protected-esterified carboxy(lower)alkylamino", "N-esterified  
 20 carboxy(lower)alkyl-N-lower alkylamino", "carboxy(lower)alkylamino", "N-protected-carboxy(lower)-alkylamino", "N-carboxy(lower)alkyl-N-lower alkylamino", "lower alkylcarbamoxy", "lower alkylcarbamoxy(lower)alkanoyloxy", "lower  
 25 alkylcarbamoxy(lower)alkoxyimino", "lower alkylthio", "N-protected-(substituted or unsubstituted N-containing heterocyclic)carbonyl(lower)alkylamino", "N-protected-carbamoyl(lower)alkylamino", "N-protected-substituted or unsubstituted lower alkylcarbamoxy(lower)alkylamino",  
 30 "N-(substituted or unsubstituted N-containing heterocyclic)carbonyl(lower)alkyl-N-lower alkylamino", "N-carbamoyl(lower)alkyl-N-lower alkylamino", "N-lower alkylcarbamoxy-N-lower alkylamino", "lower alkylcarbamoxy-(lower)alkoxyimino", "1-hydroxy(lower)alkyl", "1-(lower  
 35 alkyl)amino(lower)alkyl", "mono(lower)alkylamino",

- 28 -

"acyl(lower)alkyl", "di(lower)alkylamino", "lower alkylsulfinyl", "lower alkylsulfonyl", "lower alkylamino", "amino(lower)alkylamino", "N-protected amino(lower)alkylamino", "lower alkylsulfonyloxy",  
5 "amino(lower)alkylsulfonyl", "N-protected amino(lower)alkylsulfonyl", "lower alkylaminosulfonyl", "amino(lower)alkylsulfinyl" and "N-protected amino(lower)alkylsulfinyl" may be straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl such as methyl, ethyl, propyl, isopropyl,  
10 butyl, isobutyl, tert-butyl, pentyl, ethylpropyl, hexyl or the like.

Suitable "cyclo(lower)alkyl" and cyclo(lower)alkyl moiety in the term "cyclo(lower)alkyloxy" may be cyclo(C<sub>3</sub>-C<sub>6</sub>)alkyl such as cyclopropyl, cyclobutyl,  
15 cyclopentyl or cyclohexyl, in which preferable one is cyclopentyl or cyclohexyl.

Suitable "lower alkenyl" and lower alkenyl moiety in the term "lower alkenyloxy" may be straight and branched C<sub>2</sub>-C<sub>6</sub> alkenyl such as ethenyl, propenyl, pentenyl,  
20 isopropenyl, butenyl, hexenyl or the like, in which preferable one is ethenyl, propenyl, pentenyl or hexenyl.

Suitable "lower alkynyl" may be straight and branched C<sub>2</sub>-C<sub>6</sub> alkynyl such as ethynyl, propargyl, butynyl or the like, in which preferable one is butynyl.

25 Suitable "aryl" and aryl moiety in the terms "haloaryl", "arylsulfonyl", "acyl-substituted aryl", "ar(lower)alkoxy", "substituted ar(lower)alkoxy" and "arylsulfonyloxy" may be phenyl, naphthyl, phenyl substituted with lower alkyl [e.g. tolyl, xylyl, mesityl, cumenyl, di(tert-butyl)phenyl, etc.] and the like, in  
30 which preferable one is phenyl, tolyl or xylyl.

Suitable "substituted aryl" may be aryl substituted with suitable substituent(s) such as acyl, substituted acyl, N-protected piperazinylsulfonyl,  
35 piperazinylsulfonyl, N-lower alkylpiperazinylsulfonyl,

- 29 -

hydroxy(lower)alkyl, a heterocyclic(lower)alkyl, halogen, nitro, amino, lower alkylamino, a heterocyclic group [e.g. thiazolyl, oxazolyl, tetrazolyl, oxazolinyl, pyridyl, pyrimidinyl, pyrrolyl optionally substituted with lower alkyl and cyano, etc.], cyano, lower alkoxy or the like, in which preferable one for the substituent of alkoxy for R<sup>1</sup> is aryl substituted with N-methylpiperazinylsulfonyl, N-t-butoxycarbonyl-piperazinylsulfonyl, piperazinylsulfonyl, carboxy, esterified carboxy, N-lower alkylpiperazinylcarbonyl, lower alkanoyl, hydroxy(lower)alkyl, N-lower alkyl-piperazinyl(lower)alkyl, thiazolyl, oxazolyl, tetrazolyl, oxazolinyl, pyridyl, pyrimidinyl, pyrrolyl substituted with lower alkyl and cyano, cyano, lower alkoxy, lower alkylaminopiperidylcarbonyl, and preferable one for R<sup>4</sup> is aryl substituted with halogen, nitro, amino, lower alkylamino or lower alkoxy.

Suitable "halogen" and halo moiety in the term "haloaryl" may be fluorine, chlorine, bromine and iodine, in which preferable one is chlorine or bromine.

Suitable "lower alkylamino" and lower alkylamino moiety in the terms "lower alkylamino(lower)-alkylcarbamoxyloxy", "acyl(lower)alkylamino", "esterified carboxy(lower)alkylamino", "carboxy(lower)alkylamino", "N-containing heterocycliccarbonyl(lower)alkylamino", "carbamoxy(lower)alkylamino", "lower alkylcarbamoxy(lower)alkylamino" "amino(lower)alkylamino", "N-protected amino(lower)alkylamino", "lower alkylaminosulfonyl" and "lower alkylaminopiperidylcarbonyl" may be mono or di(lower alkyl)amino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, tert-butylamino, isobutylamino, pentylamino, hexylamino, dimethylamino, diethylamino, dipropylamino, dibutylamino, diisopropylamino, dipentylamino, dihexylamino, N-methylethylamino or the like, in which preferable one is

- 30 -

methylamino, dimethylamino or diethylamino.

Suitable "1-hydroxy(lower)alkyl" may be 1-hydroxy-(C<sub>1</sub>-C<sub>6</sub>)alkyl such as hydroxymethyl, 1-hydroxyethyl, 1-hydroxypropyl, 1-hydroxybutyl, 1-hydroxy-3-methylpropyl or the like, in which preferable one is hydroxymethyl or 1-hydroxyethyl.

Suitable "1-(lower alkyl)amino(lower)alkyl" may be 1-mono or di(C<sub>1</sub>-C<sub>6</sub> alkyl)amino(C<sub>1</sub>-C<sub>6</sub>)alkyl such as methylaminomethyl, dimethylaminomethyl, 1-methylaminoethyl, 1-dimethylaminoethyl, ethylaminomethyl, 1-ethylaminoethyl or the like, in which preferable one is methylaminomethyl, dimethylaminomethyl, 1-methylaminoethyl or 1-dimethylaminoethyl.

Suitable "heterocyclic group" may be one containing at least one hetero atom selected from nitrogen, sulfur and oxygen atom, and may include saturated or unsaturated, monocyclic or polycyclic heterocyclic group, and preferable heterocyclic group may be N-containing heterocyclic group such as unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl [e.g. 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.], tetrazolyl [e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.], etc.; saturated 3 to 7-membered heteromonocyclic group containing 1 to 4 nitrogen atoms [e.g. pyrrolidinyl, imidazolidinyl, piperidyl, piperazinyl, homopiperazinyl, etc.]; unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, imidazopyridyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl [e.g. tetrazolo[1,5-b]pyridazinyl, etc.], etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl,

- 31 -

etc.;

saturated 3 to 6-membered heteromonocyclic group

containing an oxygen atom, for example,

1H-tetrahydropyranyl, tetrahydrofuranyl, etc.;

5     unsaturated, 3 to 6-membered heteromonocyclic group  
containing 1 to 2 sulfur atoms, for example, thienyl, etc.;

unsaturated 3 to 6-membered heteromonocyclic group  
containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms,  
for example, oxazolyl, isoxazolyl, oxadiazolyl [e.g.

10     1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl,  
etc.], oxazolinyl [e.g. 2-oxazolinyl, etc.], etc.;

saturated 3 to 6-membered heteromonocyclic group  
containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms  
[e.g. morpholinyl, etc.];

15     unsaturated condensed heterocyclic group containing 1 to 2  
oxygen atoms and 1 to 3 nitrogen atoms [e.g.

benzofurazanyl, benzoxazolyl, benzoxadiazolyl, etc.];

unsaturated 3 to 6-membered heteromonocyclic group  
containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms,

20     for example, thiazolyl, thiadiazolyl [e.g.,

1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl,

1,2,5-thiadiazolyl, etc.], etc.;

saturated 3 to 6-membered heteromonocyclic group

containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms

25     [e.g., thiazolidinyl, etc.];

unsaturated condensed heterocyclic group containing 1 to 2  
sulfur atoms and 1 to 3 nitrogen atoms [e.g.,

benzothiazolyl, benzothiadiazolyl, etc.];

unsaturated condensed heterocyclic group containing 1 to 2  
oxygen atoms [e.g. benzofuranyl, benzodioxolyl, etc.] and

30     the like.

Said "heterocyclic group" may be substituted with  
lower alkyl as exemplified above or oxo, in which  
preferable one is N-methylpiperazinyl, tetrazolyl,  
35     morpholinyl, pyrrolidinyl, N-methylpiperidyl,

- 32 -

N-methylhomopiperazinyl, 1H-tetrahydropyranyl, thienyl, pyridyl, piperidyl or oxopiperidyl.

Suitable acyl and acyl moiety in the terms  
"acyl(lower)alkylsulfinyl", "acyl(lower)alkylsulfonyl",  
5 "acyloxy", "acylamino", "acyl(lower)alkanoyloxy",  
"acyl(lower)alkoxyimino", "acyl(lower)alkylamino", "N-protected-acyl(lower)alkylamino", "N-acyl(lower)alkyl-N-lower alkylamino" and "acyl(lower)alkoxy" may be carboxy, esterified carboxy, carbamoyl, lower alkylcarbamoyl, lower  
10 alkanoyl, aroyl, a heterocycliccarbonyl and the like.

The esterified carboxy may be substituted or unsubstituted lower alkoxycarbonyl [e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, tert-butoxycarbonyl, hexyloxycarbonyl,  
15 2-iodoethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, dimethylaminopropoxycarbonyl, dimethylaminoethoxycarbonyl, etc.], substituted or unsubstituted aryloxycarbonyl [e.g. phenoxycarbonyl, 4-nitrophenoxycarbonyl, 2-naphthyloxycarbonyl, etc.], substituted or unsubstituted  
20 ar(lower)alkoxycarbonyl [e.g. benzyloxycarbonyl, phenethyloxycarbonyl, benzhydryloxycarbonyl, 4-nitrobenzyloxycarbonyl, 3-methoxy-4-nitrobenzyloxy-carbonyl, etc.], N-containing heterocyclicoxycarbonyl [e.g. N-methylpiperidyloxycarbonyl, etc.] and the like, in  
25 which preferable one is lower alkoxycarbonyl, N-methylpiperidyloxycarbonyl, dimethylaminopropoxycarbonyl or dimethylaminoethoxycarbonyl.

The lower alkylcarbamoyl may be mono or di(lower alkyl)carbamoyl such as methylcarbamoyl, ethylcarbamoyl,  
30 propylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl, N-methyl-N-ethylcarbamoyl or the like.

The lower alkanoyl may be substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkanoyl such as formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl,  
35 pivaloyl, hexanoyl, trifluoroacetyl or the like, in which

- 33 -

preferable one is formyl, acetyl or butyryl.

The aroyl may be benzoyl, naphthoyl, toluoyl, di(tert-butyl)benzoyl and the like, in which preferable one is benzoyl.

5 The heterocyclic moiety in the terms "a heterocyclic-carbonyl", "heterocyclicoxycarbonylamino" and "heterocyclicsulfonyl" may be one mentioned above as a heterocyclic group.

10 Preferred "a heterocycliccarbonyl" may be N-containing heterocycliccarbonyl.

The "N-containing heterocycliccarbonyl" may be one containing at least one nitrogen atom in heterocyclic group mentioned above, in which preferable one is N-(lower alkyl)piperazinylcarbonyl (e.g. N-methylpiperazinylcarbonyl, etc.), N-(lower alkyl)-homopiperazinylcarbonyl (e.g. N-methylhomopiperazinylcarbonyl, etc.), piperazinylcarbonyl, pyrrolidinylcarbonyl, piperidylcarbonyl, morpholinocarbonyl, lower alkylpiperidylcarbonyl (e.g. methylpiperidylcarbonyl, etc.) or oxopiperidylcarbonyl.

20 Suitable "substituted acyl" may be carbamoyl substituted with amino, a heterocyclic group [e.g. N-(lower alkyl)piperazinyl, pyridyl, etc.], lower alkylsulfonyl or arylsulfonyl, substituted lower alkylcarbamoyl [e.g. N-lower alkylamino-N-lower alkylcarbamoyl, pyridyl(lower)alkylcarbamoyl, morpholino(lower)alkylcarbamoyl, bis(hydroxy(lower)alkyl)carbamoyl, hydroxy(lower)alkylcarbamoyl, carbamoyl(lower)alkylcarbamoyl, lower alkylamino(lower)alkylcarbamoyl, N-lower alkyl-N-lower alkylcarbamoyl, etc.], substituted N-containing heterocycliccarbonyl [e.g. trifluoroacetyl-piperazinylcarbonyl, pyridylpiperazinylcarbonyl, hydroxypiperidylcarbonyl, dimethylaminopiperidylcarbonyl,

35

- 34 -

diethylaminopiperidylcarbonyl,  
carbamoylpyrrolidinylcarbonyl,  
dimethylaminopiperazinylcarbonyl, hydroxyethoxyethyl-  
piperazinylcarbonyl, pyrrolidinylcarbonylmethyl-  
5 piperazinylcarbonyl, etc.), N-protected-N-containing  
heterocycliccarbonyl [e.g. N-t-butoxycarbonylpiperidyl-  
carbonyl, N-t-butoxycarbonylpiperazinylcarbonyl, etc.],  
N-protected amino(lower)alkanoyl, amino(lower)alkanoyl,  
benzyloxybenzoyl, and the like.

10 "N-Protective group" in "protected amino" may be  
common N-protective group such as substituted or  
unsubstituted lower alkanoyl [e.g. formyl, acetyl,  
propionyl, trifluoroacetyl, etc.], phthaloyl, lower  
alkoxycarbonyl [e.g. tert-butoxycarbonyl, tert-  
15 amyloxycarbonyl, etc.], substituted or unsubstituted  
aralkyloxycarbonyl [e.g. benzyloxycarbonyl,  
p-nitrobenzyloxycarbonyl, etc.],  
9-fluorenylmethoxycarbonyl, substituted or unsubstituted  
arenesulfonyl [e.g. benzenesulfonyl, tosyl, etc.],  
20 nitrophenylsulfenyl, aralkyl [e.g. trityl, benzyl, etc.]  
or the like, in which preferable one is phthaloyl, tert-  
butoxycarbonyl or 9-fluorenylmethoxycarbonyl.

"N-protective group" in "N-protected guanidino" may  
be common N-protective group such as lower alkoxycarbonyl  
25 [e.g. tert-butoxycarbonyl, etc.] or the like.

Suitable "acid residue" may be halogen [e.g. fluoro,  
chloro, bromo, iodo], arenesulfonyloxy [e.g.  
benzenesulfonyloxy, tosyloxy, etc.], alkanesulfonyloxy  
[e.g. mesyloxy, ethanesulfonyloxy, etc.], and the like, in  
30 which preferable one is halogen.

Suitable "lower alkylsulfonyl" and lower  
alkylsulfonyl moiety in the term "lower alkylsulfonyloxy"  
may be (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl such as methylsulfonyl,  
ethylsulfonyl, propylsulfonyl or the like, in which  
35 preferable one is methylsulfonyl.



- 35 -

Suitable "lower alkylene" may be straight or branched C<sub>1</sub>-C<sub>6</sub> alkylene such as methylene, ethylene, propylene or the like, in which preferable one is methylene or ethylene.

5        Suitable "lower alkenylene" may be straight or branched C<sub>2</sub>-C<sub>6</sub> alkenylene such as ethenylene, propenylene or the like, in which preferable one is ethenylene.

10        The substituent(s) on aryl for R<sup>1</sup> may be plural and in such case the substituents may be the same or different.

Preferred "aryl" for R<sup>1</sup> may be phenyl or phenyl substituted with lower alkyl.

Preferred "cyclo(lower)alkyl" for R<sup>1</sup> may be cyclopentyl.

15        Preferred "a heterocyclic group" for R<sup>1</sup> may be pyridyl or thienyl.

20        Preferred compound (I) is one having aryl (more preferably phenyl or phenyl substituted with lower alkyl) which may be substituted with lower alkoxy optionally substituted with acylamino or acyl for R<sup>1</sup>, lower alkyl for R<sup>2</sup>, hydrogen, lower alkyl or lower alkoxy for R<sup>3</sup>, hydroxy, or lower alkoxy, lower alkylthio or lower alkyl, each of which may be substituted with hydroxy, aryl, substituted aryl, acyl, amino, lower alkylamino, acylamino, protected amino or a heterocyclic group for R<sup>4</sup>, hydrogen, lower alkyl, lower alkoxy or halogen for R<sup>5</sup>, NH for A,  $\text{-}\overset{\text{O}}{\underset{\text{||}}{\text{C}}}\text{-}$  for E,  $\text{-CH=CH-}$  for X, and CH for Y.

25        More preferred compound (I) is one having phenyl or tolyl, each of which is substituted with lower alkoxy substituted with N-(lower alkyl)piperazinylcarbonyl for R<sup>1</sup>, lower alkyl for R<sup>2</sup>, hydrogen, lower alkyl or lower alkoxy for R<sup>3</sup>, lower alkoxy substituted with amino for R<sup>4</sup>, hydrogen for R<sup>5</sup>, NH for A,  $\text{-}\overset{\text{O}}{\underset{\text{||}}{\text{C}}}\text{-}$  for E,  $\text{-CH=CH-}$  for X and CH for Y.

35        Most preferred compound (I) is one having tolyl

- 36 -

which is substituted with lower alkoxy substituted with N-(lower alkyl)piperazinylcarbonyl for R<sup>1</sup>, lower alkyl for R<sup>2</sup>, lower alkoxy for R<sup>3</sup>, lower alkoxy substituted with amino for R<sup>4</sup>, hydrogen for R<sup>5</sup>, NH for A,  $\overset{\text{O}}{\parallel}\text{C}-$  for E, -CH=CH- for X and CH for Y.

Suitable pharmaceutically acceptable salts of the object compound (I) are conventional non-toxic salts and include an acid addition salt such as an inorganic acid addition salt [e.g. hydrochloride, hydrobromide, sulfate, phosphate, etc.], an organic acid addition salt [e.g. formate, acetate, trifluoroacetate, maleate, tartrate, methanesulfonate, benzenesulfonate, toluenesulfonate, etc.], a metal salt such as an alkali metal salt [e.g. sodium salt, potassium salt, etc.] and an alkaline earth metal salt [e.g. calcium salt, magnesium salt, etc.] and the like.

The processes for preparing the object compound (I) are explained in detail in the following.

#### Process 1

The object compound (Ia) or its salt can be prepared by reacting a compound (II) or its salt with a compound (III) or its reactive derivative at the carboxy group or the sulfo group, or a salt thereof.

Suitable salts of the compounds (Ia) and (II) may be the same as those exemplified for the compound (I).

Suitable salts of the compound (III) and its reactive derivative at the carboxy group or the sulfo group may be base salts as exemplified for the compound (I).

Suitable reactive derivative at the carboxy group or the sulfo group of the compound (III) may include an acid halide, an acid anhydride containing intramolecular, intermolecular and a mixed ones, an activated amide, an activated ester, and the like. Suitable examples of the

- 37 -

reactive derivatives may be an acid chloride; an acid azide; a mixed acid anhydride with an acid such as substituted phosphoric acid [e.g. dialkylphosphoric acid, phenylphosphoric acid, diphenylphosphoric acid, dibenzylphosphoric acid, halogenated phosphoric acid, etc.], dialkylphosphorous acid, sulfurous acid, thiosulfuric acid, sulfuric acid, sulfonic acid [e.g. methanesulfonic acid, etc.], aliphatic carboxylic acid [e.g. acetic acid, propionic acid, butyric acid, isobutyric acid, pivalic acid, pentanoic acid, isopentanoic acid, 2-ethylbutyric acid, trichloroacetic acid, etc.] or aromatic carboxylic acid [e.g. benzoic acid, etc.]; a symmetrical acid anhydride; an activated amide with imidazole, 4-substituted imidazole, dimethylpyrazole, triazole or tetrazole; or an activated ester [e.g. cyanomethyl ester, methoxymethyl ester, dimethyliminomethyl  $[(CH_3)_2N^+=CH-]$  ester, vinyl ester, propargyl ester, p-nitrophenyl ester, 2,4-dinitrophenyl ester, trichlorophenyl ester, pentachlorophenyl ester, mesylphenyl ester, phenylazophenyl ester, phenyl thioester, p-nitrophenyl thioester, p-cresyl thioester, carboxymethyl thioester, pyranyl ester, pyridyl ester, piperidyl ester, 8-quinolyl thioester, etc.] or an ester with an N-hydroxy compound [e.g. N,N-dimethylhydroxylamine, 1-hydroxy-2-(1H)-pyridone, N-hydroxysuccinimide, N-hydroxyphthalimide, 1-hydroxy-1H-benzotriazole, etc.], and the like. These reactive derivatives can optionally be selected from them according to the kind of the compound (III) to be used.

The reaction is usually carried out in a conventional solvent such as water, alcohol [e.g. methanol, ethanol, etc.], acetone, dioxane, acetonitrile, chloroform, methylene chloride, ethylene chloride, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide, pyridine or any other organic solvent which does not adversely influence

- 38 -

the reaction. These conventional solvents may also be used in a mixture with water.

In this reaction, when the compound (III) is used in a free acid form or its salt form, the reaction is preferably carried out in the presence of a conventional condensing agent such as N,N'-dicyclohexylcarbodiimide; N-cyclohexyl-N'-morpholinoethylcarbodiimide; N-cyclohexyl-N'-(4-diethylaminocyclohexyl)carbodiimide; N,N'-diethylcarbodiimide, N,N'-diisopropylcarbodiimide; N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide; N,N'-carbonylbis-(2-methylimidazole); pentamethyleneketene-N-cyclohexylimine; diphenylketene-N-cyclohexylimine; ethoxyacetylene; 1-alkoxy-1-chloroethylene; trialkyl phosphite; ethyl polyphosphate; isopropyl polyphosphate; phosphorus oxychloride (phosphoryl chloride); phosphorus trichloride; diphenylphosphoryl azide; diphenyl chlorophosphate; diphenylphosphinic chloride; thionyl chloride; oxalyl chloride; lower alkyl haloformate [e.g. ethyl chloroformate, isopropyl chloroformate, etc.]; triphenylphosphine; 2-ethyl-7-hydroxybenzisoxazolium salt; 2-ethyl-5-(m-sulfophenyl)isoxazolium hydroxide intramolecular salt; 1-(p-chlorobenzenesulfonyloxy)-6-chloro-1H-benzotriazole; so-called Vilsmeier reagent prepared by the reaction of N,N-dimethylformamide with thionyl chloride, phosgene, trichloromethyl chloroformate, phosphorus oxychloride, etc.; or the like.

The reaction may also be carried out in the presence of an inorganic or organic base such as an alkali metal bicarbonate, tri(lower)alkylamine, pyridine, 4-dimethylaminopyridine, N-(lower)alkylmorpholine, N,N-di(lower)alkylbenzylamine, or the like.

The reaction temperature is not critical, and the reaction is usually carried out under cooling to heating.

- 39 -

Process 2

The object compound (I) or its salt can be prepared by reacting a compound (IV) or its salt with a compound (V) or its reactive derivative at the carboxy group or a salt thereof.

Suitable salts of the compounds (IV) and (V) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as Process 1, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 1.

Process 3

The object compound (Ic) or its salt can be prepared by subjecting a compound (Ib) or its salt to deesterification reaction.

Suitable salt of the compound (Ic) may be the same as those exemplified for the compound (I).

Suitable salt of the compound (Ib) may be an acid addition salt as exemplified for the compound (I).

The reaction is carried out in accordance with a conventional method such as hydrolysis, reduction or the like.

The hydrolysis is preferably carried out in the presence of a base or an acid including Lewis acid. Suitable base may include an inorganic base and an organic base such as an alkali metal [e.g. lithium, sodium, potassium, etc.], an alkaline earth metal [e.g. magnesium, calcium, etc.], the hydroxide or carbonate or bicarbonate thereof, trialkylamine [e.g. trimethylamine, triethylamine, etc.], picoline, 1,5-diazabicyclo[4.3.0]-non-5-ene, 1,4-diazabicyclo[2.2.2]octane, 1,8-diazabicyclo[5.4.0]undec-7-ene, or the like. Suitable

- 40 -

acid may include an organic acid [e.g. formic acid, acetic acid, propionic acid, trichloroacetic acid, trifluoroacetic acid, etc.], an inorganic acid [e.g. hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, etc.] and Lewis acid [e.g. boron tribromide, etc.].

The reaction is usually carried out in a solvent such as water, an alcohol [e.g. methanol, ethanol, etc.], xylene, diethylene glycol monomethyl ethyl, methylene chloride, tetrahydrofuran, a mixture thereof or any other solvent which does not adversely influence the reaction. A liquid base or acid can be also used as the solvent.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

The reduction can be applied preferably for elimination of the ester moiety such as 4-nitrobenzyl, 2-iodoethyl, 2,2,2-trichloroethyl, or the like. The reduction method applicable for the elimination reaction may include chemical reduction and catalytic reduction.

Suitable reducing agents to be used in chemical reduction are a combination of metal [e.g. tin, zinc, iron, etc.] or metallic compound [e.g. chromium chloride, chromium acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.].

Suitable catalysts to be used in catalytic reduction are conventional ones such as platinum catalysts [e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalyst [e.g. spongy palladium, palladium black, palladium oxide, palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalyst [e.g. reduced nickel, nickel oxide, Raney nickel, etc.], cobalt catalyst [e.g.

- 41 -

reduced cobalt, Raney cobalt, etc.], iron catalyst [e.g. reduced iron, Raney iron, etc.], copper catalyst [e.g. reduced copper, Raney copper, Ullman copper, etc.] or the like.

5           The reduction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, an alcohol [e.g. methanol, ethanol, propanol, etc.], N,N-dimethylformamide, or a mixture thereof. Additionally, in case that the  
10       above-mentioned acids to be used in chemical reduction are in liquid, they can also be used as a solvent. Further, a suitable solvent to be used in catalytic reduction may be the above-mentioned solvent, and other conventional solvent such as diethyl ether, dioxane, tetrahydrofuran,  
15       etc., or a mixture thereof.

          The reaction temperature of this reduction is not critical and the reaction is usually carried out under cooling to warming.

          In this reaction, in case that the compound (Ib)  
20       having lower alkyl substituted with esterified carboxy for  $R^2$  and/or acyloxy, lower alkoxy substituted with esterified carboxy, lower alkylthio substituted with esterified carboxy, or lower alkyl substituted with esterified carboxy for  $R^4$  is used as a starting compound,  
25       the compound (Ic) having lower alkyl substituted with carboxy for  $R^2$  and/or hydroxy, lower alkoxy substituted with carboxy, lower alkylthio substituted with carboxy, or lower alkyl substituted with carboxy for  $R^4$  may be obtained according to reaction condition. This case is  
30       included within the scope of this reaction.

#### Process 4

          The object compound (Ie) or its salt can be prepared by subjecting a compound (Id) or its salt to  
35       deesterification reaction.

- 42 -

Suitable salt of the compound (Id) may be an acid addition salt as exemplified for the compound (I).

Suitable salt of the compound (Ie) may be the same as those exemplified for the compound (I).

5 This reaction can be carried out in substantially the same manner as hydrolysis in Process 3, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in hydrolysis in Process 3.

10 In this reaction, in case that the compound (Id) having aryl, haloaryl, cyclo(lower)alkyl or a heterocyclic group, each of which is substituted with esterified carboxy; lower alkenyl substituted with esterified carboxy; lower alkyl substituted with esterified carboxy,  
15 esterified carboxy(lower)alkanoyloxy or esterified carboxy(lower)alkoxyimino; lower alkylthio substituted with esterified carboxy; alkoxy substituted with esterified carboxy-substituted aryl, esterified carboxy-substituted pyridyl, esterified carboxy(lower)alkylamino,  
20 N-protected-esterified carboxy(lower)alkylamino, N-esterified carboxy(lower)alkyl-N-lower alkylamino, esterified carboxy or esterified carboxy(lower)alkoxyimino; or lower alkenyloxy substituted with esterified carboxy for R<sup>1</sup> and/or lower alkyl  
25 substituted with esterified carboxy for R<sup>2</sup> is used as a starting compound, the compound (Ie) having aryl, haloaryl, cyclo(lower)alkyl or a heterocyclic group, each of which is substituted with carboxy; lower alkenyl substituted with carboxy; lower alkyl substituted with carboxy, carboxy(lower)alkanoyloxy or  
30 carboxy(lower)alkoxyimino; lower alkylthio substituted with carboxy; alkoxy substituted with carboxy-substituted aryl, carboxy-substituted pyridyl, carboxy(lower)-alkylamino, N-protected-carboxy(lower)alkylamino,  
35 N-carboxy(lower)alkyl-N-lower alkylamino, carboxy or



- 43 -

carboxy(lower)alkoxyimino; or lower alkenyloxy substituted with carboxy for  $R^1$  and/or lower alkyl substituted with carboxy for  $R^2$  may be obtained according to reaction condition. This case is included within the scope of this reaction.

#### Process 5

The object compound (Ig) or its salt can be prepared by subjecting a compound (If) or its salt to elimination reaction of the N-protective group.

Suitable salts of the compounds (If) and (Ig) may be acid addition salts as exemplified for the compound (I).

This reaction is carried out in accordance with a conventional method such as hydrolysis, reduction or the like.

The hydrolysis is preferably carried out in the presence of a base or an acid including Lewis acid.

Suitable base may include an inorganic base and an organic base such as an alkali metal [e.g. sodium, potassium, etc.], an alkaline earth metal [e.g. magnesium, calcium, etc.], the hydroxide or carbonate or bicarbonate thereof, hydrazine, alkylamine [e.g. methylamine, trimethylamine, triethylamine, etc.], picoline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, 1,8-diazabicyclo[5.4.0]undec-7-ene, or the like.

Suitable acid may include an organic acid [e.g. formic acid, acetic acid, propionic acid, trichloroacetic acid, trifluoroacetic acid, etc.], an inorganic acid [e.g. hydrochloric acid, hydrobromic acid, sulfuric acid, hydrogen chloride, hydrogen bromide, hydrogen fluoride, etc.] and an acid addition salt compound [e.g. pyridine hydrochloride, etc.].

The elimination using trihaloacetic acid [e.g. trichloroacetic acid, trifluoroacetic acid, etc.] or the

- 44 -

like is preferably carried out in the presence of cation trapping agents [e.g. anisole, phenol, etc.].

5 The reaction is usually carried out in a solvent such as water, an alcohol [e.g. methanol, ethanol, etc.], methylene chloride, chloroform, tetrachloromethane, tetrahydrofuran, a mixture thereof or any other solvent which does not adversely influence the reaction. A liquid base or acid can be also used as the solvent. The reaction temperature is not critical and the reaction is usually carried out under cooling to heating.

10 The reduction method applicable for the elimination reaction may include chemical reduction and catalytic reduction.

15 Suitable reducing agents to be used in chemical reduction are a combination of metal [e.g. tin, zinc, iron, etc.] or metallic compound [e.g. chromium chloride, chromium acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.].

20 Suitable catalysts to be used in catalytic reduction are conventional ones such as platinum catalysts [e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalysts [e.g. spongy palladium, palladium black, palladium oxide, palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalysts [e.g. reduced nickel, nickel oxide, Raney nickel, etc.], cobalt catalysts [e.g. reduced cobalt, Raney cobalt, etc.], iron catalysts [e.g. reduced iron, Raney iron, etc.], copper catalysts [e.g. reduced copper, Raney copper, Ullman copper, etc.] and the like.

30 In case that the N-protective group is benzyl, the reduction is preferably carried out in the presence of a

35

- 45 -

combination of palladium catalysts [e.g. palladium black, palladium on carbon, etc.] and formic acid or its salt [e.g. ammonium formate, etc.].

5 The reduction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, methanol, ethanol, propanol, N,N-dimethylformamide, or a mixture thereof. Additionally, in case that the above-mentioned acids to be used in chemical reduction are in liquid, they can also be used as a solvent. Further, a suitable solvent to be used in catalytic reduction may be the above-mentioned solvent, and other conventional solvent such as diethyl ether, dioxane, tetrahydrofuran, etc. or a mixture thereof.

10 The reaction temperature of this reduction is not critical and the reaction is usually carried out under cooling to heating.

15 In this reaction, in case that the compound (If) having aryl which is substituted with alkoxy substituted with protected amino, N-protected amino(lower)alkanoylamino, N-protected piperazinylcarbonyl or N-protected guanidino for  $R^1$  is used as a starting compound, the compound (Ig) having aryl which is substituted with alkoxy substituted with amino, amino(lower)alkanoylamino, piperazinylcarbonyl or guanidino for  $R^1$  may be obtained according to reaction condition. This case is included within the scope of this reaction.

#### Process 6

30 The object compound (Ih) or its salt can be prepared by reacting a compound (Ic) or its reactive derivative at the carboxy group or a salt thereof with an amine or its salt.

35 Suitable salt of amine may be an acid addition salt as exemplified for the compound (I).

- 46 -

Suitable salts of the compounds (Ih) and (Ic) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

5        Suitable "amine" may be ammonia, substituted or unsubstituted lower alkylamine, substituted or unsubstituted N-containing heterocyclic compound, a heterocyclic group substituted with amino and the like.

10        The substituted or unsubstituted lower alkylamine may be mono or di(lower)alkylamine (e.g. methylamine, ethylamine, propylamine, isopropylamine, butylamine, isobutylamine, pentylamine, hexylamine, dimethylamine, diethylamine, dipropylamine, dibutylamine, di-isopropylamine, dipentylamine, dihexylamine, etc.), pyridyl(lower)alkylamine, (e.g. pyridylmethylamine, etc.), 15        lower alkylamino(lower)alkylamine (e.g. N-dimethylaminoethylamine, N-dimethylaminopropylamine, N-diethylaminoethyl-N-methylamine, etc.), morpholino(lower)alkylamine (e.g. morpholinoethylamine, etc.) or the like.

20        The substituted or unsubstituted N-containing heterocyclic compound may be a heterocyclic group substituted with amino (e.g. aminopyridine, N-methyl-N'-aminopiperazine, etc.), saturated 5 or 6-membered N-, or N- and S-, or N- and O-containing heterocyclic compound 25        such as pyrrolidine, imidazolidine, piperidine, piperidone, piperazine, lower alkylaminopiperidine (e.g. dimethylaminopiperidine, etc.), N-(lower)alkylhomopiperazine (e.g. N-methylhomopiperazine, etc.), N-(lower)alkylpiperazine (e.g. N-methylpiperazine, 30        N-ethylpiperazine, etc.), morpholine, thiomorpholine, N-pyridylpiperazine, N-hydroxy(lower)alkoxy(lower)-alkylpiperazine (e.g. N-hydroxyethoxyethylpiperazine, etc.), N-pyrrolidinylcarbonyl(lower)alkylpiperazine (e.g. N-pyrrolidinylcarbonylmethylpiperazine, etc.), or the 35        like, in which preferable one is N-methylpiperazine.

- 47 -

This reaction can be carried out in substantially the same manner as Process 1, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 1.

#### Process 7

The object compound (Ii) or its salt can be prepared by reacting a compound (Ie) or its reactive derivative at the carboxy group or a salt thereof with an amine or its salt.

Suitable salt of amine may be an acid addition salt as exemplified for the compound (I).

Suitable salts of the compounds (Ii) and (Ie) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manners as Processes 1 and 6, and therefore the reaction mode and reaction condition (e.g. amine, solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Processes 1 and 6.

#### Process 8

The object compound (Ik) or its salt can be prepared by subjecting a compound (Ij) or its salt to debenzylation reaction.

Suitable salts of the compounds (Ij) and (Ik) may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as hydrolysis using an acid or catalytic reduction in Process 5, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in hydrolysis using an acid or catalytic reduction in Process 5.

- 48 -

In this catalytic reduction, in case that the compound (Ij) having nitro for  $R^3$  is used as a starting compound, the compound (Ik) having amino for  $R^3$  may be obtained according to reaction condition. This case is included within the scope of this reaction.

#### Process 9

The object compound (Il) or its salt can be prepared by reacting a compound (Ika) or its salt with a compound (VI) or its salt.

Suitable salts of the compounds (Ika), (Il) and (VI) may be the same as those exemplified for the compound (I).

When the compound (VI) having halogen for  $Z^1$  is used in this reaction, the reaction is preferably carried out in the presence of a base such as an alkali metal (e.g. sodium, potassium, etc.), an alkaline earth metal (e.g. magnesium, calcium, etc.), the hydride or hydroxide or carbonate or bicarbonate thereof.

When the compound (VI) having hydroxy for  $Z^1$  is used in this reaction, the reaction is preferably carried out in the presence of diethyl azodicarboxylate and triphenylphosphine.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, dioxane, alcohol (e.g. methanol, ethanol, etc.), acetonitrile, tetrahydrofuran, acetic acid, N,N-dimethylformamide, or a mixture thereof.

The reaction temperature is not critical and the reaction can be carried out under cooling to heating.

#### Process 10

The object compound (Im) or its salt can be prepared by reacting a compound (Iga) or its salt with an acylating agent.

Suitable salts of the compounds (Iga) and (Im) may be

- 49 -

the same as those exemplified for the compound (I).

The acylating agent may include an organic acid represented by the formula :  $R^{11}-OH$ , in which  $R^{11}$  is acyl or substituted acyl as illustrated above, or its reactive derivative.

The suitable reactive derivative of organic acid may be a conventional one such as an acid halide [e.g. acid chloride, acid bromide, etc.], an acid azide, an acid anhydride containing intramolecular and intermolecular ones, an activated amide, an activated ester or the like.

When free acid is used as an acylating agent, the acylation reaction may preferably be conducted in the presence of a conventional condensing agent such as  $N,N'$ -dicyclohexylcarbodiimide or the like.

The reaction is usually carried out in a conventional solvent such as water, pyridine, acetone, dioxane, chloroform, methylene chloride, acetonitrile, ethylene chloride, tetrahydrofuran, ethyl acetate,  $N,N$ -dimethylformamide, pyridine or any other organic solvent which does not adversely influence the reaction, or a mixture thereof.

The reaction is also preferably carried out in the presence of a conventional base such as triethylamine, pyridine, sodium hydroxide or the like.

The reaction temperature is not critical, and the reaction can be carried out under cooling to heating.

#### Process 11

The object compound (In) or its salt can be prepared by reacting a compound (Igb) or its salt with lower alkanal or  $N$ -protected amino(lower)alkanal in the presence of a reducing agent.

Suitable salts of the compounds (Igb) and (In) may be the same as those exemplified for the compound (I).

Suitable lower alkanal may be  $C_1$ - $C_6$  alkanal such as

- 50 -

formaldehyde, ethanal, propanal or the like, in which preferable one is formaldehyde.

Suitable N-protected amino(lower)alkanal may be N-protected amino(C<sub>1</sub>-C<sub>6</sub>)alkanal such as phthalimidopropanal or the like.

Suitable reducing agent may be diborane, borane-organic amine complex [e.g. borane-pyridine complex, etc.], alkali metal cyanoborohydride [e.g. sodium cyanoborohydride, lithium cyanoborohydride, etc.], sodium borohydride and the like.

The reaction is preferably carried out in the presence of molecular sieves.

The reaction is usually carried out in a conventional solvent such as water, an alcohol [e.g. methanol, ethanol, etc.], dioxane, tetrahydrofuran, a mixture thereof or any other organic solvent which does not adversely influence the reaction.

The reaction may also be carried out in an acidic condition [e.g. presence of acetic acid, sulfuric acid, etc.] and the reaction temperature is not critical, and the reaction is usually carried out under cooling to warming.

#### Process 12

The object compound (Ip) or its salt can be prepared by subjecting a compound (Io) or its salt to reduction.

Suitable salts of the compounds (Io) and (Ip) may be the same as those exemplified for the compound (I).

The reduction may include chemical reduction and catalytic reduction, which are carried out in a conventional manner.

Suitable reducing agents to be used in chemical reduction are a metal [e.g. tin, zinc, iron, nickel, etc.], a combination of such metal and/or metallic compound [e.g. nickel chloride, chromium chloride,



- 51 -

chromium acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.], a combination of such metal and/or metallic compound and base [e.g. ammonia, ammonium chloride, sodium hydroxide, etc.], a metal hydride compound such as aluminum hydride compound [e.g. lithium aluminum hydride, sodium aluminum hydride, aluminum hydride, lithium trimethoxyaluminum hydride, lithium tri-t-butoxyaluminum hydride, etc.], borohydride compound [e.g. sodium borohydride, lithium borohydride, sodium cyanoborohydride, tetramethylammonium borohydride, borane, diborane, etc.], a phosphorus compound [e.g. phosphorus trichloride, phosphorus tribromide, triphenylphosphine, triethylphosphine, etc.] and the like.

Suitable catalysts to be used in catalytic reduction are conventional ones such as platinum catalysts [e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalyst [e.g. spongy palladium, palladium black, palladium oxide, palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalyst [e.g. reduced nickel, nickel oxide, Raney nickel, etc.], cobalt catalyst [e.g. reduced cobalt, Raney cobalt, etc.], iron catalyst [e.g. reduced iron, Raney iron, etc.], copper catalyst [e.g. reduced copper, Raney copper, Ullman copper, etc.], or the like.

The reduction is usually carried out in a solvent. A suitable solvent to be used may be water, and alcohol [e.g. methanol, ethanol, propanol, etc.], acetonitrile or any other conventional organic solvent such as diethyl ether, dioxane, tetrahydrofuran, etc. or a mixture thereof.

The reaction temperature is not critical, and the

- 52 -

reaction is preferably carried out under cooling to heating.

### Process 13

5       The object compound (Ir) or its salt can be prepared by subjecting a compound (Iq) or its salt to deacylation reaction.

Suitable salts of the compounds (Iq) and (Ir) may be the same as those exemplified for the compound (I).

10       This reaction can be carried out in substantially the same manner as Process 3, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 3.

### Process 14

15       The object compound (Is) or its salt can be prepared by reacting a compound (VII) or its salt with a compound (VIII) or its salt.

20       Suitable salts of the compounds (Is), (VII) and (VIII) may be the same as those exemplified for the compound (I).

25       This reaction can be carried out in substantially the same manner as Process 9, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 9.

### Process 15

30       The object compound (Iu) or its salt can be prepared by reacting a compound (It) or its salt with an oxidizing agent.

Suitable salts of the compounds (It) or (Iu) may be the same as those exemplified for the compound (I).

35       The suitable oxidizing agent may be hydrogen

- 53 -

peroxide, Jones reagent, peracid [e.g. peracetic acid, perbenzoic acid, m-chloroperbenzoic acid, etc.], chromic acid, potassium permanganate, alkali metal periodate [e.g. sodium periodate, etc.] and the like.

5           This reaction is usually carried out in a solvent which does not adversely influence the reaction such as acetic acid, dichloromethane, acetone, ethyl acetate, chloroform, water, an alcohol [e.g. methanol, ethanol, etc.], a mixture thereof or the like.

10           The reaction temperature is not critical, and the reaction is usually carried out under cooling to warming.

#### Process 16

15           The object compound (Iw) or its salt can be prepared by subjecting a compound (Iv) or its salt to catalytic reduction.

Suitable salts of the compounds (Iv) and (Iw) may be the same as those exemplified for the compound (I).

20           This reaction can be carried out in substantially the same manner as catalytic reduction in Process 5, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in catalytic reduction in Process 5.

25

#### Process 17

The object compound (Iy) or its salt can be prepared

(to be continued to the next page)

30

35

- 54 -

by subjecting a compound (Ix) or its salt to debenzylation reaction.

Suitable salts of the compounds (Ix) and (Iy) may be the same as those exemplified for the compound (I).

5 This reaction can be carried out in substantially the same manner as Process 8, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 8.

#### 10 Process 18

The object compound (Iz) or its salt can be prepared by reacting a compound (Iy) or its salt with a compound (IX) or its salt.

15 Suitable salts of the compounds (Iy), (Iz) and (IX) may be the same as those exemplified for the compound (I).

This reaction can be carried out in substantially the same manner as Process 9, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 9.

#### 20 Process 19

The object compound (I-2) or its salt can be prepared by subjecting a compound (I-1) or its salt to elimination reaction of the hydroxy protective group.

25 Suitable salts of the compounds (I-1) and (I-2) may be the same as those exemplified for the compound (I).

Suitable hydroxy protective group may be benzyloxy, acyloxy, substituted acyloxy or the like.

30 This reaction can be carried out in substantially the same manner as Processes 8 and 13, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Processes 8 and 13.

35

- 55 -

Process 20

The object compound (I-3) or its salt can be prepared by reacting a compound (I-2) or its salt with a compound (X) or its salt.

5        Suitable salts of the compounds (I-2), (I-3) and (X) may be the same as those exemplified for the compound (I).

10        This reaction can be carried out in substantially the same manner as Process 9, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 9.

Process 21

15        The object compound (I-4) or its salt can be prepared by subjecting a compound (I-3a) or its salt to deesterification reaction.

      Suitable salts of the compounds (I-3a) and (I-4) may be the same as those exemplified for the compound (I).

20        This reaction can be carried out in substantially the same manner as Process 3, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 3.

Process 22

25        The object compound (I-6) or its salt can be prepared by reacting a compound (I-5) or its salt with an alkyne compound in the presence of a palladium compound and a copper compound.

30        Suitable salts of the compounds (I-5) and (I-6) may be the same as those exemplified for the compound (I).

35        Suitable alkyne compound may be lower alkyne optionally substituted with hydroxy, amino, protected amino, lower alkylsulfonyl, arylsulfonyl or the like, in which preferable one is 3-butyne-1-ol.

- 56 -

Suitable palladium compound may be bis(triphenylphosphine)palladium(II) chloride, or the like.

5      Suitable copper compound may be copper(I) iodide, or the like.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as tetrahydrofuran, dioxane, ethylamine, or a mixture thereof.

10      The reaction temperature is not critical and the reaction is usually carried out under warming or heating.

#### Process 23

15      The object compound (I-8a) or its salt can be prepared by reacting a compound (I-7) or its salt with a compound (XI).

Suitable salts of the compounds (I-7) and (I-8a) may be the same as those exemplified for the compound (I).

20      The reaction is preferably carried out in the presence of a base such as trialkylamine (e.g. trimethylamine, triethylamine, etc.) or the like.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as tetrahydrofuran, methylene chloride or the like.

25      The reaction temperature is not critical and the reaction is usually carried out under cooling to heating.

#### Process 24

30      The object compound (I-9) or its salt can be prepared by reacting a compound (I-8) or its salt with alkali metal phthalimide.

Suitable salts of the compounds (I-8) and (I-9) may be the same as those exemplified for the compound (I).

35      The reaction is carried out in a conventional solvent which does not adversely influence the reaction such as

- 57 -

dimethyl sulfoxide, N,N-dimethylformamide, or the like.

The reaction temperature is not critical and the reaction is usually carried out under cooling to heating.

5     Process 25

The object compound (I-10) or its salt can be prepared by reacting a compound (I-6) or its salt with a reducing agent.

10     Suitable salts of the compounds (I-6) and (I-10) may be the same as those exemplified for the compound (I).

Suitable reducing agent may be a combination of nickel chloride and sodium borohydride, and the like.

15     The reaction is carried out in a conventional solvent which does not adversely influence the reaction such as an alcohol (e.g. methanol, ethanol, etc.), tetrahydrofuran, or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

20     Process 26

The object compound (I-11) or its salt can be prepared by reacting a compound (II) or its salt with a compound (XII) or its salt.

25     Suitable salts of the compounds (I-11), (II) and (XII) may be the same as those exemplified for the compound (I).

30     This reaction can be carried out in substantially the same manner as Process 11, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 11.

Process 27

35     The object compound (I-13) or its salt can be prepared by reacting a compound (I-12) or its salt with a

- 58 -

compound (XIII) in the presence of a base.

Suitable salts of the compounds (I-12) and (I-13) may be the same as those exemplified for the compound (I).

5      Suitable base may be an alkali metal (e.g. sodium, potassium, etc.), an alkali metal hydride (e.g. sodium hydride), and the like.

10      The reaction is carried out in a solvent such as N,N-dimethylformamide, tetrahydrofuran, dioxane, a mixture thereof or any other solvent which does not adversely influence the reaction.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

#### Process 28

15      The object compound (I-15) or its salt can be prepared by reacting a compound (I-14) or its salt with an acylating agent.

Suitable salts of the compounds (I-14) and (I-15) may be the same as those exemplified for the compound (I).

20      This reaction can be carried out in substantially the same manner as Process 10, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 10.

25

#### Process 29

The object compound (I-17) or its salt can be prepared by reacting a compound (I-16a) or its salt with a reducing agent.

30      Suitable salts of the compounds (I-16a) and (I-17) may be the same as those exemplified for the compound (I).

Suitable reducing agent may be alkali metal borohydride (e.g. sodium borohydride, etc.), and the like.

35      The reaction is carried out in a solvent such as an alcohol (e.g. methanol, ethanol, etc.), tetrahydrofuran,



- 59 -

or the like.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

5     Process 30

The object compound (I-18) or its salt can be prepared by reacting a compound (I-16) or its salt with an amine compound or its salt in the presence of a reducing agent.

10     Suitable salts of the compounds (I-16) and (I-18) may be the same as those exemplified for the compound (I).

Suitable amine compound may be ammonia, N-lower alkylpiperazine, and the like.

15     Suitable salt of amine compound may be an acid addition salt (e.g. acetate, hydrochloride, etc.), and the like.

20     This reaction can be carried out in substantially the same manner as Process 11, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 11.

Process 31

25     The object compound (I-20) or its salt can be prepared by reacting a compound (I-19) or its reactive derivative at the carboxy group or a salt thereof with lower alkylamino(lower)alkanol.

30     Suitable salts of the compounds (I-20) and (I-19) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

Suitable lower alkylamino(lower)alkanol may be dimethylaminoethanol, and the like.

35     This reaction can be carried out in substantially the same manner as Process 1, and therefore the reaction mode and reaction condition (e.g. solvent, reaction

- 60 -

temperature, etc.) of this reaction are to be referred to those as explained in Process 1.

### Process 32

5       The object compound (I-22) or its salt can be prepared by reacting a compound (I-21) or its salt with a reducing agent.

      Suitable salts of the compounds (I-21) and (I-22) may be the same as those exemplified for the compound (I).

10       Suitable reducing agent may be diborane, lithium aluminum hydride and the like.

      The reaction is usually carried out in a solvent which does not adversely influence the reaction such as diethyl ether, tetrahydrofuran or the like.

15       The reaction temperature is not critical and the reaction can be carried out under cooling to heating.

### Process 33

20       The object compound (I-23) or its salt can be prepared by subjecting a compound (I-22) or its salt to oxidation reaction.

      Suitable salts of the compounds (I-22) and (I-23) may be the same as those exemplified for the compound (I).

25       Suitable oxidizing agent used in this reaction may be manganese dioxide, dimethyl sulfoxide, a mixture of dimethyl sulfoxide and oxalyl chloride and the like.

      The reaction is usually carried out in a conventional solvent such as pentane, hexane, benzene, diethyl ether, dimethoxyethane, acetone, chloroform, dichloromethane or  
30       any other solvent which does not adversely influence the reaction.

      Additionally in case that the above-mentioned oxidizing agent is liquid, it can be used as a solvent.

35       In this reaction, in case that dimethyl sulfoxide or a mixture of dimethyl sulfoxide and oxalyl chloride is

- 61 -

used as an oxidizing agent, the reaction is preferably carried out in the presence of alkali metal iodide (e.g. sodium iodide, etc.) and alkali metal carbonate (e.g. sodium carbonate) or tri(lower)alkylamine (e.g. triethylamine, etc.).

The reaction temperature is not critical, and the reaction is usually carried out under cooling to heating.

#### Process 34

The object compound (I-25) or its salt can be prepared by reacting a compound (I-24) or its salt with an azide compound.

Suitable salts of the compounds (I-24) and (I-25) may be the same as those exemplified for the compound (I).

Suitable azide compound may be sodium azide, trimethyltin azide and the like.

The reaction is usually carried out in a solvent which does not adversely influence the reaction such as dioxane, an aromatic hydrocarbon (e.g. benzene, toluene, xylene) or the like.

The reaction temperature is not critical and the reaction is usually carried out under warming to heating.

#### Process 35

The object compound (I-27) or its salt can be prepared by reacting a compound (I-26) or its salt with an isourea compound.

Suitable salts of the compounds (I-26) and (I-27) may be the same as those exemplified for the compound (I).

Suitable isourea compound may be O-alkylisourea (e.g. O-methylisourea, etc.) and the like.

The reaction is usually carried out in a solvent which does not adversely influence the reaction such as an alcohol (e.g. methanol, ethanol, etc.) or the like.

The reaction temperature is not critical and the

- 62 -

reaction is usually carried out under warming to heating.

#### Process 36

5       The object compound (I-29) or its salt can be prepared by subjecting a compound (I-28) or its salt to elimination reaction of the N-protective group.

Suitable salts of the compounds (I-28) and (I-29) may be the same as those exemplified for the compound (I).

10       This reaction can be carried out in substantially the same manner as Process 5, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 5.

#### 15       Process 37

The object compound (I-31) or its salt can be prepared by reacting a compound (I-30) or its salt with N-lower alkylpiperazine, dimethylaminopiperidine, ammonia or N,N-dimethylformamide.

20       Suitable salts of the compounds (I-30) and (I-31) may be the same as those exemplified for the compound (I).

The reaction is usually carried out in a solvent which does not adversely influence the reaction such as N,N-dimethylformamide, dioxane or the like.

25       The reaction temperature is not critical and the reaction is usually carried out under warming to heating.

#### Process 38

30       The object compound (I-3a) or its salt can be prepared by reacting a compound (I-4) or its reactive derivative at the carboxy group or a salt thereof with a hydroxy compound or a diazo compound.

35       Suitable salts of the compounds (I-3a) and (I-4) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

- 63 -

Suitable reactive derivative at the carboxy group (I-4) may be acid halide (e.g. acid chloride, acid bromide, etc.) and the like.

5      Suitable hydroxy compound may be an alcohol (e.g. methanol, ethanol, etc.), phenol, naphthol and the like.

Suitable diazo compound may be methyldiazomethane, trimethylsilyldiazomethane and the like.

10      The reaction is usually carried out in a conventional solvent such as diethyl ether, tetrahydrofuran, dioxane, methylene chloride, or any other organic solvent which does not adversely influence the reaction.

Additionally, in case that the above-mentioned hydroxy compound is in liquid, it can also be used as a solvent.

15      The reaction temperature is not critical and the reaction is usually carried out under cooling to heating.

#### Process 39

20      The object compound (I-32) or its salt can be prepared by reacting a compound (I-4) or its reactive derivative at the carboxy group or a salt thereof with an amine.

25      Suitable salts of the compounds (I-32) and (I-4) and its reactive derivative at the carboxy group may be the same as those exemplified for the compound (I).

Suitable amine may be ammonia, lower alkylamine (e.g. methylamine, dimethylamine, etc.) and the like.

30      This reaction can be carried out in substantially the same manner as Process 6, and therefore the reaction mode and reaction condition (e.g. solvent, reaction temperature, etc.) of this reaction are to be referred to those as explained in Process 6.

35      The compounds obtained by the above processes can be isolated and purified by a conventional method such as

- 64 -

pulverization, recrystallization, column chromatography, reprecipitation, or the like.

5 It is to be noted that the compound (I) and the other compounds may include one or more stereoisomer(s) such as optical isomer(s) or geometrical isomer(s) due to asymmetric carbon atom(s) and double bond(s), and all of such isomers and mixture thereof are included within the scope of this invention.

10 Additionally, it is to be noted that any hydrate of the compound (I) is also included within the scope of this invention.

15 The object compound (I) and pharmaceutically acceptable salts thereof possess activities as vasopressin antagonistic activity, vasodilating activity, hypotensive activity, activity for inhibiting saccharide release in liver, activity for inhibiting growth of mesangium cells, water diuretic activity, platelet agglutination inhibitory activity, oxytocin antagonistic activity and the like, and are useful for the treatment and/or prevention of  
20 hypertension, heart failure, renal insufficiency, edema, ascites, vasopressin parasecretion syndrome, hepato-cirrhosis, hyponatremia, hypokalemia, diabetic, circulation disorder, cerebrovascular disease (e.g. cerebral edema, cerebral infarction, etc.), Meniere's  
25 syndrome (e.g. Meniere's disease, et.), motion sickness and the like in human beings and animals.

30 In order to illustrate the usefulness of the object compound (I), the pharmacological data of the compound (I) are shown in the following.

#### Test 1

35 Vasopressin 1 (V1) receptor binding

- 65 -

## (i) Test Method :

Blood was obtained by venipuncture from normal subjects. Platelet-rich plasma (PRP) was prepared by centrifugation of whole blood at 200 xg for 10 minutes. PRP was centrifuged at 45,000 xg for 30 minutes. The remaining pellet was resuspended in 10 volume of ice cold 100 mM Tris-HCl (pH 7.4) buffer (containing 5 mM MgCl<sub>2</sub>, 0.1% bovine serum albumin and 1 mM EGTA), and centrifuged at 45,000 xg for 30 minutes again. The final pellet was resuspended in 100 mM Tris-HCl buffer. The resulting membrane preparation was used immediately for the binding assay.

Competition assays were conducted at equilibrium (15 minutes at 30°C) by using 1.5 nM <sup>3</sup>H-vasopressin (40-87 Ci/mmol; New England Nuclear) in 100 mM Tris-HCl (pH 7.4) buffer. Nonspecific binding was determined by using 1 μM vasopressin. After incubation, reaction was terminated by adding 5 ml of ice-cold 100 mM Tris-HCl (pH 7.4) buffer, and then filtered rapidly through Whatman glass filter (GF/C). The filter was washed twice with the same buffer. The glass filter was mixed with liquid scintillation cocktail, and radioactivity was counted in a liquid scintillation counter. Competition activity of the test compound was represented by IC<sub>50</sub> values.

## (ii) Test Result :

Test Compound (Example No.)	IC <sub>50</sub> (nM)
5-2)	51
16	14
17-20)	31

- 66 -

Test 2

## Vasopressin 2 (V2) receptor binding

## 5 (i) Test Method :

For binding assays, the receptor cDNA was permanently expressed in Chinese hamster ovary (CHO) cells. CHO cells were transfected with a vector directing expression of the cDNA for the human V2 receptor and the clonal cell lines expressing human V2 receptor was established essentially as described previously (Nakajima, Y., et. al. J. Biol. Chem., 1992, 267, 2437).

10 DNA-transfected cells were harvested and homogenized in ice cold 250 mM sucrose buffer containing 25 mM Tris-HCl (pH 7.4), 10 mM MgCl<sub>2</sub>, 1 mM EDTA and 5 µg/ml p-amidinophenylmethylsulfonyl fluoride (A-PMSF). The homogenate was centrifuged at 500 xg for 10 minutes. The supernatant was centrifuged at 100,000 xg for 1 hour. The final pellet was suspended in 25 mM Tris-HCl (pH 7.4) buffer (containing 10 mM MgCl<sub>2</sub>, 1 mM EDTA and 5 µg/ml A-PMSF), and stored in small aliquots at -80°C.

20 Competition assays were conducted at equilibrium (2 hours at 22°C) by using 0.5 nM <sup>3</sup>H-vasopressin (40-87 Ci/mmol, New England Nuclear) in 100 mM Tris-HCl (pH 7.4) buffer (containing 5 mM MgCl<sub>2</sub>, 5 µg/ml A-PMSF, 4 µg/ml leupeptin, 40 µg/ml bacitracin, 20 µg/ml chymostatin and 0.1% bovine serum albumin). Nonspecific binding was determined by using 1 µM vasopressin. After incubation, reaction mixture was rapidly filtered through Whatman glass filter (GF/C). The filter was washed twice with the same buffer. The radioactivity was counted in a liquid scintillation counter. Competition activity of the test compound was represented by IC<sub>50</sub> values.

35



- 67 -

(ii) Test Result :

5

Test Compound (Example No.)	IC <sub>50</sub> (nM)
5-2)	1300
16	.1400
17-20)	1300

10

15

20

For therapeutic purpose, the compound (I) of the present invention can be used in a form of pharmaceutical preparation containing one of said compounds, as an active ingredient, in admixture with a pharmaceutically acceptable carrier such as an organic or inorganic solid, semi-solid or liquid excipient suitable for oral, parenteral or external(topical) administration. The pharmaceutical preparations may be capsules, tablets, dragees, granules, suppositories, solution, lotion, suspension, emulsion, ointment, gel, or the like. If desires, there may be included in these preparations, auxiliary substances, stabilizing agents, wetting or emulsifying agents, buffers and other commonly used additives.

25

30

While the dosage of the compound (I) will vary depending upon the age and condition of the patient, an average single dose of about 0.1 mg, 1 mg, 10 mg, 50 mg, 100 mg, 250 mg, 500 mg and 1000 mg of the compound (I) may be effective for treating the above-mentioned diseases. In general, amounts between 0.1 mg/body and about 1,000 mg/body may be administered per day.

The following Preparations and Examples are given for the purpose of illustrating this invention.

35

- 68 -

Preparation 1

To a solution of [N-methyl-N-(4-nitrobenzoyl)]-2-hydroxyaniline (1.2 g) in N,N-dimethylformamide (30 ml) was added potassium carbonate (1.22 g), ethyl 6-bromohexanoate (1.03 g) and sodium iodide (catalytic amount) at 60°C. The reaction mixture was stirred at same temperature for 8 hours. The reaction mixture was cooled in an ice bath and quenched with 1N hydrochloric acid (10 ml) and water (30 ml). The mixture was extracted with ethyl acetate. The organic phase was washed with water and brine. The organic solution was dried over magnesium sulfate. The solvent was removed by evaporation to give 2-(5-ethoxycarbonylpent-1-yloxy)-[N-methyl-N-(4-nitrobenzoyl)]aniline (1.7 g).

NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7.5Hz), 1.45-1.58 (2H, m), 1.67-1.76 (2H, m), 1.79-1.88 (2H, m), 2.34 (2H, t, J=7.5Hz), 3.38 (3H, s), 3.84-4.00 (2H, m), 4.13 (2H, t), 6.72-6.82 (2H, m), 7.01 (1H, d, J=7Hz), 7.17 (1H, t, J=7Hz), 7.45 (2H, d, J=8.5Hz), 7.98 (2H, d, J=8.5Hz)

Preparation 2

A solution of 3-methoxy-4-nitro-N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]-N-methylbenzamide (7.6 g) in ethanol (76 ml) was treated with 1N sodium hydroxide solution (33 ml) at ambient temperature and the mixture was stirred at the same temperature for 6 hours. The reaction was quenched by the dropwise addition of 1N hydrochloric acid (35 ml). The mixture was concentrated and the residue was dissolved in a mixture of ethyl acetate and 1N hydrochloric acid. The extracted organic layer was washed with brine and dried over magnesium sulfate. The suspension was filtered and the solvent was removed by evaporation to give 3-methoxy-4-nitro-N-[2-(5-carboxypent-1-yloxy)-4-methylphenyl]-N-

- 69 -

methylbenzamide (7.1 g) as an oil.

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.63 (2H, m), 1.66-1.91 (4H, m), 2.28 (3H, s), 2.41 (2H, t, J=7Hz), 3.34 (3H, s), 3.78 (3H, s), 3.81-3.98 (2H, m), 6.58-6.67 (2H, m), 6.89 (1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.09 (1H, s), 7.61 (1H, d, J=8Hz)

### Preparation 3

3-Methoxy-4-nitro-N-[2-(5-carboxypent-1-yloxy)-4-methylphenyl]-N-methylbenzamide (5.2 g), 1-methylpiperazine (1.45 g) and 1-hydroxybenzotriazole (1.96 g) were dissolved in N,N-dimethylformamide (50 ml) and the solution was cooled in an ice bath. To the mixture was added N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride (2.78 g) and the solution was stirred at the same temperature for 30 minutes. The reaction mixture was allowed to warm to ambient temperature and stirring was continued for additional 20 hours. The reaction mixture was diluted with ethyl acetate and the solution was washed successively with saturated sodium hydrogen carbonate and brine, and dried over sodium sulfate. The sodium sulfate was removed and the solvent was removed by evaporation to give oil. The crude material was subjected to a silica gel column chromatography (SiO<sub>2</sub>; 120 g, 2% methanol in chloroform) to give 3-methoxy-4-nitro-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yloxy]phenyl]benzamide (6.2 g).

NMR (CDCl<sub>3</sub>, δ) : 1.43-1.60 (2H, m), 1.60-1.92 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.25-2.47 (6H, m), 3.34 (3H, s), 3.44-3.54 (2H, m), 3.58-3.70 (2H, m), 3.78 (3H, s), 3.82-4.03 (2H, m), 6.56-6.66 (2H, m), 6.86 (1H, d, J=8Hz), 6.94 (1H, d, J=8Hz), 7.07 (1H, s), 7.61 (1H, d, J=8Hz)

- 70 -

Preparation 4

A mixture of 3-methoxy-4-nitro-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yloxy]phenyl]benzamide (6.2 g) and iron powder (3.43 g) in a mixture of ethanol (65 ml) and ethyl acetate (6 ml) was refluxed for 2 hours. After being cooled to ambient temperature, the solution was filtered through a bed of Celite and the filtrate was evaporated in vacuo. The residue was diluted with ethyl acetate and the solution was washed with saturated sodium hydrogen carbonate and brine, and dried over magnesium sulfate. The solvent was evaporated in vacuo to give 4-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yloxy]phenyl]benzamide (4.7 g).

NMR (CDCl<sub>3</sub>, δ) : 1.43-1.58 (2H, m), 1.61-1.91 (4H, m), 2.26 (3H, s), 2.30 (3H, s), 2.23-2.44 (6H, m), 3.29 (3H, s), 3.41-3.53 (2H, m), 3.61 (3H, s), 3.57-3.68 (2H, m), 3.75-4.03 (4H, m), 6.36-6.46 (1H, m), 6.53-6.67 (2H, m), 6.76-6.89 (3H, m)

Preparation 5

The following compounds were obtained according to a similar manner to that of Preparation 4.

- 1) 4-Amino-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7.5Hz), 1.41-1.54 (2H, m), 1.62-1.73 (2H, m), 1.75-1.84 (2H, m), 2.32 (2H, t, J=7.5Hz), 3.30 (3H, s), 3.84 (2H, br), 3.90 (2H, br), 4.13 (2H, t), 6.38 (2H, d, J=8.5Hz), 6.79 (2H, d, J=8.5Hz), 6.99 (2H, s), 7.09-7.18 (3H, m)

- 2) 4-Amino-3-methoxy-N-methyl-N-[2-[5-(4-

- 71 -

dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.29-1.95 (10H, m), 2.23-2.43 (12H, m), 2.57 (1H, m), 3.01 (1H, m), 3.31 (3H, s), 3.62 (3H, s), 3.73-4.03 (5H, m), 4.63 (1H, m), 6.42 (1H, d, J=9Hz), 6.54-6.67 (2H, m), 6.77-6.89 (3H, m)

5

3) 4-Amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

10

MASS (m/z) : 399 (M+1)

4) 4-Amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]phenylbenzamide

15

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.27 (3H, s), 2.32 (3H, s), 2.35-2.55 (4H, m), 3.31 (3H, s), 3.38-3.54 (2H, m), 3.66-3.87 (4H, m), 4.90-5.10 (2H, m), 6.38 (2H, d, J=8Hz), 6.62-6.69 (2H, m), 6.94 (1H, d, J=7Hz), 7.13 (2H, d, J=8Hz), 7.31-7.43 (4H, m)

20

5) 2-(4-Methoxycarbonyl)phenylmethoxy-4-methylamine

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.24 (3H, s), 3.90 (3H, s), 5.11 (3H, s), 6.60-6.68 (3H, m), 7.50 (2H, d, J=8Hz), 8.05 (2H, d, J=8Hz)

25

6) 4-Amino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]-phenylbenzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.28 (3H, s), 2.33 (3H, s), 2.37-2.53 (4H, m), 3.36 (3H, s), 3.41-3.54 (2H, m), 3.57 (3H, s), 3.65-3.90 (4H, m), 4.90 (1H, d, J=14Hz), 5.06 (1H, d, J=14Hz), 6.38 (1H, d, J=7Hz), 6.62-6.70 (2H, m), 6.78 (1H, d, J=7Hz), 6.84 (1H, s), 6.98 (1H, d, J=7Hz), 7.33 (2H, d, J=8Hz), 7.41 (2H, d, J=8Hz)

30

35

- 72 -

- 7) Methyl 4-[(E and Z)-2-(2-aminophenyl)ethen-1-yl]benzoate  
NMR (CDCl<sub>3</sub>, δ) : 3.72 (2H, br), 3.86 (3Hx2/3, s),  
3.90 (3Hx1/3, s), 6.57-7.43 (7H, m), 7.55 (1H, d, J=7Hz), 7.86 (1H, d, J=7Hz), 8.01 (7H, d)
- 8) 4-Amino-3-methoxy-N-[(E and Z)-2-(4-methoxycarbonylphenyl)ethen-1-yl]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 3.39 (3Hx2/3, s), 3.40 (3Hx1/3, s),  
3.50 (3Hx2/3, s), 3.51 (3Hx1/3, s), 3.81-3.96 (2H, m), 3.84 (3Hx2/3, s), 3.41 (3Hx1/3, s),  
6.30-8.05 (13H, m)
- 9) 4-Amino-3-methoxy-N-[2-(4-methoxycarbonyl)-phenylmethoxy-4-methyl]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.21 (3H, s), 3.34 (3H, s), 3.50 (3H, s), 3.83 (2H, s), 3.90 (3H, s), 4.79-5.14 (2H, m), 6.37 (1H, d, J=7Hz), 6.60 (1H, s), 6.70 (1H, d, J=7Hz), 6.77 (1H, d, J=7Hz), 6.81 (1H, s), 6.99 (1H, d, J=7Hz), 7.34 (2H, d, J=8Hz), 8.01 (2H, d, J=8Hz)
- 10) 2-[3-(Ethoxycarbonylmethyl)oxyprop-1-yl]oxyaniline  
NMR (CDCl<sub>3</sub>, δ) : 1.27 (3H, t, J=7.5Hz), 2.08-2.28 (2H, m), 3.72 (2H, t, J=7.5Hz), 3.79 (2H, s), 4.09 (2H, s), 4.14 (2H, t, J=7.5Hz), 4.21 (2H, q, J=7.5Hz), 6.65-6.82 (4H, m)
- 11) 4-Amino-3-methoxy-N-[2-[3-(ethoxycarbonylmethyl)-oxyprop-1-yl]oxy]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7.5Hz), 2.03-2.15 (2H, m), 3.31 (3H, s), 3.61 (3H, s), 3.69-3.77 (4H, m), 4.02 (2H, s), 4.20 (2H, q, J=7.5Hz), 6.41 (1H, d, J=7.5Hz), 6.64-6.89 (4H, m), 7.00 (1H, d, J=7Hz), 7.13 (1H, t, J=7Hz)

- 73 -

- 12) 2-[(E)-5-Ethoxycarbonyl-4-penten-1-yl]oxy-4-methylaniline  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.29 (3H, t, J=7.5Hz), 1.90-2.05 (2H, m), 2.23 (3H, s), 2.35-2.50 (2H, m), 3.65 (2H, br), 4.00 (2H, t, J=7.5Hz), 4.18 (2H, q, J=7.5Hz), 5.98 (1H, d, J=15Hz), 6.53-6.67 (2H, m), 6.81 (1H, s), 7.00 (1H, dt, J=15, 7.5Hz)
- 13) 4-Amino-3-methoxy-N-[2-[(E)-5-ethoxycarbonyl-4-penten-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.27 (3H, t, J=7.5Hz), 1.84-1.98 (2H, m), 2.36 (3H, s), 2.31-2.41 (2H, m), 3.29 (3H, s), 3.62 (3H, s), 3.75-3.96 (4H, m), 4.18 (2H, q, J=7.5Hz), 5.84 (1H, d, J=15Hz), 6.40 (1H, d, J=7Hz), 6.58-6.63 (2H, m), 6.78-7.01 (4H, m)
- 14) 2-(5-Ethoxycarbonylpent-1-yloxy)-4-methylaniline  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.26 (3H, t, J=7Hz), 1.45-1.60 (2H, m), 1.63-1.89 (4H, m), 2.25 (3H, s), 2.33 (2H, t, J=7Hz), 3.98 (2H, t, J=7Hz), 4.13 (2H, q, J=7Hz), 6.54-6.68 (3H, m)
- 15) 3-Methoxy-4-amino-N-(2-benzyloxy-4-methylphenyl)-N-methylbenzamide  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.28 (3H, s), 3.32 (3H, s), 3.49 (3H, s), 3.83 (2H, br), 4.80-5.11 (2H, br), 6.34 (1H, d, J=8Hz), 6.62-6.84 (5H, m), 6.92 (1H, d, J=8Hz), 7.25-7.39 (4H, m)
- 16) 4-Amino-3-methyl-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.48-1.59 (2H, m), 1.63-1.88 (4H, m), 2.00 (3H, s), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.40 (6H, m), 3.29 (3H, s), 3.43-3.48 (2H,

- 74 -

m), 3.62 (4H, br), 3.90 (2H, br), 6.32 (1H, d, J=7Hz), 6.56-6.61 (2H, m), 6.83 (1H, d, J=7Hz), 6.90 (1H, d, J=7Hz), 7.17 (1H, s)

- 5     17) 4-Amino-3-hydroxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

10     NMR (CDCl<sub>3</sub>, δ) : 1.62 (6H, br), 2.28 (3H, s), 2.31 (3H, s), 2.38-2.49 (6H, m), 3.28 (3H, s), 3.52 (2H, br), 3.67 (2H, br), 3.78 (2H, br), 3.91 (2H, br), 6.32-6.38 (1H, m), 6.57-6.67 (3H, m), 7.00-7.03 (2H, m)

#### Preparation 6

15     The following compounds were obtained by reacting the compounds, which were prepared according to a similar manner to that of Preparation 4, with hydrogen chloride.

- 20     1) Benzyl 4-amino-3-benzyloxybenzoate hydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 5.18 (2H, s), 5.25 (2H, s), 5.98 (2H, br), 6.78 (1H, d, J=7Hz), 7.29-7.52 (12H, m)

- 25     2) Methyl 2-amino-5-thiophenecarboxylate hydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 3.68 (3H, s), 5.90 (1H, d, J=5Hz), 7.32-7.37 (2H, m)

#### Preparation 7

30     The following compounds were obtained according to a similar manner to that of Preparation 1.

- 35     1) 2-(3-Hydroxyprop-1-yl)oxynitrobenzene  
NMR (CDCl<sub>3</sub>, δ) : 2.07-2.14 (2H, m), 2.22 (1H, t, J=7.5Hz), 3.90 (2H, dd, J=7.5, 7.5Hz), 4.29 (2H, t, J=7Hz), 7.01 (1H, t, J=7Hz), 7.12 (1H, t,



- 75 -

J=7Hz), 7.54 (1H, t, J=7Hz), 7.89 (1H, d, J=7Hz)

2) 3-(3-Ethoxycarbonylprop-1-yl)oxy-4-nitrotoluene

5 NMR (CDCl<sub>3</sub>, δ) : 1.24 (3H, t, J=7.5Hz), 2.09-2.19  
(2H, m), 2.56 (2H, t, J=7.5Hz), 4.08-4.20 (4H,  
m), 6.81 (1H, d, J=7Hz), 6.97 (1H, s), 7.77 (7H,  
d)

3) Benzyl 2-(3-phthalimidopropoxy)benzoate

10 NMR (CDCl<sub>3</sub>, δ) : 2.08-2.23 (2H, m), 3.85 (2H, t,  
J=7Hz), 4.07 (2H, t, J=7Hz), 5.32 (2H, s), 6.86-  
7.02 (2H, m), 7.20-7.50 (6H, m), 7.61-7.74 (2H,  
m), 7.75-7.90 (3H, m)

15 4) 2-(5-Ethoxycarbonylpent-1-yloxy)-4-methylnitrobenzene

20 NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7Hz), 1.46-1.63 (2H,  
m), 1.63-1.78 (2H, m), 1.79-1.94 (2H, m), 2.34  
(2H, t, J=7Hz), 2.40 (3H, s), 4.00-4.19 (4H, m),  
6.80 (1H, d, J=9Hz), 6.84 (1H, s), 7.76 (1H, d,  
J=9Hz)

5) 2-Benzyloxy-N-tert-butoxycarbonylaniline

25 NMR (CDCl<sub>3</sub>, δ) : 1.49 (9H, s), 5.10 (2H, s), 6.88-  
6.98 (3H, m), 7.09 (1H, s), 7.32-7.43 (5H, m),  
8.10 (1H, br)

6) Methyl 4-[N-[2-[(3-tert-butoxycarbonylaminoprop-1-yl)oxy]phenyl]-tert-butoxycarbonylamino]methyl-3-methoxybenzoate

30 NMR (CDCl<sub>3</sub>, δ) : 1.33 and 1.42 (total 18H, s),  
1.92-2.00 (2H, m), 3.26-3.32 (2H, m), 3.70 and  
3.77 (total 3H, s), 3.90 (3H, s), 4.03 (2H, br),  
4.72 (2H, br), 6.72-6.97 (3H, m), 7.10-7.23 (2H,  
m), 7.40-7.53 (2H, m), 7.62 (1H, br)

35

- 76 -

- 7) 1-Benzyloxy-2-(3-tert-butoxycarbonylamino-  
prop-1-yl)oxybenzene

NMR (CDCl<sub>3</sub>, δ) : 1.40 and 1.47 (9H, s), 1.98-2.06  
(2H, m), 3.23-3.47 (2H, m), 4.10 (2H, t, J=6Hz),  
5.18 (2H, s), 5.42 (1H, br), 6.82-6.90 (4H, m),  
7.28-7.47 (5H, m)

- 8) Methyl 4-[2-[(3-tert-butoxycarbonylamino-  
prop-1-yl)oxymethyl]-3-methoxybenzoate

NMR (CDCl<sub>3</sub>, δ) : 1.38 (9H, s), 2.02 (2H, br), 3.38  
(2H, br), 3.90-3.92 (6H, m), 4.10-4.16 (2H, m),  
5.23 (1H, s), 5.25 (1H, s), 5.44 (1H, br), 6.83-  
6.92 (4H, m), 7.53-7.57 (2H, m), 7.65-7.69 (1H,  
m)

- 9) Benzyl 3-benzyloxy-4-nitrobenzoate

NMR (CDCl<sub>3</sub>, δ) : 5.28 (2H, s), 5.89 (2H, s), 7.30-  
7.48 (9H, m), 7.70-7.73 (1H, m), 7.81-7.85 (2H,  
m)

- 10) Benzyl 3-benzyloxy-4-[2-[(3-tert-butoxycarbonylamino-  
prop-1-yl)oxy]benzoyl]aminobenzoate

NMR (CDCl<sub>3</sub>, δ) : 1.38 (9H, s), 1.60-1.70 (2H, m),  
2.95-3.02 (2H, m), 3.80 (2H, t, J=6Hz), 4.42  
(1H, br), 5.22 (2H, s), 5.38 (2H, s), 6.93 (1H,  
d, J=8Hz), 7.10 (1H, t, J=7Hz), 7.32-7.50 (12H,  
m), 7.71-7.72 (1H, m), 7.80-7.83 (1H, m), 8.23-  
8.28 (1H, m), 8.78 (1H, d, J=7Hz)

- 11) Methyl 2-[2-[(3-tert-butoxycarbonylamino-  
prop-1-yl)oxy]benzoyl]amino-5-thiophenecarboxylate

This compound was used for further reaction  
without purification.

- 77 -

Preparation 8

The following compounds were obtained according to a similar manner to that of Preparation 2.

- 5     1) 4-[N-Methyl-2-[(3-tert-butoxycarbonylamino-3-methoxybenzoic acid  
yl)oxy]benzoyl]amino-3-methoxybenzoic acid  
NMR (CDCl<sub>3</sub>, δ) : 1.45 (9H, s), 1.97-2.06 (2H, m),  
3.33-3.42 (5H, m), 3.87 (3H, s), 3.98-4.07 (2H,  
m), 5.27-5.35 (1H, br), 6.67-6.76 (2H, m), 7.03-  
10     7.19 (3H, m), 7.44-7.50 (2H, m)  
ESI-MASS (m/z) : 459 (M+H)
- 2) 4-Nitro-N-[2-(4-carboxyphenyl)methoxy-4-  
methyl]phenyl-N-methylbenzamide  
15     NMR (CDCl<sub>3</sub>, δ) : 2.27 (3H, s), 3.40 (3H, s), 4.97  
(1H, d, J=14Hz), 5.10 (1H, d, J=14Hz), 6.65 (1H,  
s), 6.68 (1H, d, J=7Hz), 7.00 (1H, d, J=7Hz),  
7.33-7.49 (4H, m), 7.97 (2H, d, J=8Hz), 8.10  
(2H, d, J=8Hz)  
20
- 3) 3-Methoxy-4-nitro-N-[2-(4-carboxy)phenylmethoxy-4-  
methyl]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.30 (3H, s), 3.42 (3H, s), 3.61  
(3H, s), 4.92 (1H, d, J=14Hz), 5.11 (1H, d,  
25     J=14Hz), 6.65 (1H, s), 6.73 (1H, d, J=7Hz), 6.86  
(1H, d, J=7Hz), 7.02-7.08 (2H, m), 7.48 (2H, d,  
J=8Hz), 7.54 (1H, d, J=7Hz), 8.16 (2H, d, J=8Hz)
- 4) 2-(4-Carboxyphenylmethyl)oxy-4-methyl-N,N-  
dimethylaniline  
30     NMR (CDCl<sub>3</sub>, δ) : 2.31 (3H, s), 2.89 (6H, s), 5.08  
(2H, s), 6.76-7.82 (2H, m), 7.03 (1H, d, J=7Hz),  
7.40 (2H, d, J=8Hz), 7.77 (2H, d, J=8Hz)
- 35     5) 2-[3-(4-Methoxyphenyl)methoxypropyl-1-yl]thiobenzoic

- 78 -

acid

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.95-2.06 (2H, m), 3.03 (2H, t, J=7.5Hz), 3.59 (2H, t, J=7.5Hz), 3.77 (3H, s), 4.46 (2H, s), 6.89 (2H, d, J=8Hz), 7.19 (1H, t, J=7Hz), 7.16 (2H, d, J=8Hz), 7.36 (1H, d, J=7Hz), 7.45 (1H, t, J=7Hz), 8.10 (1H, d, J=7Hz)

6) 4-Amino-3-methoxy-N-[2-(4-carboxy)phenylmethoxy-4-methyl]phenyl-N-methylbenzamide

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 2.21 (3H, s), 3.15 (3H, s), 3.41 (3H, s), 4.95-5.23 (2H, m), 6.33 (1H, d, J=7Hz), 6.63-6.72 (3H, m), 6.87 (1H, s), 7.04 (1H, d, J=7Hz), 7.44 (2H, d, J=8Hz), 7.95 (2H, d, J=8Hz)

7) 4-Amino-3-methoxy-N-[2-[3-(carboxymethyl)oxyprop-1-yl]oxyphenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.00-2.12 (2H, m), 3.32 (3H, s), 3.60 (3H, s), 3.63-3.74 (2H, m), 3.89-4.14 (2H, m), 4.05 (2H, s), 4.50 (2H, br), 6.40 (1H, d, J=7Hz), 6.80-6.95 (4H, m), 6.95 (1H, d, J=7Hz), 7.16 (1H, t, J=7Hz)

8) 4-Amino-3-methoxy-N-[2-[(E)-5-ethoxycarbonyl-4-penten-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.87-1.99 (2H, m), 2.28 (3H, s), 2.34-2.45 (2H, m), 3.31 (3H, s), 3.61 (3H, s), 3.71-4.00 (2H, m), 5.87 (1H, d, J=15Hz), 6.41 (1H, d, J=7Hz), 6.57-6.68 (2H, m), 6.80-7.12 (4H, m)

9) 3-(5-Carboxypent-1-yloxy)-4-(tert-butoxycarbonylamino)toluene

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.45-1.63 (11H, m), 1.64-1.95 (4H, m), 2.28 (3H, s), 2.42 (2H, t, J=7Hz), 3.99 (2H, t, J=7Hz), 6.65 (1H, s), 6.72 (1H, d, J=8Hz),

- 79 -

6.98 (1H, s), 7.87 (1H, m)

- 10) 4-[(2-Benzyloxy)benzoyl]amino-3-chlorobenzoic acid  
NMR (CDCl<sub>3</sub>, δ) : 5.49 (2H, s), 7.18 (1H, t, J=6Hz),  
7.32-7.42 (4H, m), 7.50-7.62 (3H, m), 7.89-7.93  
(2H, m), 8.10 (1H, d, J=7Hz), 8.58-8.62 (1H, m)
- 11) 4-[2-(Benzyloxy)benzoyl]amino-2-nitrobenzoic acid  
NMR (DMSO-d<sub>6</sub>, δ) : 5.22 (2H, s), 7.10 (1H, t,  
J=7Hz), 7.28-7.38 (4H, m), 7.50-7.58 (3H, m),  
7.65-7.69 (1H, m), 7.86 (2H, s), 8.16 (1H, s)
- 12) 2-[2-(Benzyloxy)benzoyl]amino-5-pyridinecarboxylic acid  
NMR (DMSO-d<sub>6</sub>, δ) : 5.18 (1H, s), 5.32 (2H, s), 6.98-  
7.20 (2H, m), 7.29-7.67 (6H, m), 7.84-7.88 (1H,  
m), 8.28-8.37 (2H, m), 8.80 (1H, s)
- 13) 4-[N-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl]oxy]phenyl]-tert-butoxycarbonylamino]methyl-3-methoxybenzoic acid  
NMR (CDCl<sub>3</sub>, δ) : 1.35 and 1.43 (total 18H, s), 1.92-  
2.00 (2H, m), 3.28 and 3.32 (total 2H, m), 3.20  
and 3.28 (total 3H, s), 4.02 (2H, br), 4.77 (2H,  
br), 6.77-7.99 (3H, m), 7.10-7.20 (2H, m), 7.44-  
7.56 (2H, m), 7.69 (1H, br)
- 14) 4-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl]oxymethyl-3-methoxybenzoic acid  
NMR (CDCl<sub>3</sub>, δ) : 1.37 (9H, s), 2.05 (2H, br), 3.40  
(2H, br), 3.93 (3H, s), 4.10-4.17 (2H, m), 5.27  
(2H, s), 5.50 (1H, br), 6.87-6.93 (4H, m), 7.59  
(2H, s), 7.72-7.77 (1H, m)
- 15) 3-Benzyloxy-4-[2-[(3-tert-butoxycarbonylamino)prop-1-

- 80 -

yl)oxy]benzoyl]aminobenzoic acid

NMR (DMSO-d<sub>6</sub>, δ) : 1.30 (9H, s), 1.62-1.72 (2H, m),  
2.88-2.92 (2H, m), 3.95 (2H, t, J=6Hz), 5.37  
(2H, s), 6.80 (1H, br), 7.13 (1H, t, J=7Hz),  
7.21 (1H, d, J=7Hz), 7.30-7.67 (9H, m), 8.08  
(1H, d, J=7Hz), 8.60 (1H, d, J=7Hz)

16) 2-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-5-thiophenecarboxylic acid

NMR (DMSO-d<sub>6</sub>, δ) : 1.32 (9H, s), 1.82-1.90 (2H, m),  
3.08-3.14 (2H, m), 4.10 (2H, t, J=6Hz), 6.81  
(1H, d, J=5Hz), 6.93-7.00 (1H, m), 7.07 (1H, t,  
J=7Hz), 7.19 (1H, d, J=7Hz), 7.50-7.58 (2H, m),  
7.67 (1H, d, J=7Hz)

#### Preparation 9

The following compounds were obtained according to a similar manner to that of Preparation 3.

1) 3-Methoxy-4-nitro-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.30-1.96 (10H, m), 2.28 (9H, s),  
2.30-2.41 (3H, m), 2.58 (1H, m), 3.02 (1H, m),  
3.33 (3H, s), 3.77 (3H, s), 3.82-4.00 (3H, m),  
4.63 (1H, m), 6.56-6.66 (2H, m), 6.84 (1H, d,  
J=9Hz), 6.93 (1H, d, J=9Hz), 7.06 (1H, s), 7.61  
(1H, d, J=9Hz)

2) 4-Nitro-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 2.26 (3H, s), 2.32 (3H, s), 2.36-  
2.57 (4H, m), 3.37 (3H, s), 3.42-3.59 (2H, m),  
3.71-3.89 (2H, m), 4.94 (1H, d, J=14Hz), 5.07  
(1H, d, J=14Hz), 6.60-6.69 (2H, m), 6.94 (1H, d,

- 81 -

J=7Hz), 7.36-7.50 (5H, m), 7.95 (2H, d, J=8Hz)

- 3) 4-Amino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]-phenylbenzamide

5  
NMR (CDCl<sub>3</sub>, δ) : 2.28 (3H, s), 2.33 (3H, s), 2.37-2.53 (4H, m), 3.36 (3H, s), 3.41-3.54 (2H, m), 3.57 (3H, s), 3.65-3.90 (4H, m), 4.90 (1H, d, J=14Hz), 5.06 (1H, d, J=14Hz), 6.38 (1H, d, J=7Hz), 6.62-6.70 (2H, m), 6.78 (1H, d, J=7Hz), 6.84 (1H, s), 6.98 (1H, d, J=7Hz), 7.33 (2H, d, J=8Hz), 7.41 (2H, d, J=8Hz)

- 4) 4-Amino-3-methoxy-N-[2-[4-(4-dimethylaminopiperidin-1-yl)carbonyl]phenylmethoxy-4-methyl]phenyl-N-methylbenzamide

15  
NMR (CDCl<sub>3</sub>, δ) : 1.14-1.58 (2H, m), 1.75-2.00 (2H, m), 2.26 (3H, s), 2.30 (3H, s), 2.40 (1H, m), 2.73-3.10 (4H, m), 3.36 (3H, s), 3.57 (3H, s), 3.87 (3H, s), 4.83-5.12 (2H, m), 6.39 (1H, d, J=7Hz), 6.61-6.71 (2H, m), 6.28 (1H, d, J=7Hz), 6.33 (1H, s), 6.97 (1H, d, J=7Hz), 7.33 (2H, d, J=8Hz), 7.40 (2H, d, J=8Hz)

- 5) 4-Amino-3-methoxy-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylmethoxyprop-1-yl]oxy]phenylbenzamide

25  
NMR (CDCl<sub>3</sub>, δ) : 1.98-2.13 (2H, m), 2.27 (3H, s), 2.29-2.38 (4H, m), 3.30 (3H, s), 3.36-3.47 (2H, m), 3.52-3.74 (4H, m), 3.60 (3H, s), 3.94-4.17 (2H, m), 4.11 (2H, s), 6.42 (1H, d, J=7Hz), 6.78-6.92 (4H, m), 7.00 (1H, d, J=7Hz), 7.14 (1H, t, J=7Hz)

- 6) 4-Amino-3-methoxy-N-[2-(E)-5-(4-

- 82 -

dimethylaminopiperidin-1-yl)carbonyl-4-penten-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide

5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.30-1.47 (2H, m), 1.80-1.98 (2H, m), 2.21 (3H, s), 2.26 (6H, s), 2.26-2.43 (2H, m), 2.45-3.67 (6H, m), 3.30 (3H, s), 3.61 (3H, s), 3.85 (2H, br), 3.85-4.04 (2H, m), 4.62 (1H, m), 6.29 (1H, d, J=15Hz), 6.41 (1H, d, J=7Hz), 6.57-6.63 (2H, m), 6.77-6.90 (4H, m)

10 7) 3-[5-(4-Dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-(tert-butoxycarbonylamino)toluene

15 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.27-2.00 (19H, m), 2.21-2.44 (12H, m), 2.58 (1H, m), 3.01 (1H, m), 3.89 (1H, m), 4.00 (2H, t, J=7Hz), 4.64 (1H, m), 6.64 (1H, s), 6.72 (1H, d, J=8Hz), 6.94 (1H, s), 7.89 (1H, m)

#### Preparation 10

The following compounds were obtained according to a similar manner to that of Example 1.

20 1) Methyl 4-(2-benzyloxybenzoyl)amino-3-methoxybenzoate

25 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 3.50 (3H, s), 3.90 (3H, s), 5.36 (2H, s), 7.08 (1H, d, J=9Hz), 7.15 (1H, t, J=9Hz), 7.33-7.49 (8H, m), 7.73 (1H, dd, J=1, 8Hz), 8.30 (1H, d, J=8Hz), 8.72 (1H, d, J=8Hz)  
ESI-MASS (m/z) : 392 (M+H)

30 2) Methyl 4-(2-acetoxybenzoyl)amino-3-methoxybenzoate

35 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.38 (3H, s), 3.92 (3H, s), 3.99 (3H, s), 7.19 (1H, d, J=8Hz), 7.38 (1H, t, J=8Hz), 7.55 (1H, t, J=8Hz), 7.60 (1H, s), 7.75 (1H, dd, J=2, 9Hz), 7.99 (1H, dd, J=1, 9Hz), 8.66 (1H, d, J=8Hz), 9.03-9.07 (1H, br s)  
ESI-MASS (m/z) : 344 (M+H)



- 83 -

- 3) 3-Methoxy-4-nitro-N-[2-(4-methoxycarbonyl)-phenylmethoxy-4-methyl]phenylbenzamide  
NMR (DMSO-d<sub>6</sub>, δ) : 2.31 (3H, s), 3.84 (3H, s), 3.98 (3H, s), 5.27 (2H, s), 6.81 (1H, d, J=7Hz), 7.00 (1H, s), 7.49 (1H, d, J=7Hz), 7.62 (2H, d, J=8Hz), 7.79 (1H, s), 7.92 (2H, d, J=8Hz), 8.00 (1H, d, J=7Hz), 9.85 (1H, s)
- 4) 4-Nitro-3-methoxy-N-[(E and Z)-2-(4-methoxycarbonylphenyl)ethen-1-yl]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 3.87 (3Hx2/3, s), 3.91 (3Hx1/3, s), 3.95 (3Hx2/3, s), 4.00 (3Hx1/3, s), 6.71-8.20 (13H, m)
- 5) 3-Methoxy-4-nitro-N-[2-[3-(ethoxycarbonylmethyl)-oxyprop-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.22 (3H, t, J=7.5Hz), 2.10-2.23 (2H, m), 3.78 (2H, t, J=7.5Hz), 4.01 (2H, s), 4.06 (3H, s), 4.14 (2H, q, J=7.5Hz), 4.26 (2H, t, J=7.5Hz), 6.91-7.06 (3H, m), 7.42 (1H, d, J=7Hz), 7.74 (1H, s), 7.93 (1H, d, J=7Hz), 8.49 (1H, d, J=7Hz), 7.78 (1H, s)
- 6) 3-Methoxy-4-nitro-N-[2-[(E)-5-ethoxycarbonyl-4-penten-1-yl]oxy-4-methyl]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.27 (3H, t, J=7.5Hz), 1.93-2.08 (2H, m), 2.27-2.50 (2H, m), 2.32 (3H, s), 4.02 (3H, s), 4.01-4.11 (2H, m), 4.18 (2H, q, J=7.5Hz), 5.88 (1H, d, J=15Hz), 6.72 (1H, s), 6.83 (1H, t, J=7Hz), 6.99 (1H, dt, J=15, 7.5Hz), 7.35 (1H, d, J=7Hz), 7.81 (1H, s), 7.92 (1H, d, J=7Hz), 8.28 (1H, d, J=7Hz), 8.45 (1H, s)
- 7) 4-Benzyloxy-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-

- 84 -

methylphenyl]benzamide

5 NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.35-1.49 (2H, m), 1.49-1.63 (2H, m), 1.64-1.79 (2H, m), 2.23 (3H, s), 2.37 (2H, t, J=7Hz), 2.72 (3H, m), 2.78-3.11 (2H, m), 3.16 (3H, s), 3.28-3.60 (5H, m), 3.71-4.13 (5H, m), 4.43 (1H, m), 4.99 (2H, s), 6.63 (1H, d, J=8Hz), 6.80 (2H, d, J=2Hz), 6.86 (2H, s), 6.98 (1H, d, J=8Hz), 7.26-7.44 (5H, m)

10 8) 3-Methoxy-4-nitro-N-(2-benzyloxy-4-methylphenyl)-benzamide

15 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.38 (3H, s), 3.90 (3H, s), 5.12 (2H, s), 6.88 (1H, s), 7.30 (1H, s), 7.51 (4H, s), 7.59 (1H, s), 7.82 (1H, d, J=8Hz), 8.37 (1H, d, J=8Hz), 8.53 (1H, br)

20 9) 3-Methyl-4-nitro-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

25 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.48-1.60 (2H, m), 1.69-1.77 (2H, m), 1.79-1.90 (2H, m), 2.25 (3H, s), 2.29 (3H, s), 2.33-2.42 (6H, m), 2.47 (3H, s), 3.32 (3H, s), 3.45-3.50 (2H, m), 3.58-3.63 (2H, m), 3.82-3.95 (2H, m), 6.55-6.59 (2H, m), 6.83 (1H, d, J=7Hz), 7.14 (1H, d, J=7Hz), 7.37 (1H, s), 7.70 (1H, d, J=7Hz)

30 10) Ethyl 4-[(2-benzyloxy)benzoyl]amino-3-chlorobenzoate

35 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.38 (3H, t, J=7Hz), 4.34 (2H, q, J=7Hz), 5.38 (1H, s), 5.39 (1H, s), 7.03-7.16 (2H, m), 7.33-7.50 (6H, m), 7.92-7.99 (2H, m), 8.24-8.32 (1H, m), 8.73-8.29 (1H, m)

11) 3-Hydroxy-4-nitro-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

- 85 -

- 5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.48-1.60 (2H, m), 1.68-1.80 (2H, m), 1.82-1.91 (2H, m), 2.28 (3H, s), 2.30 (3H, s), 2.37-2.42 (6H, m), 3.32 (3H, s), 3.48-3.50 (2H, m), 3.62-3.68 (2H, m), 3.90-3.97 (2H, m), 6.57-6.58 (2H, m), 6.80-6.87 (2H, m), 7.08-7.10 (1H, m), 7.85 (1H, d, J=7Hz)
- 10 12) Ethyl 4-[2-(benzyloxy)benzoyl]amino-2-nitrobenzoate  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.32 (3H, t, J=7Hz), 4.32 (2H, q, J=7Hz), 5.22-5.30 (2H, m), 7.12-7.27 (2H, m), 7.37-7.69 (9H, m), 8.20-8.34 (1H, m)
- 15 13) Methyl 2-[2-(benzyloxy)benzoyl]amino-5-pyridinecarboxylate  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 3.92 (3H, s), 5.12 (1H, s), 5.36 (2H, s), 6.90-7.01 (1H, m), 7.10-7.18 (2H, m), 7.32-7.55 (5H, m), 8.27-8.34 (2H, m), 8.46 (1H, d, J=6Hz), 8.87-8.88 (1H, m)
- 20 14) Benzyl 4-(2-acetoxybenzoyl)amino-3-benzyloxybenzoate  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.05 (3H, s), 5.20 (2H, s), 5.87 (2H, s), 7.13 (1H, d, J=8Hz), 7.32-7.47 (10H, m), 7.50-7.57 (1H, m), 7.73 (1H, s), 7.80 (1H, d, J=8Hz), 7.96 (1H, d, J=8Hz), 8.68 (1H, d, J=7Hz), 9.13 (1H, s)
- 25 15) Methyl 2-(2-acetoxybenzoyl)amino-5-thiophenecarboxylate  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.39 (3H, s), 3.88 (3H, s), 6.69 (1H, d, J=5Hz), 7.19-7.21 (1H, m), 7.35-7.30 (1H, m), 7.52-7.59 (1H, m), 7.63-7.66 (1H, m), 7.92-7.95 (1H, m), 9.18 (1H, s)
- 30

Preparation 11

35 The following compound was obtained by reacting the

- 86 -

compound, which was prepared according to a similar manner to that of Example 1, with hydrogen chloride.

5 4-Benzyloxy-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

10 NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.35-1.49 (2H, m), 1.49-1.63 (2H, m), 1.64-1.79 (2H, m), 2.23 (3H, s), 2.37 (2H, t, J=7Hz), 2.72 (3H, m), 2.78-3.11 (2H, m), 3.16 (3H, s), 3.28-3.60 (5H, m), 3.71-4.13 (5H, m), 4.43 (1H, m), 4.99 (2H, s), 6.63 (1H, d, J=8Hz), 6.60 (2H, d, J=2Hz), 6.86 (2H, s), 6.98 (1H, d, J=8Hz), 7.26-7.44 (5H, m)

15 Preparation 12

The following compound was obtained according to a similar manner to that of Example 4.

20 4-[2-[(3-tert-Butoxycarbonylamino-3-methoxybenzoic acid

25 NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.35 (9H, s), 2.04 (2H, quintet, J=7Hz), 3.13 (2H, q, J=7Hz), 3.98 (3H, s), 4.29 (2H, t, J=7Hz), 6.95-7.00 (1H, m), 7.16 (1H, t, J=8Hz), 7.28 (1H, d, J=8Hz), 7.57-7.65 (3H, m), 8.11 (1H, dd, J=1, 8Hz), 8.63 (1H, d, J=8Hz)  
ESI-MASS (m/z) : 445 (M+H)

Preparation 13

30 The following compounds were obtained according to a similar manner to that of Example 10.

1) 4-Hydroxy-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

35 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.44-1.59 (2H, m), 1.62-1.92 (4H,

- 87 -

m), 2.22-2.45 (12H, m), 3.31 (3H, s), 3.42-3.53 (2H, m), 3.58-3.74 (5H, m), 3.77-4.02 (2H, m), 6.53-6.70 (3H, m), 6.80-6.96 (3H, m)

- 5        2) Methyl 4-(N-methyl-2-hydroxybenzoylamino)-3-methoxybenzoate

NMR (CDCl<sub>3</sub>, δ) : 3.37 (3H, s), 3.69 (3H, s), 3.91 (3H, s), 6.38 (1H, t, J=8Hz), 6.72 (1H, d, J=8Hz), 6.91 (1H, d, J=8Hz), 7.15 (1H, t, J=8Hz), 7.21 (1H, d, J=9Hz), 7.49 (1H, d, J=1Hz), 7.62 (1H, dd, J=1, 9Hz)

ESI-MASS (m/z) : 316 (M+H)

- 15        3) 4-Hydroxy-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.25-2.00 (10H, m), 2.06-2.40 (6H, m), 2.52 (1H, m), 2.73 (6H, br s), 3.02 (1H, m), 3.30 (3H, s), 3.67 (3H, s), 3.76-4.07 (3H, m), 4.82 (1H, m), 6.56-6.72 (3H, m), 6.78-6.96 (3H, m)

- 25        4) Methyl 4-[N-(2-hydroxyphenyl)-tert-butoxycarbonylamino]methyl-3-methoxybenzoate

NMR (CDCl<sub>3</sub>, δ) : 1.38 (9H, s), 3.82 and 3.83 (total 3H, s), 3.90 and 3.91 (total 3H, s), 4.88 (2H, s), 6.80-6.87 (1H, m), 6.95 (1H, br), 7.03-7.12 (2H, m), 7.25-7.30 (2H, m), 7.48-7.50 (1H, m), 7.58-7.60 (1H, m)

- 30        5) 2-(3-tert-Butoxycarbonylamino)prop-1-yl)oxyphenol

NMR (CDCl<sub>3</sub>, δ) : 1.45 (9H, s), 1.95-2.07 (2H, m), 3.25-3.45 (2H, m), 4.10 (2H, t, J=6Hz), 4.68 (1H, br), 6.22 (1H, br), 6.78-6.97 (4H, m)

35

Preparation 14

The following compounds were obtained according to a similar manner to that of Example 12.

- 5      1) Methyl 4-[N-methyl-2-[(3-tert-butoxycarbonylamino-prop-1-yl)oxy]benzoyl]amino-3-methoxybenzoate  
NMR (CDCl<sub>3</sub>, δ) : 1.43 (9H, s), 1.95-2.05 (2H, m),  
3.30-3.40 (5H, m), 3.83 (3H, s), 3.85 (3H, s),  
3.96-4.04 (2H, m), 5.23-5.32 (1H, br), 6.65-6.73  
10      (2H, m), 7.00-7.16 (3H, m), 7.38-7.45 (2H, m)  
ESI-MASS (m/z) : 473 (M+H)
- 15      2) Methyl 4-[2-[(3-tert-butoxycarbonylamino-prop-1-yl)oxy]benzoyl]amino-3-methoxybenzoate  
NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 2.13-2.21 (2H, m),  
3.33 (2H, q, J=7Hz), 3.92 (3H, s), 4.00 (3H, s),  
4.29 (2H, t, J=7Hz), 4.72-4.78 (1H, br), 7.03  
(1H, d, J=8Hz), 7.23 (1H, t, J=8Hz), 7.49 (1H,  
t, J=8Hz), 7.60 (1H, s), 7.75 (1H, d, J=8Hz),  
20      8.27 (1H, d, J=8Hz), 8.77 (1H, d, J=8Hz)  
ESI-MASS (m/z) : 459 (M+H)
- 25      3) 4-Nitro-N-[2-(4-methoxycarbonylphenyl)methoxy-4-methyl]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.27 (3H, s), 3.40 (3H, s), 3.94  
(3H, s), 4.95 (1H, d, J=14Hz), 5.09 (1H, d,  
J=14Hz), 6.62 (1H, s), 6.69 (1H, d, J=7Hz), 6.97  
(1H, d, J=7Hz), 7.31-7.49 (4H, m), 7.95 (2H, d,  
J=8Hz), 8.10 (2H, d, J=8Hz)
- 30

Preparation 15

The following compound was obtained according to a similar manner to that of Example 16.

- 35      4-Amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-

- 89 -

methylnpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide  
dihydrochloride

5 NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.33-1.64 (4H, m), 1.64-1.81 (2H, m), 2.20 (3H, s), 2.29-2.43 (2H, m), 2.73 (3H, s), 2.79-3.10 (4H, m), 3.14 (3H, s), 3.22-3.56 (4H, m), 3.62 (3H, s), 3.72-4.18 (3H, m), 4.42 (1H, m), 6.62 (1H, d, J=8Hz), 6.74-6.92 (3H, m), 6.92-7.10 (2H, m)

10 Preparation 16

The following compounds were obtained according to a similar manner to that of Example 43.

- 1) Methyl 4-(2-hydroxybenzoyl)amino-3-methoxybenzoate  
15 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 3.93 (3H, s), 4.03 (3H, s), 6.96 (1H, t, J=8Hz), 7.04 (1H, d, J=8Hz), 7.47 (1H, t, J=8Hz), 7.54 (1H, dd, J=1, 8Hz), 7.62 (1H, s), 7.76 (1H, d, J=8Hz), 8.51 (1H, d, J=8Hz), 8.85-8.89 (1H, br s)
- 20 ESI-MASS (m/z) : 302 (M+H)
- 2) Benzyl 3-benzyloxy-4-(2-hydroxybenzoyl)aminobenzoate  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 5.23 (2H, s), 5.38 (2H, s), 6.82 (1H, t, J=7Hz), 7.01 (1H, d, J=7Hz), 7.30-7.49 (12H, m), 7.70-7.73 (1H, m), 7.80-7.83 (1H, m), 7.52 (1H, d, J=7Hz), 8.95 (1H, s)
- 25 3) Methyl 2-(2-hydroxybenzoyl)amino-5-thiophenecarboxylate  
NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 3.79 (3H, s), 6.95-7.03 (3H, m), 7.42-7.48 (1H, m), 7.62-7.64 (1H, m), 7.88 (1H, d, J=7Hz)
- 30

35 Preparation 17

The following compound was obtained according to a

- 90 -

similar manner to that of Example 30.

3-Methoxy-4-nitro-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]-phenylbenzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.29 (3H, s), 2.35 (3H, s), 2.38-2.54 (4H, m), 3.39 (3H, s), 3.43-3.53 (2H, m), 3.66 (3H, s), 3.71-3.88 (2H, m), 4.92 (1H, d, J=14Hz), 5.07 (1H, d, J=14Hz), 6.65-6.72 (2H, m), 6.87 (1H, d, J=7Hz), 6.98 (1H, d, J=7Hz), 7.03 (1H, s), 7.37 (2H, d, J=8Hz), 7.45 (2H, d, J=8Hz), 7.56 (1H, d, J=7Hz)

#### Preparation 18

To a mixture of 2-(4-methoxycarbonylphenyl)methoxy-4-methylaniline (420 mg) and 37% formaldehyde solution (69.7 mg) in a mixture of methanol (10 ml) and acetic acid (0.1 ml) was added sodium cyanoborohydride (146 mg) and the mixture was stirred at ambient temperature for 3 hours. The solution was diluted with ethyl acetate (30 ml) and washed successively with saturated aqueous sodium hydrogen carbonate, water and brine. The organic solution was dried over magnesium sulfate and the solvent was evaporated in vacuo. The residue was purified by silica gel column (chloroform) to give 2-(4-methoxycarbonylphenyl)methoxy-4-methyl-N-methylaniline (356 mg).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.22 (3H, s), 2.80 (3H, s), 3.91 (3H, s), 5.11 (2H, s), 6.53 (1H, d, J=7Hz), 6.63 (1H, s), 6.72 (1H, d, J=7Hz), 7.49 (2H, d, J=8Hz), 8.04 (2H, d, J=8Hz)

#### Preparation 19

A solution of 2-benzyloxy-N-tert-butoxycarbonylaniline (1 g) in N,N-dimethylformamide (40



- 91 -

ml) was treated with sodium hydride (147 mg, 60% w/w in mineral oil) at 0°C. The reaction mixture was stirred at 0°C for 30 minutes and then at ambient temperature for 1 hour. Methyl 4-bromomethyl-3-methoxybenzoate (909 mg) was added, and the mixture was stirred at ambient temperature for 30 minutes. The reaction was quenched with water and the mixture diluted with ethyl acetate. The organic phase was washed with 0.5N hydrochloric acid, saturated aqueous sodium hydrogen carbonate, and brine. The solution was concentrated in vacuo and the residue was purified by silica gel column chromatography (hexane:ethyl acetate = 9:1) to give methyl 4-[N-[2-(benzyloxy)phenyl]-tert-butoxycarbonylamino]methyl-3-methoxybenzoate (1.38 g).

NMR (CDCl<sub>3</sub>, δ) : 1.32 and 1.40 (total 9H, s), 3.65 and 3.71 (total 3H, s), 3.90 (3H, s), 4.77 (2H, s), 5.07 (2H, s), 6.78-7.00 (3H, m), 7.09-7.20 (1H, m), 7.27-7.55 (8H, m)

#### Preparation 20

The following compounds were obtained according to a similar manner to that of Preparation 19.

1) 4-Nitro-3-methoxy-N-[(E and Z)-2-(4-methoxycarbonylphenyl)ethen-1-yl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 3.40 (3Hx2/3, s), 3.49 (3Hx1/3, s), 3.54 (3Hx1/3, s), 3.60 (3Hx2/3, s), 3.86 (3Hx2/3, s), 3.95 (3Hx1/3, s), 6.41-8.07 (7H, m)

2) 3-Methoxy-4-nitro-N-[2-[3-(ethoxycarbonylmethyl)-oxyprop-1-yl]oxy]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.27 (3H, t, J=7.5Hz), 2.04-2.17 (2H, m), 3.37 (3H, s), 3.71 (2H, t, J=7.5Hz), 3.76 (3H, s), 4.06 (3H, s), 4.20 (2H, q, J=7.5Hz), 6.78-7.01 (4H, m), 7.04 (1H, s), 7.19 (1H, t, J=7Hz), 7.60 (1H, d, J=7Hz)

- 92 -

## 3) 3-Methoxy-4-nitro-N-(2-benzyloxy-4-methylphenyl)-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 2.28 (3H, s), 3.39 (3H, s), 3.58 (3H, s), 4.85 (1H, d, J=12Hz), 5.07 (1H, d, J=12Hz), 6.68 (2H, s), 6.83 (1H, d, J=9Hz), 6.96 (1H, d, J=9Hz), 7.00 (1H, s), 7.30-7.44 (5H, m), 7.52 (1H, d, J=9Hz)

Preparation 21

To an ice bath cooled solution of methyl 2-(3-hydroxyprop-1-yl)thiobenzoate (3.7 g) in N,N-dimethylformamide (30 ml) was added sodium hydride (60% in oil, 719 mg) and the solution was stirred at the same temperature for 30 minutes. 4-Methoxybenzyl chloride (2.56 g) was added to the solution and the mixture was stirred at ambient temperature for 5 hours. The mixture was diluted with ethyl acetate (100 ml) and the solution was washed with water and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo to give a crude oil. The crude product was purified by silica gel column chromatography (hexane:ethyl acetate = 10:1) to give methyl 2-[3-(4-methoxyphenyl)methoxyprop-1-yl]thiobenzoate (2.13 g).

NMR (CDCl<sub>3</sub>, δ) : 1.94-2.07 (2H, m), 3.03 (2H, t, J=7.5Hz), 3.58 (2H, t, J=7.5Hz), 3.80 (3H, s), 3.90 (3H, s), 4.39 (2H, q, J=7.5Hz), 4.45 (2H, s), 6.87 (2H, d, J=8Hz), 7.13 (1H, t, J=7Hz), 7.21-7.46 (4H, m), 7.96 (1H, d, J=7Hz)

Preparation 22

The following compound was obtained according to a similar manner to that of Preparation 21.

2-[3-(Ethoxycarbonylmethyl)oxyprop-1-yl]oxynitrobenzene

- 93 -

NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7.5Hz), 2.08-2.20  
(2H, m), 3.73 (2H, t, J=7.5Hz), 4.06 (2H, s),  
4.13-4.32 (4H, m), 7.01 (1H, m), 7.10 (1H, d,  
J=7Hz), 7.50 (1H, t, J=7Hz), 7.82 (1H, d, J=7Hz)

5

#### Preparation 23

To an ice bath cooled solution of 3-methoxy-4-nitro-  
N-[2-(4-methoxycarbonyl)phenylmethoxy-4-methyl]-  
phenylbenzamide (7.67 g) in N,N-dimethylformamide (50 ml)  
10 was added sodium hydride (60% in oil, 817 mg) and the  
solution was stirred at the same temperature for 30  
minutes. Iodomethane (1.27 ml) was added to the solution  
and the mixture was stirred at ambient temperature for 2  
hours. The mixture was diluted with ethyl acetate (100  
15 ml) and the solution was washed with water and brine. The  
organic phase was dried over magnesium sulfate and the  
solvent was evaporated in vacuo to give an oil. The oil  
was solidified with diethyl ether to give 3-methoxy-4-  
nitro-N-[2-(4-methoxycarbonyl)phenylmethoxy-4-  
20 methyl]phenyl-N-methylbenzamide (6.65 g).

NMR (CDCl<sub>3</sub>, δ) : 2.28 (3H, s), 3.40 (3H, s), 3.60  
(3H, s), 3.94 (3H, s), 4.91 (1H, d, J=14Hz),  
5.09 (1H, d, J=14Hz), 6.64 (1H, s), 6.71 (1H, d,  
J=7Hz), 6.84 (1H, d, J=7Hz), 7.00-7.04 (2H, m),  
25 7.42 (2H, d, J=8Hz), 7.52 (1H, d, J=7Hz), 8.08  
(2H, d, J=8Hz)

#### Preparation 24

The following compound was obtained according to a  
30 similar manner to that of Preparation 23.

3-Methoxy-4-nitro-N-[2-[(E)-5-ethoxycarbonyl-4-  
penten-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.28 (3H, t, J=7.5Hz), 1.90-2.00  
35 (2H, m), 2.00 (3H, s), 2.34-2.45 (2H, m), 3.35

- 94 -

(3H, s), 3.77 (3H, s), 3.84-3.97 (2H, m), 4.19 (2H, q, J=7.5Hz), 5.88 (1H, d, J=15Hz), 6.58-6.64 (2H, m), 6.84-7.02 (3H, m), 7.07 (1H, s), 7.60 (1H, d, J=7Hz)

5

Preparation 25

The following compound was obtained according to a similar manner to that of Example 45.

10

2-[5-(4-Dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylaniline

NMR (CDCl<sub>3</sub>, δ) : 1.18-2.00 (10H, m), 2.14-2.69 (13H, m), 2.99 (1H, m), 3.44-4.07 (5H, m), 4.64 (1H, m), 6.45-6.70 (3H, m)

15

Preparation 26

The following compound was obtained according to a similar manner to that of Example 38.

20

2-Hydroxy-N-tert-butoxycarbonylaniline

NMR (CDCl<sub>3</sub>, δ) : 1.55 (9H, s), 6.63 (1H, s), 6.82-6.89 (1H, m), 6.97-6.99 (1H, m), 7.02-7.08 (2H, m), 8.13 (1H, br)

25

Preparation 27

The following compound was obtained according to a similar manner to that of Example 87.

30

Methyl 2-nitro-5-thiophenecarboxylate

NMR (CDCl<sub>3</sub>, δ) : 3.95 (3H, s), 7.70 (1H, d, J=5Hz), 7.86-7.88 (1H, m)

Preparation 28

To a suspension of phosphonium bromide (1.9 g) in tetrahydrofuran (35 ml) at 0°C was added 1.0M lithium

35

- 95 -

bis(trimethylsilyl)amide in tetrahydrofuran (7.88 ml) over 5 minutes period. After 40 minutes, the cooling bath was removed and the red suspension was stirred for 15 minutes at ambient temperature. The suspension was recooled to  
5 -78°C, and a solution of 2-[3-(phthalimido)prop-1-yl]oxybenzaldehyde (1.16 g) in 10 ml of tetrahydrofuran (plus a 5 ml rinse) was added via cannula. The red reaction mixture was stirred at 0°C to ambient temperature. After 20 hours, the solution was quenched by  
10 0.5N hydrochloric acid at 0°C. The resulting mixture was concentrated and extracted with chloroform. The organic extract was washed with brine and dried over magnesium sulfate, filtered, and concentrated to give 4-[2-[2-[3-(phthalimido)prop-1-yl]oxy]phenyl]vinyl-3-methoxybenzoic  
15 acid (2.4 g).

NMR (DMSO-d<sub>6</sub>, δ) : 1.99-2.22 (2H, m), 3.72-3.94 (5H, m), 3.98-4.17 (2H, m), 6.38-7.88 (11H, m)

#### Preparation 29

20 To a suspension of sodium hydride (60% oil suspension, 88.3 mg) in N,N-dimethylformamide (6 ml) was added a solution of methyl 4-(2-benzyloxybenzoyl)amino-3-methoxybenzoate (600 mg) in N,N-dimethylformamide (4 ml) and the mixture was stirred at 0°C for 1 hour. Methyl  
25 iodide (0.14 ml) was added dropwise to the above solution and the mixture was stirred at 0°C for 30 minutes. The reaction temperature was raised to ambient temperature over 30 minutes and the reaction was quenched with 1N hydrochloric acid, and then the resulting solution was  
30 extracted with ethyl acetate. Drying, filtering and removal of solvents afforded a crude product. The crude product was purified by column chromatography (eluent; hexane:ethyl acetate = 3:1) to give methyl 4-(N-methyl-2-benzyloxybenzoylamino)-3-methoxybenzoate (650 mg).

35 NMR (CDCl<sub>3</sub>, δ) : 3.35 (3H, s), 3.72 (3H, s), 3.87

- 96 -

(3H, s), 4.93-5.00 (2H, m); 6.65 (1H, d, J=8Hz),  
6.76 (1H, t, J=8Hz), 7.00-7.12 (2H, m), 7.18-  
7.23 (1H, m), 7.30-7.43 (6H, m), 8.02 (1H, s)  
ESI-MASS (m/z) : 406 (M+H)

5

### Preparation 30

To a solution of (S)-1,3-butanediol (1.0 g) and  
triethylamine (1.12 g) in dichloromethane (30 ml) was  
added portionwise p-toluenesulfonyl chloride (2.12 g) at  
0°C, and then the mixture was stirred at ambient  
temperature for 3 hours and stand overnight. The  
resulting solution was diluted with dichloromethane (30  
ml) and the organic layer was washed successively with 1N  
hydrochloric acid, saturated sodium bicarbonate aqueous  
solution and brine. Drying, filtering and removal of  
solvents afforded (S)-3-hydroxybutyl p-toluenesulfonate  
(2.26 g).

15

NMR (CDCl<sub>3</sub>, δ) : 1.20 (3H, d, J=8Hz), 1.63-1.77 (1H,  
m), 1.78-1.93 (1H, m), 2.47 (3H, s), 3.89-4.00  
(1H, m), 4.08-4.16 (1H, m), 4.20-4.29 (1H, m),  
7.37 (2H, d, J=9Hz), 7.80 (2H, d, J=9Hz)

20

### Preparation 31

A mixture of (S)-3-hydroxybutyl p-toluenesulfonate  
(2.25 g) and phthalimide potassium salt (3.41 g) in N,N-  
dimethylformamide (40 ml) was stirred at 60°C for 3.5  
hours. The resulting mixture was diluted with water (50  
ml) and the aqueous layer was extracted with ethyl  
acetate. Drying, filtering and removal of solvents  
afforded a crude product. The crude product was  
chromatographed on silica gel (eluent; hexane-ethyl  
acetate = 2:1) to give (S)-4-(phthalimido-1-yl)-2-butanol  
(910 mg).

30

NMR (CDCl<sub>3</sub>, δ) : 1.22 (3H, d, J=7Hz), 1.64-1.88 (2H,  
m), 2.73 (1H, d, J=4Hz), 3.68-3.78 (1H, m),

35

- 97 -

3.82-3.89 (2H, m), 7.70-7.77 (2H, m), 7.83-7.89  
(2H, m)

Preparation 32

5 To an ice-bath cooled solution of 4-methoxycarbonyl-  
phenylmethyl-tri-phenylphosphonium bromide (9.75 g) in  
N,N-dimethylacetamide (50 ml) was added potassium tert-  
butoxide (2.23 g). After being stirred in an ice-bath for  
10 30 minutes, 2-nitrobenzaldehyde (3.0 g) was added to the  
solution and the mixture was stirred at the same  
temperature for 1 hour. The mixture was diluted with  
ethyl acetate and the solution was washed with water and  
brine. The organic solution was dried over magnesium  
sulfate and the solvent was evaporated in vacuo to give an  
15 oil. The crude oil was subjected to silica gel column  
(10% ethyl acetate in n-hexane). Trans isomer was eluted  
first (1.4 g) and next cis and trans mixture (3.7 g).

Methyl 4-[(E)-2-(2-nitrophenyl)ethen-1-yl]benzoate  
20 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 3.92 (3H, s), 7.10 (1H, d, J=15Hz),  
7.41-7.50 (2H, m), 7.55-7.79 (4H, m), 8.00 (1H,  
d, J=7Hz), 8.07 (2H, d, J=8Hz)

Methyl 4-[E and Z]-2-(2-nitrophenyl)ethen-1-  
25 yl]benzoate  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 3.83 (3Hx2/3 (Z), s), 3.91 (3Hx1/3  
(E), s), 6.79 (1Hx2/3, d, J=12Hz), 6.98-8.14  
(9H+1/3H, m)

30 Preparation 33

To a solution of 3-(3-ethoxycarbonylprop-1-yl)oxy-4-  
nitrotoluene (2.67 g) in dichloromethane (30 ml) was added  
diisobutylaluminum hydride (1.5 M solution in toluene, 7  
ml) at -78°C and the solution was stirred at the same  
35 temperature for 2 hours. The reaction was quenched with

- 98 -

addition of small amount of water and a mixture of chloroform (30 ml) and 1N hydrochloric acid (20 ml) was added. The organic phase was separated and washed with water and brine. The organic solution was dried over magnesium sulfate and the solvent was evaporated in vacuo to give an oil. A mixture of the crude aldehyde and carbethoxymethylene triphenylphosphorane (3.49 g) in tetrahydrofuran (20 ml) was stirred at ambient temperature overnight and the solvent was evaporated in vacuo. The residue was subjected to silica gel column and the column was eluted with 10% ethyl acetate in n-hexane to give 3-[(E)-5-ethoxycarbonyl-4-penten-1-yl]oxy-4-nitrotoluene (2.29 g).

NMR (CDCl<sub>3</sub>, δ) : 1.27 (2H, t, J=7.5Hz), 1.93-2.04 (2H, m), 2.37 (3H, s), 2.40-2.50 (2H, m), 4.09 (2H, t, J=7.5Hz), 4.18 (2H, q, J=7.5Hz), 5.89 (1H, d, J=15Hz), 6.80 (1H, d, J=7Hz), 6.82 (1H, s), 7.00 (1H, dt, J=15, 7.5Hz), 7.78 (1H, d, J=7Hz)

#### Preparation 34

A 300 ml of hydrogenation bottle was flushed with nitrogen, and 10% palladium on carbon (1.5 g) was added into the bottle. A solution of benzyl 2-(3-phthalimidopropoxy)benzoate (1.50 g) in methanol (50 ml) and 1,4-dioxane (50 ml) was added to the bottle, along with one drop of acetic acid. The mixture was shaken in a Parr apparatus at 3 atm of hydrogen at 35°C for 8 hours. The catalyst was removed by filtration through a bed of Celite, and washed with 1,4-dioxane (20 ml x 2). The combined solution was concentrated with a rotary evaporator to give crude solid. The crude solid in methanol (57 ml) and 1,4-dioxane (10 ml) was heated and the product was recrystallized on cooling. The crystal was collected by filtration, washed with cold methanol (5



- 99 -

ml) and airdried to give 2-(3-phthalimidopropoxy)-benzoic acid (4.18 g).

mp : 155-157°C

5 NMR (DMSO-d<sub>6</sub>, δ) : 1.98-2.14 (2H, m), 3.79 (2H, t, J=7Hz), 4.08 (2H, t, J=7Hz), 6.99 (1H, dd, J=8, 8Hz), 7.08 (1H, d, J=8Hz), 7.47 (1H, m), 7.62 (1H, d, J=8Hz), 7.77-7.92 (4H, m)

### Preparation 35

10 A mixture of 4-amino-3-methoxy-N-[2-(4-carboxyphenylmethyl)oxy-4-methylphenyl]-N-methylbenzamide (500 mg), ethanolamine (109 mg), triphenylphosphine (936 mg) and carbon tetrachloride (0.57 ml) in a mixture of  
15 pyridine and acetonitrile (1:1, 15 ml) was stirred at ambient temperature for 18 hours. The solvent was evaporated and the residue was purified on silica gel column chromatography (SiO<sub>2</sub> 0-10% methanol in chloroform) to give 4-amino-3-methoxy-N-[2-[4-[N-(2-hydroxyethyl)-  
20 carbamoyl]phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide (392 mg).

NMR (CDCl<sub>3</sub>, δ) : 2.27 (3H, s), 3.33 (3H, s), 3.48 (3H, s), 3.60 (2H, q, J=5Hz), 3.78-3.84 (2H, m), 4.97 (2H, br), 6.35 (1H, d, J=8Hz), 6.61 (1H, s), 6.68-6.79 (3H, m), 7.04 (1H, d, J=8Hz), 7.11 (1H, br), 7.26 (2H, d, J=8Hz), 7.76 (2H, d, J=8Hz)  
25

### Preparation 36

To an ice-cooled 4-amino-3-methoxy-N-[2-[4-[N-(2-hydroxyethyl)carbamoyl]phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide (387 mg) was added dropwise thionyl  
30 chloride (129 mg), and the mixture was stirred at ambient temperature for 1 hour. The resulting mixture was added aqueous sodium hydrogen carbonate solution (15 ml). The  
35 solution was extracted with ethyl acetate (10 ml x 3).

- 100 -

The organic layer was washed with brine and dried over magnesium sulfate. The solvent was removed under reduced pressure to give 4-amino-3-methoxy-N-[2-[4-(2-oxazolin-2-yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide (315 mg).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.26 (3H, s), 3.35 (3H, s), 3.52 (3H, s), 4.08 (2H, t, J=10Hz), 4.25 (2H, t, J=10Hz), 4.94 (1H, br), 5.07 (1H, br), 6.40 (1H, d, J=8Hz), 6.40-6.88 (4H, m), 7.00 (1H, d, J=8Hz), 7.36 (2H, d, J=8Hz), 7.96 (2H, d, J=8Hz)

#### Preparation 37

To a solution of 3-bromopropylamine hydrobromide (5.0 g) and diisopropylethylamine (5.90 g) in dichloromethane (80 ml) was added portionwise 9-fluorenylmethoxycarbonyl chloride (5.91 g) and the mixture was stirred at ambient temperature for 3 hours and stand overnight. The resulting mixture was diluted with dichloromethane (50 ml) and the organic layer was washed successively with 1N hydrochloric acid and brine. Drying, filtering and removal of solvents afforded a crude product. The crude product was triturated with diethyl ether-hexane (1:5) to give 3-(9-fluorenylmethoxycarbonylamino)propyl bromide (7.82 g).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.02-2.12 (2H, m), 3.30-3.45 (4H, m), 4.21 (1H, t, J=8Hz), 4.44 (2H, d, J=8Hz), 4.82-4.90 (1H, br), 7.32 (2H, t, J=8Hz), 7.40 (2H, t, J=8Hz), 7.58 (2H, d, J=8Hz), 7.78 (2H, d, J=8Hz)

ESI-MASS (m/z) : 360 (M+H)

#### Preparation 38

To a solution of thiosalicylic acid (500 mg) in ethanol (15 ml) and 2N sodium hydroxide aqueous solution (3.2 ml) was added 3-(9-fluorenylmethoxycarbonylamino)-

- 101 -

propyl bromide at ambient temperature and the suspension was stirred for 2 hours. The resulting clear solution was diluted with water (20 ml) and acidified with 1N hydrochloric acid (6.5 ml). White crystals were collected by filtration and the solid was washed with ethanol-water (1:3, 15 ml) and then with n-hexane - diethyl ether (2:1, 15 ml) to give 2-[3-(9-fluorenylmethoxycarbonylamino)-propylthio]benzoic acid (1.07 g).

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.69-1.79 (2H, m), 2.90 (2H, t, J=8Hz), 3.08-3.18 (2H, m), 4.21 (1H, t, J=6Hz), 4.32 (2H, d, J=6Hz), 7.20 (1H, t, J=8Hz), 7.28-7.45 (6H, m), 7.50 (1H, t, J=8Hz), 7.68 (2H, d, J=8Hz), 7.85-7.91 (3H, m)

ESI-MASS (m/z) : 434 (M+H)

#### Example 1

To a mixture of 2-benzyloxybenzoic acid (1.17 g) and oxalyl chloride (0.536 ml) in dichloromethane (30 ml) was added 2 drops of N,N-dimethylformamide and the mixture was stirred at ambient temperature for 1 hour. After removing a solvent by evaporation, a solution of residual acid chloride in dichloromethane (5 ml) was added to a mixture of 4-amino-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide (1.97 g) and triethylamine (1.07 ml) in dichloromethane (5 ml) and the resulting solution was stirred at ambient temperature for 3 hours. The reaction mixture was washed successively with 1N hydrochloric acid, water (20 ml) and brine (20 ml), and dried over magnesium sulfate. The solvent was evaporated to give an oil and the crude product was purified by silica gel column (chloroform) to give 4-(2-benzyloxybenzoyl)amino-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide (2.89 g) as a colorless oil.

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.23 (3H, t, J=7.5Hz), 1.41-1.54

- 102 -

(2H, m), 1.63-1.75 (2H, m), 1.75-1.85 (2H, m),  
2.32 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.80-3.95  
(2H, br), 4.12 (2H, q, J=7.5Hz), 5.18 (2H, s),  
6.82-6.90 (2H, m), 6.92-7.00 (3H, m), 7.07-7.19  
5 (5H, m), 7.38-7.52 (6H, m), 8.27 (1H, d, J=7Hz)

Example 2

The following compounds were obtained according to a  
similar manner to that of Example 1.

10

- 1) 4-(2-Benzyloxybenzoyl)amino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylaminoprop-1-yloxy]-phenyl]benzamide

15

NMR (CDCl<sub>3</sub>, δ) : 2.00 (2H, m), 2.26 (3H, s),  
2.32-2.39 (4H, m), 3.32 (3H, s), 3.34-3.41 (6H,  
m), 3.81-4.02 (2H, m), 5.20 (2H, s), 6.78-7.26  
(9H, m), 7.38-7.53 (7H, m), 8.27 (1H, d, J=7Hz)

20

- 2) 3-Methoxy-4-(2-nitrobenzoyl)amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

25

NMR (CDCl<sub>3</sub>, δ) : 1.43-1.60 (2H, m), 1.61-1.90 (4H,  
m), 2.30 (6H, s), 2.31-2.44 (6H, m), 3.33 (3H,  
s), 3.44-3.53 (2H, m), 3.57-3.67 (2H, m), 3.71  
(3H, s), 3.81-4.03 (2H, m), 6.56-6.69 (2H, m),  
6.82-6.99 (2H, m), 7.03 (1H, s), 7.57-7.66 (2H,  
m), 7.67-7.76 (1H, m), 8.02-8.13 (2H, m), 8.21  
(1H, d, J=8Hz)

30

- 3) 4-(2-Methoxybenzoyl)amino-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]benzamide

35

NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7Hz), 1.43-1.59 (2H,  
m), 1.63-1.90 (4H, m), 2.26 (3H, s), 2.34 (2H,  
t, J=7Hz), 3.32 (3H, s), 3.79-3.99 (2H, m), 4.02  
(3H, s), 4.11 (2H, q, J=7Hz), 6.53-6.66 (2H, m),

- 103 -

6.87 (1H, d, J=8Hz), 7.01 (1H, d, J=8Hz), 7.12 (1H, dd, J=8, 8Hz), 7.29-7.40 (2H, m), 7.42-7.56 (3H, m), 8.18-8.28 (1H, m), 9.81 (1H, br s)

- 5      4) 4-(2-Benzyloxybenzoyl)amino-3-methoxy-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7Hz), 1.42-1.58 (2H, m), 1.62-1.90 (4H, m), 2.32 (2H, t, J=7Hz), 3.28 (3H, s), 3.33 (3H, s), 3.78-4.03 (2H, m), 4.12 (2H, q, J=7Hz), 5.30 (2H, s), 6.72-7.22 (8H, m), 7.28-7.55 (6H, m), 8.20-8.29 (1H, m), 8.38 (1H, d, J=8Hz)
- 10
- 15      5) 4-[2-(Acetyloxy)benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.32-2.01 (10H, m), 2.21-2.46 (15H, m), 1.57 (1H, m), 3.02 (1H, m), 3.32 (3H, s), 3.79 (3H, s), 3.83-4.03 (3H, m), 4.69 (1H, m), 6.54-6.67 (2H, m), 6.80-8.33 (8H, m)
- 20
- 25      6) 4-[2-(Acetyloxy)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7Hz), 1.44-1.91 (6H, m), 2.21-2.41 (8H, m), 3.32 (3H, s), 3.80 (3H, s), 3.82-4.03 (2H, m), 6.54-6.67 (2H, m), 6.80-6.95 (2H, m), 7.07 (1H, s), 7.15 (1H, d, J=8Hz), 7.35 (1H, m), 7.51 (1H, m), 7.94 (1H, m), 8.28 (1H, d, J=8Hz), 8.87 (1H, s)
- 30
- 35      7) 4-(2-Benzyloxybenzoyl)amino-2-chloro-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7Hz), 1.47-1.98 (6H, m), 2.36 (2H, t, J=7Hz), 3.34 (3H, s), 3.96 (2H,

- 104 -

t, J=7Hz), 4.14 (2H, q, J=7Hz), 5.17 (2H, s),  
6.64-6.82 (3H, m), 6.96 (1H, d, J=8Hz), 7.02-  
7.21 (5H, m), 7.41-7.62 (6H, m), 8.25 (1H, m)

- 5      8) 4-(2-Acetoxybenzoyl)amino-3-methoxy-N-methyl-N-(2-methylphenyl)benzamide

NMR (CDCl<sub>3</sub>, δ) : 2.21 (3H, s), 2.31 (3H, s), 3.38  
(3H, s), 3.73 (3H, s), 6.87 (1H, d, J=8Hz), 7.00  
(1H, s), 7.03-7.24 (5H, m), 7.29-7.43 (1H, m),  
10      7.51 (1H, dd, J=8, 8Hz), 7.92 (1H, d, J=8Hz),  
8.30 (1H, d, J=8Hz), 8.87 (1H, br s)

- 15      9) 4-(3-Benzyloxybenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.32-1.42 (2H, m), 1.50-1.58 (2H,  
m), 1.67-1.90 (6H, m), 2.28 (3H, s), 2.29 (6H,  
s), 2.37 (2H, t, J=8Hz), 2.52-2.62 (1H, m),  
2.98-3.07 (1H, m), 3.34 (3H, s), 3.78 (3H, s),  
20      3.85-3.98 (3H, m), 4.59-4.67 (1H, m), 5.12 (2H,  
s), 6.58 (1H, d, J=8Hz), 6.63 (1H, d, J=8Hz),  
6.84 (1H, d, J=8Hz), 6.92 (1H, d, J=8Hz), 7.02  
(1H, s), 7.12-7.17 (1H, m), 7.33-7.50 (8H, m),  
8.28 (1H, d, J=8Hz), 8.48 (1H, s)

25      ESI-MASS (m/z) : 721 (M+H)

- 30      10) 4-(2-Benzyloxybenzoyl)amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7.5Hz), 1.43-1.56  
(2H, m), 1.65-1.84 (4H, m), 2.25 (3H, s), 2.32  
(2H, t, J=7.5Hz), 3.29 (3H, s), 3.77-3.93 (2H,  
m), 4.12 (2H, q, J=7.5Hz), 5.19 (2H, s), 6.51  
(2H, m), 6.81 (1H, d, J=7Hz), 6.98 (2H, d,  
J=8Hz), 7.07-7.19 (4H, m), 7.39-7.53 (6H, m),  
35      8.27 (1H, d, J=7Hz)

- 105 -

- 11) 4-(2-Iodobenzoyl)amino-N-[2-(4-methoxyphenyl)-methoxy]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 3.35 (3H, s), 3.82 (3H, s), 4.90-5.05 (2H, m), 6.83 (1H, t, J=7Hz), 6.89-6.96 (3H, m), 7.04 (1H, d, J=7Hz), 7.10-7.19 (2H, m), 7.22-7.32 (4H, m), 7.37-7.48 (3H, m), 7.53 (1H, s), 7.88 (1H, d, J=7Hz)
- 12) 3-Methoxy-4-[2-(4-methoxyphenylmethyl)oxybenzoyl]-amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.27 (3H, s), 2.31 (3H, s), 2.35-2.52 (2H, m), 3.24 (3H, s), 3.37 (3H, s), 3.40-3.53 (2H, m), 3.62-3.81 (2H, m), 3.39 (3H, s), 4.89 (1H, d, J=14Hz), 5.06 (1H, d, J=14Hz), 5.21 (2H, s), 6.61-6.70 (2H, m), 6.80-7.18 (7H, m), 7.30-7.45 (7H, m), 8.22 (1H, d, J=7Hz), 8.31 (1H, d, J=7Hz)
- 13) 4-[2-(E)-(2-Ethoxycarbonyl)ethen-1-yl]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.30 (3H, t, J=7.5Hz), 1.49-1.60 (2H, m), 1.67-1.77 (2H, m), 1.79-1.90 (2H, m), 2.29 (6H, sx2), 2.33-2.43 (6H, m), 3.33 (3H, s), 3.45-3.53 (2H, m), 3.60-3.67 (2H, m), 3.71 (3H, s), 3.85-4.01 (2H, m), 4.23 (2H, q, J=7.5Hz), 6.40 (1H, d, J=15Hz), 6.58-6.67 (2H, m), 6.86 (1H, d, J=7Hz), 6.92 (1H, d, J=7Hz), 7.02 (1H, s), 7.40-7.52 (2H, m), 7.58 (1H, d, J=7Hz), 7.68 (1H, d, J=7Hz), 8.02-8.15 (2H, m), 8.27 (1H, d, J=7Hz)
- 14) 4-(2-Dimethylamino-4-methyl)phenoxyethyl-N-[2-(5-ethoxycarbonyl)pent-1-yl]oxy-4-methyl]phenylbenzamide

- 106 -

5 NMR (CDCl<sub>3</sub>, δ) : 1.20 (3H, t, J=7.5Hz), 1.45-1.58 (2H, m), 1.66-1.77 (2H, m), 1.80-1.95 (2H, m), 2.25 (3H, s), 2.29-2.34 (2H, m), 2.31 (3H, s), 2.80 (6H, s), 4.00-4.16 (4H, m), 5.20 (2H, s), 6.68-6.89 (5H, m), 7.58 (2H, d, J=8Hz), 7.88 (2H, d, J=8Hz), 8.37 (1H, d, J=7Hz), 8.50 (1H, s)

10 15) 4-(2-Benzyloxy)benzoylamino-3-methoxy-N-[(E and Z)-2-(4-methoxycarbonylphenyl)ethen-1-yl]phenyl-N-methylbenzamide

15 NMR (CDCl<sub>3</sub>, δ) : 3.06 (3Hx2/3, s), 3.10 (3Hx1/3, s), 3.40 (3Hx2/3, s), 3.43 (3Hx1/3, s), 3.46 (3Hx2/3, s), 3.91 (3Hx1/3, s), 5.20 (2Hx2/3, s), 5.27 (2Hx1/3, s), 6.38-8.37 (22H, m)

20 16) 3-Methoxy-4-[2-[3-(4-methoxyphenyl)methoxyprop-1-yl]thiobenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]-phenylbenzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.44-1.58 (2H, m), 1.61-1.73 (2H, m), 1.68-1.92 (4H, m), 2.25 (3H, s), 2.27 (3H, s), 2.30-2.41 (6H, m), 2.99 (2H, t, J=7.5Hz), 3.30 (3H, s), 3.43-3.52 (4H, m), 3.57-3.66 (2H, m), 3.70 (3H, s), 3.78 (3H, s), 3.82-3.90 (2H, m), 4.38 (2H, s), 6.53-6.65 (2H, m), 6.79-6.93 (3H, m), 7.02 (1H, s), 7.17-7.29 (4H, m), 7.33-7.45 (2H, m), 7.65 (1H, d, J=7Hz), 8.29 (1H, d, J=7Hz), 8.80 (1H, s)

30 17) 4-(2,4-Dimethoxybenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

35 NMR (CDCl<sub>3</sub>, δ) : 1.43-1.57 (2H, m), 1.64-1.72 (2H, m), 1.72-1.91 (2H, m), 2.24 (3H, s), 2.27 (3H, s)



- 107 -

- 5 s), 2.30-2.40 (6H, m), 3.31 (3H, s), 3.42-3.50 (2H, m), 3.59-3.65 (2H, m), 3.77 (3H, s), 3.80 (3H, s), 3.80-4.02 (2H, m), 3.96 (3H, s), 6.52-6.63 (2H, m), 6.81-7.04 (5H, m), 7.79 (1H, m), 8.38 (1H, d, J=7Hz)
- 10 18) 4-[2-(Acetoxy)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide
- 15 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.61 (2H, m), 1.64-1.93 (4H, m), 2.22-2.46 (15H, m), 3.33 (3H, s), 3.44-3.53 (2H, m), 3.58-3.68 (2H, m), 3.79 (3H, s), 3.82-4.04 (2H, m), 6.54-6.68 (2H, m), 6.80-6.95 (2H, m), 7.04 (1H, s), 7.14 (1H, d, J=8Hz), 7.35 (1H, m), 7.51 (1H, m), 7.92 (1H, m), 8.29 (1H, br d, J=8Hz), 8.86 (1H, s)
- 20 19) 4-(2-Benzyloxy-4-methylbenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide
- 25 NMR (CDCl<sub>3</sub>, δ) : 1.48-1.61 (2H, m), 1.69-1.91 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.44 (6H, m), 2.38 (3H, s), 3.20 (3H, s), 3.32 (3H, s), 3.50 (2H, t, J=5Hz), 3.64 (2H, t, J=4Hz), 3.85-4.06 (2H, m), 4.89 (2H, s), 6.60-6.68 (2H, m), 6.82-6.95 (4H, m), 7.18 (1H, dd, J=2, 7Hz), 7.27-7.40 (5H, m), 7.98 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)
- 30 20) 4-(2-Benzyloxy-4-methylbenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide
- 35 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.86 (6H, m), 2.28 (3H, s), 2.30 (3H, s), 2.36 (3H, s), 2.32-2.48 (6H, m), 3.30 (3H, s), 3.45-3.51 (2H, m), 3.60-3.66 (2H,

- 108 -

m), 3.63 (3H, s), 3.79-4.00 (2H, m), 5.24 (2H, d, J=9Hz), 6.56-6.68 (2H, m), 6.80-6.93 (5H, m), 7.31-7.58 (5H, m), 8.11 (1H, d, J=8Hz), 8.36 (1H, d, J=8Hz)

5

21) 4-(2-Benzyloxy-5-methylbenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide

10

NMR (CDCl<sub>3</sub>, δ) : 1.47-1.99 (8H, m), 2.28 (3H, s), 2.31 (3H, s), 2.31-2.45 (6H, m), 3.25 (3H, s), 3.29 (2H, s), 3.48 (3H, s), 3.48-3.53 (2H, m), 3.60-3.64 (2H, m), 3.82-4.01 (2H, m), 5.27 (2H, s), 6.54-6.63 (2H, m), 6.81-6.95 (4H, m), 7.19-7.47 (7H, m), 8.02 (1H, s), 8.36 (1H, d, J=8Hz)

15

22) 4-(2-Benzyloxy-4-chlorobenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide

20

NMR (CDCl<sub>3</sub>, δ) : 1.46-1.86 (6H, m), 2.15-2.31 (2H, m), 2.29 (6H, s), 2.30-2.58 (4H, m), 3.28 (3H, s), 3.49 (2H, t, J=5Hz), 3.60 (3H, s), 3.61 (2H, t, J=5Hz), 3.85-4.00 (2H, m), 5.15 (2H, s), 6.54-6.67 (2H, m), 6.83-7.16 (4H, m), 7.34-7.49 (7H, m), 8.01 (1H, d, J=8Hz), 8.19 (1H, d, J=8Hz)

25

23) 4-(2-Benzyloxy-4-methoxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide

30

NMR (CDCl<sub>3</sub>, δ) : 1.44-1.59 (2H, m), 1.63-1.84 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.25-2.40 (6H, m), 3.28 (3H, s), 3.30 (3H, s), 3.48 (2H, t, J=4Hz), 3.62 (2H, t, J=4Hz), 3.89-4.01 (2H, m), 5.26 (2H, s), 6.52-6.67 (4H, m), 6.81-6.92 (4H, m), 7.35-7.48 (5H, m), 8.21 (1H, d, J=9Hz), 8.38

35

- 109 -

(1H, d, J=8Hz)

- 24) 4-(2-Acetoxybenzoyl)amino-3-methoxy-N-(2-benzyloxy-4-methylphenyl)-N-methylbenzamide

5 NMR (CDCl<sub>3</sub>, δ) : 2.29 (3H, s), 3.39 (3H, s), 3.60 (3H, s), 4.88 (1H, d, J=12Hz), 5.02 (1H, d, J=12Hz), 6.68-6.73 (2H, m), 6.82 (1H, d, J=8Hz), 7.02 (1H, s), 7.11-7.20 (2H, m), 7.31-7.42 (5H, m), 7.46-7.53 (1H, m), 7.93 (1H, d, J=8Hz), 8.27 (1H, d, J=8Hz), 8.86 (1H, br)

- 25) 4-(2-Acetoxybenzoyl)amino-3-methoxy-N-[2-[4-(2-oxazolin-2-yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide

15 NMR (CDCl<sub>3</sub>, δ) : 2.27 (3H, s), 2.31 (3H, s), 3.39 (3H, s), 3.63 (3H, s), 4.07 (2H, t, J=10Hz), 4.42 (2H, t, J=10Hz), 4.91 (1H, d, J=12Hz), 5.11 (1H, d, J=12Hz), 6.61 (1H, br), 6.77 (1H, d, J=8Hz), 6.82-7.15 (5H, m), 7.24-7.50 (4H, m), 7.90 (2H, d, J=8Hz), 8.20 (1H, d, J=8Hz)

- 26) 4-[2-[3-(9-Fluorenylmethyl)oxycarbonylaminoprop-1-yl]thiobenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.32-1.92 (12H, m), 2.29 (9H, s), 2.39 (2H, t, J=5Hz), 2.60 (1H, t, J=10Hz), 2.90-3.12 (3H, m), 3.29 (2H, q, J=5Hz), 3.33 (3H, s), 3.75 (3H, s), 3.82-4.00 (4H, m), 4.38 (2H, t, J=4Hz), 6.55-6.67 (3H, m), 6.83 (1H, d, J=8Hz), 6.92 (1H, d, J=8Hz), 7.02 (1H, s), 7.25-7.46 (6H, m), 7.59 (2H, d, J=7Hz), 7.63 (1H, d, J=8Hz), 7.77 (2H, d, J=8Hz), 8.30 (1H, d, J=8Hz), 8.70 (1H, s)

35

- 110 -

- 27) 4-[2-(Acetyloxy)benzoyl]amino-3-methyl-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.53 (2H, br), 1.63-1.89 (4H, m),  
2.22 (3H, s), 2.30 (3H, s), 2.36 (3H, s), 2.22-  
2.50 (10H, m), 3.32-3.38 (3H, m), 3.52-3.57 (2H,  
m), 3.67 (2H, br), 3.95 (2H, br), 6.61 (2H, s),  
6.83-6.93 (2H, m), 7.02-7.20 (2H, m), 7.32-7.58  
10 (2H, m), 7.68 (1H, d, J=7Hz), 7.85 (1H, br)

- 28) 4-[(2-Benzyloxy)benzoyl]amino-3-[(2-benzyloxy)-benzoyl]oxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.44-1.53 (2H, m), 1.60-1.87 (4H,  
m), 2.28 (3H, s), 2.29 (3H, s), 2.32-2.38 (7H,  
m), 3.33 (3H, s), 3.43 (2H, br), 3.60 (2H, br),  
3.90 (2H, br), 4.79 (2H, s), 4.93 (2H, s), 6.11-  
6.20 (3H, m), 6.82-7.43 (18H, m), 7.83-7.88 (1H,  
m), 8.12-8.15 (1H, m), 8.37-8.42 (1H, m)

- 29) 4-[4-(Benzyloxy)benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.30-1.45 (1H, m), 1.47-1.58 (2H,  
m), 1.60-1.75 (4H, m), 1.78-1.91 (2H, m), 2.27  
(9H, s), 2.30-2.40 (3H, m), 2.50-2.63 (1H, m),  
2.95-3.07 (1H, m), 3.30 (3H, s), 3.77 (3H, s),  
3.82-3.98 (4H, m), 4.56-4.67 (1H, m), 5.11 (2H,  
s), 6.56-6.62 (2H, m), 6.80-6.93 (2H, m), 7.00-  
30 7.05 (3H, m), 7.34-7.45 (4H, m), 7.78-7.82 (2H,  
m), 8.22-8.30 (1H, m), 8.46 (1H, s)

- 30) 4-[4-(Benzyloxy)benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

35

- 111 -

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.59 (2H, m), 1.69-1.90 (4H, m), 2.28 (3H, s), 2.30 (3H, s), 2.35-2.42 (6H, m), 3.31 (3H, s), 3.48-3.50 (2H, m), 3.62-3.66 (2H, m), 3.78 (3H, s), 3.82-4.00 (2H, m), 5.13 (2H, s), 6.57-6.60 (2H, m), 6.81-6.92 (2H, m), 7.00-7.02 (3H, m), 7.30-7.43 (5H, m), 7.78-7.82 (2H, m), 8.27 (1H, d, J=7Hz), 8.43 (1H, s)

31) 4-[2-(Benzyloxy)benzoyl]amino-2-nitro-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.30-1.44 (2H, m), 1.50-1.94 (8H, m), 2.20 (3H, s), 2.27 (6H, s), 2.30-2.43 (3H, m), 2.52-2.63 (1H, m), 2.97-3.10 (1H, m), 3.32 (3H, s), 3.85-3.97 (4H, m), 4.57-4.68 (1H, m), 5.20 (2H, s), 6.41-6.48 (2H, m), 6.52 (1H, s), 6.90-6.93 (1H, m), 7.11-7.20 (3H, m), 7.32 (1H, s), 7.48-7.59 (6H, m), 7.69-7.73 (1H, m), 8.29 (1H, d, J=7Hz)

32) 2-[2-(Benzyloxy)benzoyl]amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-5-pyridinecarboxamide

NMR (CDCl<sub>3</sub>, δ) : 1.30-1.44 (2H, m), 1.44-1.60 (2H, m), 1.60-1.95 (6H, m), 2.20 and 2.28 (total 9H, s), 2.29-2.41 (3H, m), 2.47-2.64 (1H, m), 2.93-3.09 (1H, m), 3.32 (3H, s), 3.79-3.98 (4H, m), 4.57-4.69 (1H, m), 4.97-5.17 (1H, m), 5.32 (1H, s), 6.39-6.50 (1H, m), 6.60-6.78 (2H, m), 6.85-6.90 (1H, m), 7.00-7.12 (2H, m), 7.27-7.50 (7H, m), 7.56-8.25 (2H, m)

### Example 3

To a mixture of 2-benzyloxybenzoic acid (1.55 g) and oxalyl chloride (1.18 ml) in dichloromethane (30 ml) was

- 112 -

added 1 drop of N,N-dimethylformamide and the mixture was stirred at ambient temperature for 1 hour. After removing a solvent by evaporation, a solution of residual acid chloride in dichloromethane (30 ml) was added to a mixture of 4-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yloxy]phenyl]benzamide (3.28 g) and pyridine (1.1 ml) in dichloromethane (50 ml) and the mixture was stirred at ambient temperature for 2.5 hours. The mixture was washed with saturated sodium hydrogen carbonate solution and brine, and dried over sodium sulfate. The solvent was removed by evaporation and purified by silica gel column chromatography (SiO<sub>2</sub>; 85 g, 2% methanol in dichloromethane) to give 4-(2-benzyloxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yloxy]-4-methylphenyl]benzamide (4.5 g).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.44-1.59 (2H, m), 1.62-1.90 (4H, m), 2.27 (3H, s), 2.28 (3H, s), 2.30-2.43 (6H, m), 3.30 (3H, s), 3.32 (3H, s), 3.43-3.53 (2H, m), 3.57-3.67 (2H, m), 3.78-4.03 (2H, m), 5.30 (2H, s), 6.52-6.66 (2H, m), 6.78-6.96 (3H, m), 7.04 (1H, d, J=9Hz), 7.10 (1H, dd, J=9, 9Hz), 7.30-7.49 (6H, m), 8.20-8.28 (1H, m), 8.37 (1H, d, J=9Hz)

#### Example 4

A solution of 4-(2-benzyloxybenzoyl)amino-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide (2.80 g) in a mixture of ethanol (50 ml) and 1N sodium hydroxide solution (10 ml) was stirred at ambient temperature for 4 hours. After removing ethanol by evaporation, the aqueous solution was adjusted to pH 2 with 1N hydrochloric acid and the mixture was extracted with chloroform (30 x 2). The organic phase was washed with water (40 ml) and brine (30 ml), and dried over magnesium sulfate. The solvent

- 113 -

was evaporated to give 4-(2-benzyloxybenzoyl)-  
amino-N-methyl-N-[2-(5-carboxypent-1-yloxy)phenyl]-  
benzamide (1.76 g) as a colorless oil.

5 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.57 (2H, m), 1.66-1.83 (4H,  
m), 2.37 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.78-  
3.96 (2H, br), 5.17 (2H, s), 6.75-6.82 (2H, m),  
6.93-7.02 (3H, m), 7.10-7.22 (5H, m), 7.36-7.51  
(6H, m), 8.28 (1H, d, J=7Hz)

10 Example 5

The following compounds were obtained according to a  
similar manner to that of Example 4.

15 1) 4-[2-(Carboxymethoxy)benzoyl]amino-N-methyl-N-[2-[5-  
(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-  
benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.63 (2H, m), 1.73 (2H, m), 1.85  
(2H, m), 2.28 (3H, s), 2.35-2.41 (6H, m), 3.36  
(3H, s), 3.47 (2H, m), 3.61 (2H, m), 3.91 (2H,  
m), 4.76 (2H, s), 6.72-6.82 (2H, m), 6.86-7.01  
(2H, m), 7.07-7.18 (2H, m), 7.35 (2H, d,  
J=8.5Hz), 7.47 (1H, t, J=7Hz), 7.72 (2H, d,  
J=8.5Hz), 8.25 (1H, d, J=7Hz)

25 2) 4-(2-Aminobenzoyl)amino-N-methyl-N-[2-(5-carboxypent-  
1-yloxy)-4-methylphenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.59 (2H, m), 1.64-1.85 (4H,  
m), 2.27 (3H, s), 2.38 (2H, t, J=7Hz), 3.32 (3H,  
s), 3.73-4.00 (2H, m), 6.56-6.76 (4H, m), 6.93  
(1H, d, J=9Hz), 7.18-7.48 (6H, m), 7.86 (1H, br  
s)

35 3) 4-(2-Methoxybenzoyl)amino-N-methyl-N-[2-(5-  
carboxypent-1-yloxy)-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.46-1.62 (2H, m), 1.65-1.88 (4H,

- 114 -

5 m), 2.26 (3H, s), 2.39 (2H, t, J=7Hz), 3.33 (3H, s), 3.73-4.00 (2H, m), 4.01 (3H, s), 6.54-6.68 (2H, m), 6.91 (1H, br d, J=9Hz), 6.99 (1H, d, J=9Hz), 7.10 (1H, dd, J=9, 9Hz), 7.35 (2H, br d, J=9Hz), 7.41-7.57 (3H, m), 8.17-8.27 (1H, m), 9.84 (1H, br s)

10 4) 4-(2-Benzyloxybenzoyl)amino-3-methoxy-N-methyl-N-[2-(5-carboxypent-1-yloxy)phenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.43-1.60 (2H, m), 1.62-1.88 (4H, m), 2.38 (2H, t, J=7Hz), 3.28 (3H, s), 3.34 (3H, s), 3.76-4.02 (2H, m), 5.28 (2H, s), 6.74-6.85 (2H, m), 6.86-6.97 (2H, m), 6.97-7.20 (4H, m), 7.28-7.50 (6H, m), 8.16-8.27 (1H, m), 8.36 (1H, d, J=8Hz)

20 5) 4-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl]oxy]benzoyl]amino-3-methoxy-N-methyl-N-[2-(5-carboxypent-1-yloxy)-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.31-1.96 (17H, m), 2.00-2.48 (6H, m), 3.14-3.39 (5H, m), 3.62-4.07 (5H, m), 4.10-4.30 (2H, m), 4.86 (1H, m), 6.52-6.72 (2H, m), 6.81-7.16 (5H, m), 7.37-7.53 (2H, m), 8.11-8.51 (2H, m)

25 6) 4-(2-Benzyloxybenzoyl)amino-2-chloro-N-methyl-N-[2-(5-carboxypent-1-yloxy)phenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.50-1.67 (2H, m), 1.68-1.98 (4H, m), 2.42 (2H, t, J=7Hz), 3.34 (3H, s), 3.99 (2H, t, J=7Hz), 5.16 (2H, s), 6.65-6.80 (3H, m), 6.98 (1H, d, J=8Hz), 7.02-7.22 (5H, m), 7.40-7.61 (6H, m), 8.24 (1H, m)

30 7) 4-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl]oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-(5-carboxypent-



- 115 -

1-yloxy)phenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 1.45-1.80 (8H, m),  
2.18-2.27 (2H, m), 2.32-2.40 (2H, m), 3.25-3.35  
(2H, m), 3.48 (3H, s), 3.80 (3H, s), 3.93 (2H,  
t, J=6Hz), 4.19-4.28 (2H, m), 4.73-4.83 (1H,  
br), 6.73-6.80 (3H, m), 6.93-7.12 (6H, m), 7.46  
(1H, t, J=8Hz), 8.17-8.27 (1H, m)

ESI-MASS (m/z) : 686 (M+Na)

- 10 8) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-[2-(5-carboxypent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.31-1.80 (6H, m), 1.95-2.07 (4H, m), 2.22 (3H, s), 2.86 (2H, t, J=7.5Hz), 3.16 (3H, s), 3.70 (1H, m), 3.93 (1H, m), 4.16 (2H, t, J=7.5Hz), 6.65 (1H, d, J=7Hz), 6.78 (1H, s), 7.00-7.10 (2H, m), 7.20 (1H, d, J=7Hz), 7.23 (2H, d, J=8Hz), 7.43-7.62 (4H, m)

- 20 9) 4-[2-[3-(tert-Butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-N-[2-(5-carboxypent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.36-1.50 (2H, m), 1.41 (9H, s), 1.50-1.62 (2H, m), 1.66-1.84 (2H, m), 2.05-2.19 (2H, m), 2.25 (3H, s), 2.36-2.44 (2H, m), 3.23-3.41 (2H, m), 3.31 (3H, s), 3.77-4.00 (2H, m), 4.16-4.29 (2H, m), 4.88 (1H, br), 6.53-6.67 (2H, m), 6.98 (2H, d, J=8Hz), 7.08 (1H, m), 7.30-7.53 (3H, m), 8.11 (1H, m)

- 30 10) 4-[(2-Benzyloxy)benzoyl]amino-N-[2-(3-carboxyprop-1-yl)oxy]phenyl-N-methylbenzamide

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.90-2.01 (2H, m), 2.42 (2H, t, J=7.5Hz), 3.20 (3H, s), 3.85-4.02 (2H, m), 5.20 (2H, s), 6.85 (1H, t, J=7Hz), 6.98 (1H, d,

35

- 116 -

J=7Hz), 7.09 (1H, t, J=7Hz), 7.15-7.37 (6H, m),  
7.49 (2H, d, J=8Hz), 7.62 (1H, d, J=7Hz)

- 5 11) 4-(2-Iodobenzoyl)amino-N-[2-(5-carboxypent-1-yl)oxy]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.45-1.58 (2H, m), 2.65-2.75 (2H, m), 2.75-2.84 (2H, m), 2.35 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.82-3.98 (2H, m), 6.77-6.86 (2H, m), 7.04 (1H, d, J=7Hz), 7.09-7.21 (2H, m), 7.28-7.48 (5H, m), 7.82-7.90 (2H, m)
- 15 12) 4-(2-Dimethylamino-4-methyl)phenoxyethyl-N-[2-(5-carboxypent-1-yl)oxy]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.38-1.52 (2H, m), 1.59-1.69 (2H, m), 1.72-1.85 (2H, m), 2.23 (3H, s), 2.25 (3H, s), 2.30 (2H, t, J=7.5Hz), 2.75 (6H, s), 3.33 (3H, s), 3.11-3.25 (2H, m), 3.88-4.00 (2H, m), 5.02 (2H, s), 6.56-6.67 (3H, m), 6.71 (1H, d, J=7Hz), 6.90-6.99 (2H, m), 7.24 (2H, d, J=8Hz), 7.38 (2H, d, J=8Hz)
- 20 13) 3-Methoxy-4-[2-[1-(tert-butoxycarbonyl)piperidin-4-yl]oxybenzoyl]amino-N-[2-(5-carboxypent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.40-1.57 (2H, m), 1.45 (9H, s), 1.60-1.94 (6H, m), 2.01-2.22 (2H, m), 2.29 (3H, s), 2.38 (2H, t, J=7.5Hz), 2.97-3.20 (2H, m), 3.33 (3H, s), 3.41 (2H, t, J=7.5Hz), 3.71 (3H, s), 3.78-4.00 (2H, m), 4.67 (1H, m), 6.60-6.65 (2H, m), 6.87-7.12 (5H, m), 7.44 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)
- 30 14) 3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)amino-1-methylprop-1-yl]oxybenzoyl]amino-N-[2-(5-carboxypent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide
- 35

- 117 -

- 5 NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.42 (3H, d, J=7.5Hz), 1.43-1.96 (8H, m), 2.25 (3H, s), 2.33-2.42 (2H, m), 3.11-3.33 (2H, m), 3.33 (3H, s), 3.65-3.97 (5H, m), 4.70 (1H, m), 6.53-6.70 (2H, m), 6.79-7.13 (4H, m), 7.44 (1H, t, J=7Hz), 8.23 (1H, m), 8.39 (1H, m)
- 10 15) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-[2-(3-carboxypyrid-6-yl)methoxy-4-methylphenyl]-N-methylbenzamide
- 15 NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 2.05-2.16 (2H, m), 2.27 (3H, s), 3.28 (2H, br), 3.42 (3H, br), 3.58 (3H, br), 3.86-4.00 (2H, m), 4.10-4.25 (2H, m), 4.95 (1H, br), 5.16 (1H, br), 6.62 (3H, br), 6.86-7.18 (4H, m), 7.41 (3H, br), 8.14 (1H, br), 8.33 (1H, br), 9.17 (1H, br)
- 20 16) 4-[2-(E)-(2-Carboxyethen-1-yl)benzoylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenyl]benzamide
- 25 NMR (CDCl<sub>3</sub>, δ) : 1.50-2.00 (6H, m), 2.27-2.52 (10H, m), 2.60-2.81 (2H, m), 3.31 (3H, s), 3.43-3.66 (2H, m), 3.83-4.22 (7H, m), 5.60 (1H, m), 6.57 (1H, m), 6.65-6.76 (4H, m), 7.01-7.12 (2H, m), 7.21 (1H, d, J=7Hz), 7.42-7.60 (3H, m), 7.85 (1H, m)
- 30 17) 4-[2-(3-Carboxyprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide
- 35 NMR (CDCl<sub>3</sub>, δ) : 1.44-1.57 (2H, m), 1.64-1.75 (2H, m), 1.75-1.87 (2H, m), 2.20 (3H, s), 2.34 (3H, s), 2.35-2.50 (6H, m), 2.61-2.74 (2H, m), 3.30 (3H, s), 3.33-3.46 (2H, m), 3.49-3.69 (4H, m), 3.75 (3H, s), 3.90-4.02 (2H, m), 4.17-4.27 (2H,

- 118 -

m), 6.56-6.72 (2H, m), 6.83-6.92 (2H, m), 6.93-7.00 (2H, m), 7.07 (1H, t, J=7Hz), 7.43 (1H, t, J=7Hz), 7.43 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

5

18) 4-[2-(Carboxymethoxy)benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

10

NMR (CDCl<sub>3</sub>, δ) : 1.51-1.92 (6H, m), 2.02 (3H, s), 2.30 (3H, s), 2.32 (2H, t, J=5Hz), 2.43-2.68 (4H, m), 3.33 (3H, s), 3.40-3.55 (4H, m), 3.72 (3H, s), 3.75-4.07 (2H, m), 4.73 (2H, s), 6.57-6.68 (2H, m), 6.81-7.10 (6H, m), 7.35-7.45 (1H, m), 8.18 (1H, d, J=7Hz), 8.32 (1H, d, J=8Hz)

15

#### Example 6

A mixture of 4-[2-[3-(phthalimido)prop-1-yl]oxy]-benzoylamino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (470 mg) and hydrazine hydrate (158 mg) in ethanol (5 ml) was stirred at ambient temperature for 6 hours and filtered through a bed of Celite. The filtrate was evaporated and the residue was subjected to silica gel column. The column was eluted with a mixture of chloroform, methanol and aqueous ammonia (10:1:0.1). The object fractions were evaporated to give 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]-amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide (256 mg) as a colorless amorphous.

20

25

30

35

NMR (CDCl<sub>3</sub>, δ) : 1.56 (2H, m), 1.74 (2H, m), 1.87 (2H, m), 2.09 (2H, m), 2.29 (3H, s), 2.34-2.43 (6H, m), 2.97 (2H, t, J=7.5Hz), 3.33 (3H, s), 3.50 (2H, m), 3.65 (2H, m), 3.96 (2H, m), 4.30 (2H, t, J=7.5Hz), 6.73-6.83 (2H, m), 6.95-7.03 (2H, m), 7.77-7.16 (2H, m), 7.34 (2H, d,

- 119 -

J=8.5Hz), 7.42-7.50 (3H, m), 8.22 (1H, d, J=7Hz)

Example 7

The following compounds were obtained according to a similar manner to that of Example 6.

- 1) 4-[2-(3-Aminoprop-1-yl)oxy]benzoylamino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylamino]prop-1-yloxy]phenyl]benzamide

10 NMR (CDCl<sub>3</sub>, δ) : 2.00 (2H, m), 2.10 (2H, m), 2.27 (3H, s), 2.34-2.39 (4H, m), 2.98 (2H, t, J=7.5Hz), 3.35 (3H, s), 3.35-3.61 (6H, m), 3.98 (2H, m), 4.30 (2H, t, J=7.5Hz), 6.80-6.91 (2H, m), 7.02 (2H, d, J=7Hz), 7.07-7.21 (3H, m),  
15 7.33-7.51 (5H, m), 8.22 (1H, d, J=7Hz)

- 2) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.53 (2H, m), 1.70 (2H, m), 1.84 (2H, m), 2.07 (2H, m), 2.26 (3H, s), 2.28 (3H, s), 2.31-2.40 (6H, m), 2.90 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.49 (2H, m), 3.60 (2H, m), 3.89 (3H, s), 3.82-3.99 (2H, m), 4.28 (2H, t, J=7.5Hz), 6.54-6.64 (2H, m), 6.82-6.94 (2H, m),  
25 7.00-7.11 (3H, m), 7.45 (1H, m), 8.20 (1H, m), 8.39 (1H, m)

- 3) (R)-4-[2-[(4-Aminobut-2-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.42 and 1.45 (total 3H, s), 1.50-1.89 (8H, m), 2.02-2.12 (1H, m), 2.29 (3H, s), 2.31 (3H, s), 2.33-2.42 (6H, m), 2.84-2.90 (2H, m),  
35 3.33 (3H, s), 3.46-3.52 (2H, m), 3.60-3.67 (2H, m)

- 120 -

m), 3.80 (3H, s), 3.87-4.00 (2H, m), 4.78-4.87 (1H, m), 6.58 (1H, d, J=7Hz), 6.65 (1H, s), 6.82-6.92 (2H, m), 7.03-7.10 (3H, m), 7.45 (1H, t, J=8Hz), 8.21 (1H, dd, J=1, 8Hz), 8.40 (1H, d, J=7Hz)

ESI-MASS (m/z) : 674 (M+H)

- 4) (R)-4-[2-[(4-Aminobut-2-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.43 and 1.45 (total 3H, s), 1.46-1.91 (12H, m), 2.02-2.12 (1H, m), 2.29 (9H, s), 2.30-2.41 (4H, m), 2.52-2.63 (1H, m), 2.87 (2H, t, J=8Hz), 2.97-3.07 (1H, m), 3.35 (3H, s), 3.80 (3H, s), 3.87-3.98 (4H, m), 4.59-4.68 (1H, m), 4.79-4.88 (1H, m), 6.59 (1H, d, J=8Hz), 6.64 (1H, s), 6.83-6.93 (2H, m), 7.05-7.10 (3H, m), 7.45 (1H, t, J=8Hz), 8.23 (1H, d, J=9Hz), 8.42 (1H, d, J=8Hz)

ESI-MASS (m/z) : 702 (M+H)

- 5) (S)-4-[2-[(4-Aminobut-2-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.43 and 1.45 (total 3H, s), 1.46-1.92 (12H, m), 2.02-2.13 (1H, m), 2.28 (9H, s), 2.30-2.40 (4H, m), 2.52-2.63 (1H, m), 2.86 (2H, t, J=8Hz), 2.97-3.07 (1H, m), 3.35 (3H, s), 3.81 (3H, s), 3.87-3.98 (4H, m), 4.60-4.68 (1H, m), 4.79-4.89 (1H, m), 6.59 (1H, d, J=8Hz), 6.54 (1H, s), 6.83-6.93 (2H, m), 7.05-7.10 (3H, m), 7.46 (1H, t, J=8Hz), 8.23 (1H, d, J=9Hz), 8.43 (1H, d, J=8Hz)

- 121 -

ESI-MASS (m/z) : 702 (M+H)

- 6) 4-[2-[4-Aminobut-1-yl]oxybenzoyl]amino-N-methyl-N-[2-  
[5-(4-methylpiperazin-1-yl)carbonylpent-1-  
yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.47-1.74 (8H, m), 1.77-1.88 (2H, m), 1.95-2.06 (2H, m), 2.27 (3H, s), 2.31-2.40 (4H, m), 2.78 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.45-3.50 (2H, m), 3.58-3.65 (2H, m), 3.84-3.98 (2H, m), 4.20 (2H, t, J=7.5Hz), 6.72-6.80 (2H, m), 6.93-7.00 (2H, m), 7.04-7.14 (2H, m), 7.30 (2H, d, J=8Hz), 7.40-7.48 (3H, m), 8.19 (1H, d, J=7Hz)

- 7) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.24 (3H, t, J=7.5Hz), 1.44-1.56 (2H, m), 1.63-1.87 (6H, m), 2.06-2.16 (2H, m), 2.28 (3H, s), 2.33 (2H, t, J=7.5Hz), 2.97 (2H, t, J=7.5Hz), 3.30 (3H, s), 3.82-3.96 (2H, m), 4.11 (2H, q, J=7.5Hz), 4.30 (2H, t, J=7.5Hz), 6.54-6.60 (2H, m), 6.83 (1H, d, J=7Hz), 7.00 (1H, d, J=7Hz), 7.09 (1H, t, J=7Hz), 7.30 (2H, d, J=8Hz), 7.41-7.48 (3H, m), 8.20 (1H, d, J=7Hz)

- 8) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylprop-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 2.03-2.17 (2H, m), 2.29 (3H, s), 2.33-2.42 (2H, m), 2.53 (2H, t, J=7.5Hz), 2.96 (2H, t, J=7.5Hz), 3.38 (3H, s), 3.46-3.53 (2H, m), 3.59-3.68 (2H, m), 3.92-4.08 (2H, m), 4.28 (2H, t, J=7.5Hz), 6.77-6.83 (2H, m), 6.98-7.18

- 122 -

(4H, m), 7.31 (2H, d, J=8Hz), 7.43-7.50 (3H, m),  
8.20 (1H, d, J=7Hz)

- 5 9) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]-phenylmethoxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.02-2.14 (2H, m), 2.37 (3H, s),  
2.30 (3H, s), 2.32-2.51 (4H, m), 2.94 (2H, t,  
J=7.5Hz), 3.35 (3H, s), 3.41-3.57 (2H, m), 3.67-  
10 3.86 (2H, m), 4.30 (2H, t, J=7.5Hz), 4.96 (1H,  
d, J=14Hz), 5.08 (1H, d, J=14Hz), 6.63-6.71 (2H,  
m), 6.95-7.02 (2H, m), 7.11 (1H, t, J=7Hz), 7.31  
(2H, d, J=8Hz), 7.36-7.50 (7H, m), 8.22 (1H, d,  
J=7Hz)  
15
- 10) 4-[2-(4-Amino-1-butyn-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.42-1.90 (10H, m), 2.28 (3H, s),  
2.32-2.41 (6H, m), 3.37 (3H, s), 3.46-3.51 (2H,  
20 m), 3.59-3.67 (2H, m), 3.82-4.02 (2H, m), 6.73-  
6.82 (2H, m), 7.00 (1H, d, J=7Hz), 7.08-7.20  
(2H, m), 7.35-7.64 (5H, m), 7.81-7.88 (2H, m)
- 25 11) 4-[2-(4-Aminobut-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide  
MASS (m/z) : 614 (M+1)
- 30 12) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.01-2.11 (2H, m), 2.28 (3H, s),  
2.31 (3H, s), 2.33-2.51 (4H, m), 2.90 (2H, t,  
J=7.5Hz), 3.39 (3H, s), 3.40-3.52 (2H, m), 3.61-  
35



- 123 -

- 3.86 (2H, m), 3.67 (3H, s), 4.79 (2H, t, J=7.5Hz), 4.90 (1H, d, J=14Hz), 5.08 (1H, d, J=14Hz), 6.61-6.70 (2H, m), 7.86 (1H, d, J=7Hz), 6.94-7.10 (4H, m), 7.31-7.46 (5H, m), 8.20 (1H, d, J=7Hz), 8.37 (1H, d, J=7Hz)
- 13) 3-Methoxy-4-[2-(3-aminoprop-1-yl)oxy]phenylmethyl]-amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide
- 10 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.54 (2H, m), 1.62-1.71 (2H, m), 1.76-1.85 (2H, m), 1.87-2.00 (2H, m), 2.27 (3H, s), 2.30 (3H, s), 2.31-2.40 (4H, m), 2.90 (2H, t, J=7.5Hz), 3.28 (3H, s), 3.45-3.50 (2H, m), 3.57-3.64 (2H, m), 3.61 (3H, s), 3.80-3.97 (2H, m), 4.07 (2H, t, J=7.5Hz), 4.27 (2H, s), 4.70 (1H, br), 6.37 (1H, d, J=7Hz), 6.59 (1H, d, J=7Hz), 6.62 (1H, s), 6.78 (1H, s), 6.82-6.90 (4H, m), 7.16-7.71 (2H, m)
- 15
- 20 14) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[4-(4-methylpiperazin-1-yl)carbonyl]-phenyleth-1-yl]phenylbenzamide
- 25 NMR (CDCl<sub>3</sub>, δ) : 2.00-2.11 (2H, m), 2.29 (3H, s), 2.32-2.50 (4H, m), 2.61-2.93 (6H, m), 3.32 (3H, s), 3.35-3.89 (2H, m), 3.59-3.81 (2H, m), 3.71 (3H, s), 4.22-4.32 (2H, m), 6.83 (1H, d, J=7Hz), 6.94-7.33 (11H, m), 7.43 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.39 (1H, d, J=7Hz)
- 30 15) 4-[2-(3-Aminoprop-1-yl)thiobenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide
- MASS (m/z) : 676 (M+1)
- 35 16) 4-[2-(3-Aminoprop-1-yl)sulfonylbenzoyl]amino-3-

- 124 -

methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

MASS (m/z) : 724 (M+1)

- 5     17) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-(4-dimethylaminopiperidin-1-yl)carbonyl]-phenylmethoxy-4-methyl]phenyl-N-methylbenzamide  
MASS (m/z) : 708 (M+1)
- 10    18) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylmethoxyprop-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.00-2.14 (4H, m), 2.23 (3H, s),  
15                2.29-2.38 (4H, m), 2.88 (2H, t, J=7.5Hz), 3.35 (3H, s),  
                 3.37-3.45 (2H, m), 3.54-3.61 (2H, m), 3.66-3.76 (2H, m),  
                 3.77 (3H, s), 3.94-4.17 (4H, m), 4.30 (2H, t, J=7.5Hz),  
                 6.75-7.18 (8H, m), 7.45 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz),  
20                8.42 (1H, d, J=7Hz)
- 25    19) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[(E)-5-(4-dimethylaminopiperidin-1-yl)carbonyl-4-penten-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide  
MASS (m/z) : 686 (M+1)
- 30    20) 3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-N-[2-(4-aminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.41 (9H, s), 1.50-1.67 (2H, m),  
                 1.77-1.89 (2H, m), 2.06-2.21 (2H, m), 2.27 (3H, s),  
                 2.80 (2H, t, J=7.5Hz), 3.23-3.36 (2H, m), 3.36 (3H, s),  
                 3.80 (3H, s), 3.84-4.03 (2H, m), 4.26 (2H, t, J=7.5Hz),  
                 6.57-6.68 (2H, m), 6.83-7.15 (5H, m), 7.45 (1H, t, J=7Hz),  
35                8.21 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 125 -

- 21) 4-[2-(3-Amino-1-methylprop-1-yl)oxybenzoyl]amino-3-methoxy-N-(2-benzyloxy-4-methyl)phenyl-N-methylbenzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.41 (3H, d, J=7.5Hz), 1.70-1.83 (1H, m), 1.96-2.10 (1H, m), 2.26 (3H, s), 2.80-2.89 (2H, m), 3.37 (3H, s), 3.62 (3H, s), 4.82 (1H, m), 4.89 (1H, d, J=14Hz), 5.07 (1H, d, J=14Hz), 6.63-6.72 (2H, m), 7.86 (1H, d, J=7Hz), 6.98 (1H, d, J=7Hz), 7.02-7.11 (3H, m), 7.28-7.49 (6H, m), 8.22 (1H, d, J=7Hz), 8.37 (1H, d, J=7Hz)

- 22) 4-[2-(4-Aminobut-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.46-2.03 (10H, m), 2.24 (3H, s), 2.28 (3H, s), 2.31-2.40 (6H, m), 2.73 (2H, t, J=7.5Hz), 3.31 (3H, s), 3.44-3.50 (2H, m), 3.59-3.65 (2H, m), 3.77 (3H, s), 3.83-4.00 (2H, m), 4.20 (2H, t, J=7.5Hz), 6.58 (1H, d, J=7Hz), 7.61 (1H, s), 6.85 (1H, d, J=7Hz), 6.90 (1H, d, J=7Hz), 6.87-7.10 (3H, m), 7.45 (1H, t, J=7Hz), 8.21 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 25 23) 4-[2-(3-Aminoprop-1-yl)oxy-3-methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.49-1.91 (6H, m), 1.96-2.07 (2H, m), 2.27 (3H, s), 2.30 (3H, s), 2.35 (3H, s), 2.32-2.40 (6H, m), 2.95 (2H, t, J=5Hz), 3.32 (3H, s), 3.46-3.53 (2H, m), 3.60-3.67 (2H, m), 3.81 (3H, s), 3.85-4.02 (4H, m), 6.56-6.66 (2H, m), 6.82-7.18 (4H, m), 7.33 (1H, d, J=8Hz), 7.80 (1H, d, J=7Hz), 8.36 (1H, d, J=7Hz)

35

- 126 -

- 24) 4-[2-(3-Aminoprop-1-yl)oxy-4-methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.46-1.90 (6H, m), 2.13-2.25 (2H, m), 2.26 (3H, s), 2.28 (3H, s), 2.30-2.58 (6H, m), 2.37 (3H, s), 2.99 (2H, t, J=5Hz), 3.30 (3H, s), 3.49 (3H, s), 3.49 (2H, t, J=5Hz), 3.61 (2H, t, J=5Hz), 3.79 (3H, s), 3.83-3.92 (2H, m), 4.28 (2H, t, J=5Hz), 6.56-6.65 (2H, m), 6.80-6.93 (4H, m), 7.00 (1H, s), 8.02 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)
- 25) 4-[2-(3-Aminoprop-1-yl)oxy-5-methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.49-1.90 (6H, m), 1.98-2.20 (4H, m), 2.28 (3H, s), 2.29 (3H, s), 2.31 (3H, s), 2.31-2.42 (4H, m), 2.95 (2H, t, J=5Hz), 3.31 (3H, s), 3.50 (2H, t, J=4Hz), 3.62 (2H, t, J=4Hz), 3.79 (3H, s), 3.80-4.00 (2H, m), 4.25 (2H, t, J=5Hz), 6.57-6.68 (2H, m), 6.82-7.04 (4H, m), 7.24 (1H, d, J=8Hz), 7.95 (1H, s), 8.39 (1H, d, J=8Hz)
- 26) 4-[2-(3-Aminoprop-1-yl)oxy-4-chlorobenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.48-1.60 (2H, m), 1.62-1.90 (2H, m), 2.10 (2H, t, J=6Hz), 2.27 (3H, s), 2.29 (3H, s), 2.30-2.41 (4H, m), 2.93 (2H, t, J=5Hz), 3.31 (3H, s), 3.45-3.53 (2H, m), 3.58-3.66 (2H, m), 3.78 (3H, s), 3.82-4.01 (2H, m), 4.29 (2H, t, J=5Hz), 6.55-6.68 (2H, m), 6.80-6.91 (2H, m), 6.99-7.10 (4H, m), 8.13 (1H, d, J=8Hz), 8.36 (1H, d, J=8Hz)

- 127 -

27) 4-[2-(3-Aminoprop-1-yl)oxy-4-methoxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.89 (6H, m), 2.04-2.15 (2H, m), 2.28 (3H, s), 2.30 (3H, s), 2.31-2.42 (6H, m), 2.93 (2H, t, J=5Hz), 3.31 (3H, s), 3.44-3.52 (2H, m), 3.57-3.65 (2H, m), 3.79 (3H, s), 3.83 (3H, s), 3.83-4.00 (2H, m), 4.26 (2H, t, J=5Hz),  
10 7.50-7.68 (4H, m), 6.82-6.95 (2H, m), 7.03 (3H, s), 8.16 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

#### Example 8

A mixture of 4-(2-benzyloxybenzoyl)amino-N-methyl-N-[2-(5-carboxypent-1-yloxy)phenyl]benzamide (1.76 g), N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride (714 mg), N-methylpiperazine (311 mg) and 1-hydroxybenzotriazol (504 mg) in N,N-dimethylformamide (20 ml) was stirred at ambient temperature for 2 hours and the mixture was diluted with ethyl acetate (40 ml). The solution was washed successively with saturated aqueous sodium hydrogen carbonate solution (40 ml), water (40 ml) and brine (40 ml), and dried over magnesium sulfate. The solvent was evaporated to give 4-(2-benzyloxybenzoyl)amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (1.98 g) as a colorless oil.

25 NMR (CDCl<sub>3</sub>, δ) : 1.46-1.58 (2H, m), 1.64-1.88 (4H, m), 2.30 (3H, s), 2.32-2.41 (6H, m), 3.32 (3H, s), 3.49 (2H, m), 3.62 (2H, m), 3.81-4.00 (2H, br), 5.20 (2H, s), 6.73-6.82 (2H, m), 6.94-7.00 (3H, m), 7.08-7.20 (5H, m), 7.40-7.53 (6H, m),  
30 8.28 (1H, d, J=7Hz)

#### Example 9

The following compound was obtained by using 4-[2-(carboxymethoxy)benzoyl]amino-N-methyl-N-[2-[5-(4-

- 128 -

methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide as a starting compound according to a similar manner to that of Example 8.

- 5            4-[2-[(4-Methyloperazin-1-yl)carbonyl-methoxy]-benzoyl]amino-N-methyl-N-[2-[5-(4-methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide  
             MASS : 699 (M+1)

10    Example 10

             A solution of 4-(2-benzyloxybenzoyl)amino-N-methyl-N-[2-[5-(4-methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (1.90 g) in methanol (30 ml) was hydrogenated under atmospheric pressure in the presence of palladium hydroxide (400 mg) for 6 hours and the catalyst was removed by filtration. The filtrate was evaporated to give 4-(2-hydroxybenzoyl)amino-N-methyl-N-[2-[5-(4-methyloperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (1.60 g) as a colorless amorphous.

20            NMR (CDCl<sub>3</sub>, δ) : 1.51 (2H, m), 1.66 (2H, m), 1.79 (2H, m), 2.30 (2H, m), 2.63 (3H, s), 2.82-2.95 (4H, m), 3.33 (3H, s), 3.72 (2H, m), 3.86 (2H, m), 3.99 (2H, m), 6.78-6.93 (3H, m), 7.05 (2H, m), 7.17 (1H, t, J=7Hz), 7.27 (2H, d, J=8.5Hz), 7.40 (1H, t, J=7Hz), 7.53 (2H, d, J=8.5Hz), 7.91 (1H, m), 9.21 (1H, br)

25            Example 11

             The following compounds were obtained according to a similar manner to that of Example 10.

- 30            1) 4-(2-Hydroxybenzoyl)amino-N-methyl-N-[2-[3-(4-methyloperazin-1-yl)carbonylamino-prop-1-yloxy]phenyl]benzamide  
35            NMR (CDCl<sub>3</sub>, δ) : 2.00 (2H, m), 2.71 (3H, s), 2.90-

- 129 -

3.09 (4H, m), 3.33 (3H, s), 3.50-3.80 (6H, m),  
3.97 (2H, m), 6.76-7.03 (5H, m), 7.11-7.22 (2H,  
m), 7.29-7.44 (3H, m), 7.45-7.54 (2H, m), 7.88  
(1H, m)

5

- 2) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yloxy]-4-methylphenyl]benzamide

10 NMR (CDCl<sub>3</sub>, δ) : 1.46-1.62 (2H, m), 1.65-1.90 (4H,  
m), 2.29 (3H, s), 2.30-2.43 (2H, m), 2.82 (3H,  
s), 2.88-3.30 (4H, m), 3.31 (3H, s), 3.48 (3H,  
s), 3.79 (3H, s), 3.77-4.07 (6H, m), 6.58-6.69  
(2H, m), 6.84-7.08 (5H, m), 7.43 (1H, dd, J=9,  
9Hz), 7.52 (1H, d, J=9Hz), 8.20 (1H, d, J=9Hz),  
15 8.82 (1H, br s)

- 3) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.63 (2H, m), 1.64-1.90 (4H,  
m), 2.38 (2H, t, J=7Hz), 2.78 (3H, s), 2.90-3.31  
(4H, m), 3.33 (3H, s), 3.77 (3H, s), 3.80-4.07  
(6H, m), 6.77-7.11 (7H, m), 7.12-7.23 (1H, m),  
7.37-7.58 (2H, m), 8.21 (1H, d, J=9Hz), 8.79  
25 (1H, s)

- 4) 2-Chloro-4-[2-(hydroxy)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-phenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.38-1.98 (6H, m), 2.21-2.46 (2H,  
m), 2.73 (3H, br s), 2.92-3.25 (4H, m), 3.36  
(3H, s), 3.70-4.20 (6H, m), 6.67-6.82 (2H, m),  
6.82-7.08 (4H, m), 7.08-7.20 (2H, m), 7.21-7.50  
(2H, m), 7.70 (1H, br s), 7.92 (1H, br d,  
35 J=8Hz), 9.48 (1H, s)

- 130 -

- 5) 4-(3-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.34-1.58 (4H, m), 1.64-1.97 (6H, m), 2.28 (3H, s), 2.32 (6H, s), 2.33-2.38 (3H, m), 2.51-2.61 (1H, m), 2.97-3.06 (1H, m), 3.34 (3H, s), 3.78-3.81 (3H, br s), 3.85-3.97 (3H, m), 4.60-4.69 (1H, m), 6.58-6.65 (2H, m), 6.84-7.06 (4H, m), 7.38-7.60 (3H, m), 8.17-8.23 (1H, m)

10 ESI-MASS (m/z) : 631 (M+H)

- 6) 4-(2-Hydroxybenzoyl)amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

15 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.23 (3H, t, J=7.5Hz), 1.41-1.53 (2H, m), 1.62-1.84 (4H, m), 2.27 (3H, s), 2.32 (2H, t, J=7.5Hz), 3.31 (3H, s), 3.78-3.97 (2H, m), 4.13 (2H, q, J=7.5Hz), 6.56-6.61 (2H, m), 6.84-6.91 (2H, m), 7.02 (1H, d, J=7Hz), 7.28-7.45 (4H, m), 7.62 (1H, d, J=7Hz), 8.47 (1H, s)

- 7) 4-(2-Hydroxybenzoyl)amino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylprop-1-yl]oxy]-phenylbenzamide

25 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 2.00 (2H, m), 2.71 (3H, s), 2.90-3.09 (4H, m), 3.33 (3H, s), 3.50-3.80 (6H, m), 3.97 (2H, m), 6.76-7.03 (5H, m), 7.11-7.22 (2H, m), 7.29-7.44 (3H, m), 7.45-7.54 (2H, m), 7.88 (1H, m)

- 8) 4-(2-Hydroxy)benzoylamino-3-methoxy-N-methyl-N-[2-[4-(4-methylpiperazin-1-yl)carbonyl]phenyleth-1-yl]phenylbenzamide

35 MASS (m/z) : 607 (M+1)



- 131 -

- 9) 4-(2-Hydroxy-3-methylbenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.50-1.90 (6H, m), 2.27 (6H, s),  
5 2.28 (3H, s), 2.33-2.40 (4H, m), 2.70-2.78 (2H, m), 3.30 (3H, s), 3.80 (3H, s), 3.85-4.10 (6H, m), 6.59-6.65 (2H, m), 6.77-6.97 (6H, m), 8.19 (1H, d, J=8Hz), 8.70 (1H, br s)
- 10 10) 4-(2-Hydroxy-4-methylbenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.49-1.91 (6H, m), 2.24 (3H, s),  
15 2.29 (3H, s), 2.32 (3H, s), 2.30-2.42 (6H, m), 3.32 (3H, s), 3.49 (2H, t, J=5Hz), 3.63 (2H, t, J=5Hz), 3.80 (3H, s), 3.88-4.01 (2H, m), 6.68-6.65 (2H, m), 6.80 (1H, s), 6.84 (1H, d, J=8Hz), 6.93 (1H, d, J=7Hz), 7.03 (1H, s), 7.37 (1H, d, J=7Hz), 8.19 (1H, d, J=8Hz), 8.71 (1H, br)
- 20 11) 4-(2-Hydroxy-4-methylbenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.50-1.91 (10H, m), 2.28 (3H, s),  
25 2.34 (3H, s), 2.35 (3H, s), 2.30-2.41 (6H, m), 2.80 (2H, br), 3.31 (3H, s), 3.80 (3H, s), 3.81-4.09 (4H, m), 6.60-6.68 (2H, m), 6.84-7.02 (4H, m), 7.20-7.30 (2H, m), 8.20 (1H, br), 8.37 (1H, br)
- 30 12) 4-(2-Hydroxy-4-chlorobenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.46-1.89 (6H, m), 2.23-2.45 (6H, m),  
35 2.27 (3H, s), 2.32 (3H, s), 3.30 (3H, s),

- 132 -

3.44-3.68 (4H, m), 3.80 (3H, s), 3.80-3.99 (2H, m), 6.53-6.65 (2H, m), 6.72-7.03 (5H, m), 7.41 (1H, d, J=8Hz), 8.12 (1H, d, J=8Hz), 8.74 (1H, br)

5

- 13) 4-(2-Hydroxy-4-methoxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-ylcarbonyl)pent-1-yl]oxy-4-methylphenyl]benzamide

10

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.60 (2H, m), 1.63-1.84 (4H, m), 2.28 (3H, s), 2.37 (2H, t, J=5Hz), 2.25-2.40 (6H, m), 2.79 (3H, s), 3.30 (3H, s), 3.79 (3H, s), 3.82 (3H, s), 3.90-4.01 (2H, m), 6.44-6.50 (2H, m), 6.60-6.66 (2H, m), 6.88-6.97 (3H, m), 7.41 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz), 8.40 (1H, br)

15

- 14) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonyl]pent-1-yloxy]phenyl]benzamide

20

NMR (CDCl<sub>3</sub>, δ) : 1.22-1.45 (2H, m), 1.45-1.58 (2H, m), 1.62-1.78 (2H, m), 1.80-1.96 (4H, m), 2.30 (6H, s), 2.30-2.40 (3H, m), 2.50-2.62 (1H, m), 2.97-2.37 (1H, m), 3.37 (3H, s), 3.78 (3H, s), 3.82-4.02 (4H, m), 4.57-4.68 (1H, m), 6.77-7.02 (8H, m), 7.10-7.20 (1H, m), 7.37-7.45 (1H, m), 7.46-7.62 (1H, m), 8.20 (1H, br)

25

- 15) 4-(2-Hydroxybenzoyl)amino-3-chloro-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonyl]pent-1-yloxy]-4-methylphenyl]benzamide

30

NMR (CDCl<sub>3</sub>, δ) : 1.30-2.08 (10H, m), 2.20-2.60 (13H, m), 2.89-3.05 (1H, m), 3.30 (3H, s), 3.82-4.02 (4H, m), 4.62-4.79 (1H, m), 6.62 (2H, s), 6.73-7.02 (4H, m), 7.28-7.57 (3H, m), 7.99 (1H, d, J=7Hz), 8.42 (1H, d, J=7Hz)

35

- 133 -

- 16) 3-Ethoxy-4-(2-hydroxybenzoyl)amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.40 (3H, t, J=6Hz), 1.47-1.57 (2H, m), 1.65-1.72 (2H, m), 1.78-1.88 (2H, m), 2.27 (3H, s), 2.30 (3H, s), 2.31-2.42 (7H, m), 3.30 (3H, s), 3.48-3.50 (2H, m), 3.52-3.65 (2H, m), 3.82-4.02 (4H, m), 6.58-6.61 (2H, m), 6.82-6.94 (3H, m), 6.98-7.02 (2H, m), 7.40-7.47 (2H, m), 10 8.20 (1H, d, J=7Hz), 8.83 (1H, s)

- 17) 3-Hydroxy-4-(2-hydroxybenzoyl)amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.62 (2H, br), 1.75 (2H, br), 1.85 (2H, br), 2.27 (3H, s), 2.30 (3H, s), 2.42 (7H, br), 3.30 (3H, s), 3.53 (2H, br), 3.68 (3H, br), 3.90 (1H, br), 6.52 (1H, s), 6.63-6.73 (2H, m), 6.87 (1H, t, J=7Hz), 6.97 (1H, d, J=7Hz), 7.08 (1H, d, J=7Hz), 7.15 (1H, s), 7.38 (1H, t, J=7Hz), 7.58 (1H, d, J=7Hz), 7.98 (1H, br), 9.02 (1H, br) 20

- 18) 2-(2-Hydroxybenzoyl)amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-5-pyridinecarboxamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.32-2.15 (10H, m), 2.29-2.42 (12H, m), 2.47-2.62 (1H, m), 2.95-3.09 (1H, m), 3.32 (3H, s), 3.75-4.10 (4H, m), 4.58-4.77 (2H, m), 30 6.33-8.47 (15H, m)

- 19) 4-(4-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

35 NMR (CDCl<sub>3</sub>, δ) : 1.38-1.55 (4H, m), 1.62-1.72 (2H,

- 134 -

m), 1.72-1.83 (2H, m), 1.83-1.97 (2H, m), 2.27 (3H, s), 2.32-2.37 (8H, m), 2.43-2.60 (2H, m), 2.93-3.05 (1H, m), 3.31 (3H, s), 3.70 (3H, s), 3.78-3.95 (4H, m), 4.60-4.70 (1H, m), 6.57-6.60 (2H, m), 6.80-6.97 (5H, m), 7.67 (2H, d, J=7Hz), 8.22 (1H, d, J=7Hz), 8.40 (1H, s)

20) 4-(4-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.47-1.58 (2H, m), 1.67-1.75 (2H, m), 1.75-1.87 (2H, m), 2.27 (3H, s), 2.32 (3H, s), 2.38-2.48 (7H, m), 3.35 (3H, s), 3.48-3.53 (2H, m), 3.60-3.70 (2H, m), 3.70 (3H, s), 3.80-3.90 (1H, m), 3.90-4.00 (1H, m), 3.58-3.60 (2H, m), 6.82-6.97 (5H, m), 7.68 (2H, d, J=7Hz), 8.24 (1H, d, J=7Hz), 8.40 (1H, s)

#### Example 12

A solution of 4-(2-hydroxybenzoyl)amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (400 mg) in N,N-dimethylformamide (15 ml) was added potassium carbonate (99 mg) and N-(3-bromopropyl)phthalimide (192 mg) and the mixture was stirred at 60°C for 4 hours. The mixture was poured into water (30 ml) and the aqueous solution was extracted with ethyl acetate (20 ml x 2). The organic phase was washed with water (20 ml) and brine (20 ml), and dried over magnesium sulfate. The solvent was evaporated to give 4-[2-[3-(phthalimido)prop-1-yl]oxy]benzoylamino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (484 mg) as a colorless amorphous.

NMR (CDCl<sub>3</sub>, δ) : 1.56 (2H, m), 1.63-1.76 (4H, m), 1.86 (2H, m), 2.30 (3H, s), 2.32-2.41 (6H, m), 3.35 (3H, s), 3.50 (2H, m), 3.63 (2H, m), 3.83-

- 135 -

3.97 (4H, m), 4.20 (2H, t, J=7.5Hz), 6.73-6.81  
(2H, m), 6.92 (1H, d, J=7Hz), 7.00-7.14 (3H, m),  
7.32 (2H, d, J=8.5Hz), 7.42 (1H, m), 7.50 (2H,  
d, J=8.5Hz), 7.65-7.74 (4H, m), 8.08 (1H, d,  
J=7Hz), 9.69 (1H, s)

### Example 13

The following compounds were obtained according to a  
similar manner to that of Example 12.

- 1) 4-[2-(Ethoxycarbonylmethoxy)benzoyl]amino-N-methyl-N-  
[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-  
yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.31 (3H, t, J=7.5Hz), 1.62 (2H,  
m), 1.71 (2H, m), 1.83 (2H, m), 2.29 (3H, s),  
2.33-2.41 (6H, m), 3.35 (3H, s), 3.49 (2H, m),  
3.62 (2H, m), 3.93 (2H, m), 4.33 (2H, q,  
J=7.5Hz), 4.76 (2H, s), 6.72-6.82 (2H, m), 6.87  
(1H, d, J=7Hz), 7.00 (1H, d, J=7Hz), 7.07-7.18  
(2H, m), 7.33 (2H, d, J=8.5Hz), 7.46 (1H, t,  
J=7Hz), 7.71 (2H, d, J=8.5Hz), 8.26 (1H, d,  
J=7Hz)

- 2) 4-[2-(3-Piperidinoprop-1-yloxy)benzoyl]amino-N-  
methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-  
1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.45 (2H, m), 1.50-1.60 (4H, m),  
1.71 (2H, m), 1.85 (2H, m), 2.14 (2H, m), 2.28  
(3H, s), 2.30-2.41 (10H, m), 2.49 (2H, t,  
J=7.5Hz), 3.34 (3H, s), 3.49 (2H, m), 3.63 (2H,  
m), 3.94 (2H, m), 4.23 (2H, t, J=7.5Hz), 6.73-  
6.82 (2H, m), 6.96-7.02 (2H, m), 7.04-7.15 (2H,  
m), 7.32 (2H, d, J=8.5Hz), 7.43-7.50 (3H, m),  
8.22 (1H, d, J=7Hz)

- 136 -

- 3) 4-[2-[2-(Dimethylamino)eth-1-yloxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]phenyl]benzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.56 (2H, m), 1.70 (2H, m), 1.85 (2H, m), 2.23 (6H, s), 2.30 (3H, s), 2.33-2.41 (6H, m), 2.78 (2H, t, J=7.5Hz), 3.35 (3H, s), 3.50 (2H, m), 3.64 (2H, m), 3.93 (2H, m), 4.22 (2H, t, J=7.5Hz), 6.74-6.81 (2H, m), 6.95-7.01 (2H, m), 7.06-7.15 (2H, m), 7.30 (2H, d, J=8.5Hz), 7.43 (1H, m), 7.56 (2H, d, J=8.5Hz), 8.21 (1H, d, J=7Hz)

- 15 4) 4-[2-[3-(Phthalimido)prop-1-yl]oxy]benzoylamino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)-carbonylaminoprop-1-yloxy]phenyl]benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 2.01 (2H, m), 2.25 (3H, s), 2.25-2.38 (6H, m), 3.33-3.45 (6H, m), 3.35 (3H, s), 3.87-4.00 (4H, m), 4.21 (2H, t, J=7.5Hz), 6.78-7.00 (3H, m), 7.06-7.20 (3H, m), 7.33-7.56 (4H, m), 7.65-7.75 (4H, m), 7.86 (1H, m), 8.10 (1H, d, J=7Hz), 9.73 (1H, br)

- 25 5) 4-[2-[3-(Phthalimido)prop-1-yl]benzoylamino]-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]-4-methylphenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.46-1.62 (2H, m), 1.63-1.93 (4H, m), 2.10-2.46 (14H, m), 3.33 (3H, s), 3.40-3.53 (2H, m), 3.57-3.68 (2H, m), 3.78 (3H, s), 3.79-4.04 (4H, m), 4.26 (2H, t, J=7Hz), 6.54-6.68 (2H, m), 6.74-7.11 (5H, m), 7.37-7.48 (1H, m), 7.52-7.63 (3H, m), 7.66-7.77 (1H, m), 7.80-7.90 (1H, m), 8.06-8.23 (2H, m)

- 35 6) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-

- 137 -

yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.47-1.63 (2H, m), 1.63-1.93 (6H, m), 2.29 (3H, s), 2.29-2.44 (6H, m), 3.36 (3H, s), 3.44-3.53 (2H, m), 3.58-3.68 (2H, m), 3.76 (3H, s), 3.81-4.05 (4H, m), 4.27 (2H, t, J=7Hz), 6.74-6.91 (3H, m), 6.92-7.20 (5H, m), 7.38-7.48 (1H, m), 7.58 (3H, s), 7.68-7.77 (1H, m), 7.82-7.90 (1H, m), 8.09-8.16 (1H, m), 8.20 (1H, d, J=9Hz)

7) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.32-2.00 (12H, m), 2.16-2.48 (12H, m), 2.57 (1H, m), 3.02 (1H, m), 3.33 (3H, s), 3.78 (3H, s), 3.80-4.05 (5H, m), 4.27 (2H, t, J=7Hz), 4.64 (1H, m), 6.56-6.70 (2H, m), 6.78-7.12 (5H, m), 7.43 (1H, m), 7.59 (2H, s), 7.66-7.91 (2H, m), 8.05-8.24 (2H, m)

8) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.26 (3H, t, J=7Hz), 1.34-1.92 (17H, m), 2.23-2.40 (5H, m), 3.20-3.40 (5H, m), 3.78 (3H, s), 3.82-4.01 (2H, m), 4.12 (2H, q, J=7Hz), 4.25 (2H, t, J=7Hz), 4.78 (1H, m), 6.52-6.69 (2H, m), 6.79-7.15 (5H, m), 7.40-7.52 (2H, m), 8.21 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

9) 2-Chloro-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.51-1.67 (2H, m), 1.68-1.82 (2H, m), 1.82-2.01 (2H, m), 2.22-2.48 (11H, m), 3.38

- 138 -

5 (3H, s), 3.47-3.56 (2H, m), 3.58-3.69 (2H, m), 3.90 (2H, t, J=7Hz), 3.94-4.11 (2H, m), 4.21 (2H, t, J=7Hz), 6.69-6.82 (2H, m), 6.93 (1H, d, J=8Hz), 7.02-7.20 (4H, m), 7.30 (1H, m), 7.43 (1H, m), 7.68 (4H, s), 8.07 (1H, m), 9.62 (1H, s)

10 10) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]-benzoyl]amino-3-methoxy-N-methyl-N-(2-methylphenyl)-benzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.41 (9H, s), 2.02-2.18 (2H, m), 2.21 (3H, s), 3.21-3.34 (2H, m), 3.39 (3H, s), 3.75 (3H, s), 4.24 (2H, t, J=7Hz), 4.74 (1H, m), 6.83-7.22 (9H, m), 7.44 (1H, m), 8.20 (1H, m), 8.42 (1H, d, J=8Hz)

20 11) 4-[3-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]-benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.32-1.45 (2H, m), 1.43 (9H, s), 1.49-1.58 (2H, m), 1.64-1.90 (6H, m), 1.97-2.03 (2H, m), 2.29 (3H, s), 2.30 (6H, s), 2.33-2.39 (3H, m), 2.51-2.61 (1H, m), 2.97-3.07 (1H, m), 3.28-3.38 (2H, m), 3.33 (3H, s), 3.79 (3H, s), 3.86-3.97 (3H, m), 4.08 (2H, t, J=7Hz), 4.59-4.67 (1H, m), 4.70-4.78 (1H, m), 6.57-6.64 (2H, m), 6.84 (1H, d, J=8Hz), 6.93 (1H, d, J=8Hz), 7.02 (1H, s), 7.03-7.07 (1H, m), 7.33-7.40 (3H, m), 8.27 (1H, d, J=8Hz), 8.49 (1H, s)

30 ESI-MASS (m/z) : 788 (M+1)

35 12) 4-[2-[4-(Phthalimido)but-1-yl]oxybenzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide



- 139 -

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.60 (2H, m), 1.65-1.77 (4H, m),  
 1.80-2.06 (6H, m), 2.29 (3H, s), 2.33-2.41 (6H,  
 m), 3.38 (3H, s), 3.45-3.51 (2H, m), 3.60-3.67  
 (2H, m), 3.78 (2H, t, J=7.5Hz), 3.88-4.00 (2H,  
 m), 4.23 (2H, d, J=7.5Hz), 6.73-6.42 (2H, m),  
 6.99 (2H, d, J=8Hz), 7.08-7.17 (2H, m), 7.36  
 (2H, d, J=8Hz), 7.44-7.50 (3H, m), 7.68-7.77  
 (2H, m), 7.81-7.91 (2H, m), 8.22 (1H, d, J=7Hz)

- 10 13) 4-[2-[3-(Phthalimido)prop-1-yl]oxybenzoyl]amino-N-[2-  
 (5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl]-N-  
 methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.24 (3H, t, J=7.5Hz), 1.45-1.57  
 (2H, m), 1.64-1.88 (4H, m), 2.25 (3H, s), 2.28-  
 2.37 (4H, m), 3.31 (3H, s), 3.84-3.95 (4H, m),  
 4.10 (2H, q, J=7.5Hz), 4.20 (2H, t, J=7.5Hz),  
 6.52-6.62 (2H, m), 6.88 (1H, d, J=7Hz), 6.92  
 (1H, d, J=7Hz), 7.07 (1H, t, J=7Hz), 7.31 (2H,  
 d, J=8Hz), 7.39-7.50 (3H, m), 7.62-7.64 (4H, m),  
 8.10 (1H, d, J=7Hz), 9.68 (1H, s)

- 14) 4-[2-[3-(Phthalimido)prop-1-yl]oxybenzoyl]amino-N-  
 methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylprop-  
 1-yl]oxy]phenylbenzamide

MASS (m/z) : 718 (M+1)

- 15) 4-[2-[3-(Phthalimido)prop-1-yl]oxybenzoyl]amino-N-  
 methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-  
 yl)carbonyl]phenylmethoxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 2.25 (3H, s), 2.25-2.31 (2H, m),  
 2.31 (3H, s), 2.36-2.51 (4H, m), 3.38 (3H, s),  
 3.63-3.85 (4H, m), 3.91 (2H, t, J=7.5Hz), 4.20  
 (2H, t, J=7.5Hz), 4.98 (1H, d, J=14Hz), 5.08  
 (1H, d, J=14Hz), 6.63-6.70 (2H, m), 6.90-7.00  
 (2H, m), 7.09 (1H, t, J=7Hz), 7.32 (2H, d,

- 140 -

J=8Hz), 7.40-7.77 (7H, m), 8.10 (1H, d, J=7Hz),  
9.70 (1H, s)

- 5 16) 4-[2-(3-Hydroxyprop-1-yl)oxybenzoyl]amino-N-methyl-N-  
[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-  
yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.43-1.56 (2H, m), 1.60-1.86 (4H,  
m), 2.11-2.23 (2H, m), 2.14 (3H, s), 2.37-2.90  
(6H, m), 3.33 (3H, s), 3.40-3.47 (2H, m), 3.51-  
10 3.59 (2H, m), 3.86 (2H, t, J=7.5Hz), 3.86-4.00  
(2H, m), 4.32 (2H, t, J=7.5Hz), 6.78-6.85 (2H,  
m), 6.99-7.19 (4H, m), 7.31 (2H, d, J=8Hz),  
7.41-7.53 (3H, m), 8.21 (1H, d, J=8Hz)
- 15 17) 4-[2-(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-  
methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-  
carbonylpent-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.43-1.54 (2H, m), 1.60-1.70 (2H,  
m), 1.74-1.85 (2H, m), 2.10-2.21 (2H, m), 2.26  
20 (6H, sx2), 2.30-2.41 (6H, m), 3.32 (3H, s),  
3.40-3.48 (2H, m), 3.55-3.61 (2H, m), 3.77 (3H,  
s), 3.77-4.00 (4H, m), 4.31 (2H, t, J=7.5Hz),  
6.57-6.63 (2H, m), 6.85-6.92 (2H, m), 7.00-7.11  
(3H, m), 7.44 (1H, t, J=7Hz), 8.20 (1H, d,  
25 J=7Hz), 8.40 (1H, d, J=7Hz)
- 30 18) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-  
amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-  
yl)carbonyl]phenylmethoxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.19-2.32 (2H, m), 2.25 (3H, s),  
2.33 (3H, s), 2.36-2.52 (4H, m), 3.33-3.50 (2H,  
m), 3.39 (3H, s), 3.67 (3H, s), 3.71-3.91 (4H,  
m), 4.28 (2H, t, J=7.5Hz), 4.95 (1H, d, J=14Hz),  
5.09 (1H, d, J=14Hz), 6.62-6.72 (2H, m), 6.81  
35 (1H, d, J=7Hz), 6.93-7.08 (4H, m), 7.34-7.47

- 141 -

(4H, m), 7.59 (1H, m), 7.68-7.75 (2H, m), 7.82-7.88 (2H, m), 8.10-8.19 (2H, m)

- 5 19) 4-[2-(3-Ethoxycarbonylprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide
- 10 NMR (CDCl<sub>3</sub>, δ) : 1.22 (3H, t, J=7.5Hz), 1.45-1.57 (2H, m), 1.63 (3H, s), 1.63-1.73 (2H, m), 1.76-1.88 (2H, m), 2.20-2.32 (2H, m), 2.24 (3H, s), 2.27 (3H, s), 2.32-2.40 (6H, m), 2.50 (2H, t, J=7.5Hz), 3.31 (3H, s), 3.43-3.50 (2H, m), 3.58-3.67 (2H, m), 3.78 (3H, s), 3.83-4.00 (2H, m), 4.12 (2H, q, J=7.5Hz), 4.22 (2H, t, J=7.5Hz), 6.57 (1H, d, J=7Hz), 6.62 (1H, s), 6.80-6.90 (2H, m), 6.97-7.11 (3H, m), 7.45 (1H, t, J=7Hz), 8.21 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)
- 15 20) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxy]-phenylmethylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]-phenylbenzamide
- 20 NMR (CDCl<sub>3</sub>, δ) : 1.44-1.57 (2H, m), 1.62-1.72 (2H, m), 1.72-1.95 (4H, m), 2.18 (2H, t, J=7.5Hz), 2.25 (3H, s), 2.28 (3H, s), 2.28-2.43 (4H, m), 3.28 (3H, s), 3.43-3.50 (2H, m), 3.57-3.65 (2H, m), 3.58 (3H, s), 3.80-3.96 (2H, m), 4.02 (2H, t, J=7.5Hz), 4.24 (2H, s), 4.80 (1H, s), 6.27 (1H, d, J=7Hz), 6.60 (1H, d, J=7Hz), 6.64 (1H, s), 6.80-6.95 (5H, m), 7.12-7.21 (2H, m), 7.64-7.88 (4H, m)
- 25 21) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-N-methyl-N-[2-[4-(4-methylpiperazin-1-yl)carbonyl]phenyleth-1-yl]phenylbenzamide
- 30 NMR (CDCl<sub>3</sub>, δ) : 2.20-2.50 (6H, m), 2.29 (3H, s),

- 142 -

5 2.61-2.94 (6H, m), 3.30 (3H, s), 3.37-3.68 (2H, m), 3.68 (3H, s), 3.68-3.92 (2H, m), 4.20-4.30 (2H, m), 6.80 (1H, d, J=7Hz), 6.90-6.98 (2H, m), 7.05 (1H, t, J=7Hz), 7.10-7.49 (9H, m), 7.53-7.89 (4H, m), 8.12 (1H, d, J=7Hz), 8.20 (1H, d, J=7Hz)

10 22) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-N-[2-[4-(4-dimethylaminopiperidin-1-yl)carbonyl]phenylmethoxy-4-methyl]phenyl-N-methylbenzamide

MASS (m/z) : 824 (M+1)

15 23) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylmethoxyprop-1-yl]oxy]phenylbenzamide

20 NMR (CDCl<sub>3</sub>, δ) : 2.04-2.17 (2H, m), 2.25 (3H, s), 2.28-2.40 (6H, m), 3.33 (3H, s), 3.38-3.46 (2H, m), 3.54-3.62 (2H, m), 3.66-3.76 (2H, m), 3.74 (3H, s), 3.80-3.90 (2H, m), 3.98-4.11 (4H, m), 4.28 (2H, t, J=7.5Hz), 6.78-7.10 (7H, m), 7.14 (1H, t, J=7Hz), 7.43 (1H, t, J=7Hz), 7.55 (2H, s), 7.68-7.75 (1H, m), 7.81-7.90 (1H, m), 8.13 (1H, d, J=7Hz), 8.20 (1H, d, J=7Hz)

25 24) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-N-[2-[(E)-5-(4-dimethylaminopiperidin-1-yl)carbonyl-4-penten-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.27-1.47 (2H, m), 1.83-2.02 (2H, m), 2.10-2.48 (6H, m), 2.23 (3H, s), 2.26 (6H, s), 2.50-4.13 (8H, m), 3.32 (3H, s), 3.78 (3H, s), 4.26 (2H, t, J=7.5Hz), 4.62 (2H, m), 6.32 (1H, d, J=15Hz), 6.57-6.67 (2H, m), 6.80-7.16 (5H, m), 7.44 (1H, t, J=7Hz), 7.53-7.88 (5H, m),

35

- 143 -

7.57 (2H, s), 8.09-8.19 (2H, m)

- 25) 3-Methoxy-4-[2-(pyrid-3-yl)methoxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.44-1.57 (2H, m), 1.63-1.72 (2H, m), 1.75-1.86 (2H, m), 2.27 (3H, s), 2.30 (3H, s), 2.32-2.40 (6H, m), 3.29 (3H, s), 3.31 (3H, s), 3.45-3.51 (2H, m), 3.58-3.65 (2H, m), 3.80-4.00 (2H, m), 5.30 (2H, s), 6.58 (1H, d, J=7Hz), 6.61 (1H, s), 6.83 (1H, d, J=7Hz), 6.88-6.92 (2H, m), 7.05 (1H, d, J=7Hz), 7.14 (1H, t, J=7Hz), 7.29 (1H, m), 7.46 (1H, t, J=7Hz), 7.79 (1H, d, J=7Hz), 8.22 (1H, d, J=7Hz), 8.37 (1H, d, J=7Hz), 8.62 (6H, d, J=7Hz), 8.73 (1H, s)

- 26) 3-Methoxy-4-[2-[4-(phthalimido)but-1-yl]oxybenzoyl]-amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.46-1.59 (2H, m), 1.65-1.74 (2H, m), 1.78-2.03 (6H, m), 2.26 (3H, s), 2.30 (3H, s), 2.32-2.41 (6H, m), 3.31 (3H, s), 3.43-3.50 (2H, m), 3.60-3.65 (2H, m), 3.74 (2H, t, J=7.5Hz), 3.77 (3H, s), 3.82-4.01 (2H, m), 4.22 (2H, t, J=7.5Hz), 6.58 (1H, d, J=7Hz), 6.63 (1H, s), 6.85 (1H, d, J=7Hz), 6.90 (1H, d, J=7Hz), 7.00 (1H, d, J=7Hz), 7.02 (1H, s), 7.08 (1H, t, J=7Hz), 7.45 (1H, t, J=7Hz), 7.70-7.76 (2H, m), 7.80-7.87 (2H, m), 8.21 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 27) 4-[2-(3-Dimethylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.50-1.91 (6H, m), 2.07-2.18 (2H,

- 144 -

- 5  
2.25 (6H, s), 2.28 (3H, s), 2.30 (3H, s),  
2.32-2.44 (2H, m), 2.48 (2H, t, J=5Hz), 3.32  
(3H, s), 3.49 (2H, t, J=5Hz), 3.63 (2H, t,  
J=3Hz), 3.78 (3H, s), 3.81-3.92 (2H, m), 4.25  
(2H, t, J=5Hz), 6.54-6.64 (2H, m), 6.80-6.91  
(2H, m), 6.99-7.11 (4H, m), 7.40-7.48 (1H, m)  
8.18 (1H, d, J=7Hz), 8.38 (1H, d, J=7Hz)
- 10  
28) 4-[2-(Ethoxycarbonylmethoxy)benzoyl]amino-3-methoxy-  
N-methyl-N-[2-[5-(4-methylpiperazin-1-  
yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.23 (3H, t, J=5Hz), 1.50-1.91 (6H,  
m), 2.28 (3H, s), 2.31 (3H, s), 2.35-2.47 (6H,  
m), 3.33 (3H, s), 3.52 (2H, t, J=5Hz), 3.67 (2H,  
t, J=5Hz), 3.76 (3H, s), 3.84-4.02 (2H, m), 4.24  
15 (2H, q, J=5Hz), 4.85 (2H, s), 6.55-6.67 (2H, m),  
6.81-7.19 (6H, m), 7.41-7.49 (1H, m), 8.20 (1H,  
d, J=8Hz), 8.34 (1H, d, J=7Hz)
- 20  
29) 4-[2-[3-(Phthalimido-1-yl)prop-1-yloxy]-3-  
methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-  
methylpiperazin-1-yl)carbonylpent-1-yloxy-4-  
methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.50-1.92 (6H, m), 2.14-2.44 (8H,  
25 m), 2.25 (3H, s), 2.28 (3H, s), 2.36 (3H, s),  
3.33 (3H, s), 3.50 (2H, t, J=5Hz), 3.58 (2H, t,  
J=5Hz), 3.63 (2H, t, J=5Hz), 3.81 (3H, s), 3.81-  
4.03 (8H, m), 6.55-6.68 (2H, m), 6.82-7.38 (6H,  
m), 7.59-7.88 (5H, m), 8.32 (1H, d, J=8Hz)
- 30  
30) 4-[2-[3-(Phthalimido-1-yl)prop-1-yl]oxy-4-  
methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-  
methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-  
methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.50-1.91 (8H, m), 2.27 (3H, s),  
35

- 145 -

2.30 (3H, s), 2.31-2.42 (6H, m), 2.38 (3H, s),  
3.32 (3H, s), 3.50 (2H, t, J=5Hz), 3.63 (2H, t,  
J=5Hz), 3.78 (3H, s), 3.85-4.02 (6H, m), 4.28  
(2H, t, J=5Hz), 6.58-6.67 (2H, m), 6.77 (1H, s),  
5 6.80-6.92 (4H, m), 7.00 (1H, s), 7.58 (4H, s),  
8.01 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz)

31) 4-[2-[3-(Phthalimido-1-yl)propyloxy]-5-  
methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-  
10 methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-  
methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.52-1.91 (10H, m), 2.25 (3H, s),  
2.30 (3H, s), 2.31 (3H, s), 2.31-2.45 (2H, m),  
3.31 (3H, s), 3.50 (2H, t, J=4Hz), 3.59 (2H, t,  
15 J=5Hz), 3.64 (2H, t, J=4Hz), 3.78 (3H, s), 3.85-  
4.02 (4H, m), 4.24 (2H, t, J=5Hz), 6.58 (2H, m),  
6.81-6.92 (3H, m), 7.00 (1H, s), 7.25 (1H, d,  
J=8Hz), 7.59 (3H, s), 7.71-7.79 (1H, m), 7.82-  
7.89 (1H, m), 7.92 (1H, s), 8.20 (1H, d, J=8Hz)

20 32) 4-[2-[3-(Phthalimido-1-yl)propyloxy]-4-  
chlorobenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-  
methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-  
methylphenyl]benzamide  
25 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.92 (6H, m), 2.25 (3H, s),  
2.30 (3H, s), 2.29-2.44 (8H, m), 3.32 (3H, s),  
3.46-3.54 (2H, m), 3.61-3.68 (2H, m), 3.78 (3H,  
s), 3.80-4.01 (4H, m), 4.25 (2H, t, J=5Hz),  
6.56-6.77 (2H, m), 6.79-7.04 (7H, m), 7.44 (2H,  
30 s), 7.70-7.78 (1H, m), 7.81-7.88 (1H, m), 8.06  
(1H, d, J=8Hz)

33) 4-[2-[3-(Phthalimido-1-yl)propyloxy]-4-  
methoxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-  
35 methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-

- 146 -

methylphenyl]benzamide

5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.49-1.90 (6H, m), 2.15-2.24 (2H, m), 2.28 (3H, s), 2.32 (3H, s), 2.30-2.42 (6H, m), 3.33 (3H, s), 3.50 (2H, t, J=4Hz), 3.60 (2H, t, J=5Hz), 3.63 (2H, t, J=4Hz), 3.79 (3H, s), 3.85 (3H, s), 3.82-4.02 (6H, m), 4.24 (2H, t, J=5Hz), 6.57-6.68 (2H, m), 6.82 (1H, d, J=8Hz), 6.89 (1H, d, J=8Hz), 7.00 (1H, s), 7.57 (2H, s), 7.71-7.76 (2H, m), 7.82-7.88 (2H, m), 8.11 (1H, d, J=9Hz), 8.17 (1H, d, J=8Hz)

15 34) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-(2-benzyloxy-4-methylphenyl)-N-methylbenzamide

20 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 2.10 (2H, t, J=5Hz), 2.29 (3H, s), 3.28 (2H, q, J=5Hz), 3.39 (3H, s), 3.62 (3H, s), 4.21 (2H, t, J=5Hz), 4.90 (1H, d, J=13Hz), 5.08 (1H, d, J=13Hz), 6.63-6.71 (3H, m), 6.87 (1H, d, J=7Hz), 6.96-7.11 (6H, m), 7.31-7.48 (6H, m), 8.21 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

25 35) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-[2-[4-(2-oxazolin-2-yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide

30 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 2.09-2.17 (2H, m), 2.28 (3H, s), 3.27 (1H, q, J=5Hz), 3.40 (3H, s), 3.65 (3H, s), 4.05 (2H, t, J=10Hz), 4.23 (2H, t, J=5Hz), 4.40 (2H, t, J=10Hz), 4.88 (1H, d, J=12Hz), 5.08 (1H, d, J=12Hz), 6.62 (1H, s), 6.68 (1H, d), 6.97-7.11 (6H, m), 7.32 (1H, d, J=8Hz), 7.41 (1H, d, J=8Hz), 7.92 (2H, d, J=8Hz), 8.21 (1H, d, J=8Hz), 8.37 (1H, d, J=8Hz)

35 36) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]oxybenzoyl]-



- 147 -

amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.22-1.47 (2H, m), 1.47-1.80 (6H, m), 1.80-1.92 (4H, m), 2.29 (6H, s), 2.31-2.41 (3H, m), 2.50-2.63 (1H, m), 2.95-3.07 (1H, m), 3.36 (3H, s), 3.48 (1H, s), 3.49 (1H, s), 3.75 (3H, s), 3.82-4.03 (4H, m), 4.22-4.30 (2H, m), 4.60-4.70 (1H, m), 6.78-6.90 (3H, m), 6.92-7.20 (4H, m), 7.40-7.50 (1H, m), 7.55-7.63 (3H, m), 10 7.70-7.80 (1H, m), 7.82-7.90 (1H, m), 8.10-8.22 (2H, m)

37) 3-Methyl-4-[2-[[3-(phthalimido)prop-1-yl]oxy]-benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
15 MASS (m/z) : 774 (M+H)

38) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-3-chloro-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
20 NMR (CDCl<sub>3</sub>, δ) : 1.32-1.45 (2H, m), 1.41 (9H, s), 1.48-1.59 (2H, m), 1.62-1.91 (8H, m), 2.08-2.18 (2H, m), 2.27 (6H, s), 2.28 (3H, s), 2.30-2.40 (3H, m), 2.52-2.61 (1H, m), 2.97-3.07 (1H, m), 25 3.22-3.30 (2H, m), 3.30 (3H, s), 3.83-4.00 (3H, m), 4.30 (2H, t, J=6Hz), 4.57-4.68 (1H, m), 6.60-6.63 (2H, m), 6.87-6.90 (1H, m), 7.02-7.15 (3H, m), 7.46-7.57 (2H, m), 8.20-8.22 (1H, m), 30 8.40 (1H, d, J=7Hz)

39) 3-Ethoxy-4-[2-[[3-(phthalimido)prop-1-yl]oxy]-benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
35 MASS (m/z) : 804 (M+H)

- 148 -

- 40) 3-(3-tert-Butoxycarbonylaminoprop-1-yl)oxy-4-[2-[(3-tert-butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
5 NMR (CDCl<sub>3</sub>, δ) : 1.40 and 1.43 (total 18H, s), 1.49-1.60 (2H, m), 1.62-1.98 (6H, m), 2.00-2.10 (2H, m), 2.27 (3H, s), 2.29 (3H, s), 2.31-2.41 (6H, m), 3.17-3.29 (4H, m), 3.30 (3H, s), 3.45-3.50 (2H, m), 3.59-3.69 (2H, m), 3.84-4.05 (4H, m), 4.22-4.30 (2H, m), 5.04 (2H, br), 6.55-6.63 (2H, m), 6.85 (1H, d, J=7Hz), 6.93 (1H, d, J=7Hz), 6.98-7.03 (2H, m), 7.09 (1H, t, J=7Hz), 7.43 (1H, t, J=7Hz), 8.14 (1H, d, J=7Hz), 8.36 (1H, d, J=7Hz)  
10
- 41) 2-Amino-4-[2-[(3-tert-butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
15 NMR (CDCl<sub>3</sub>, δ) : 1.25-1.39 (2H, m), 1.42 and 1.46 (total 9H, s), 1.48-1.60 (2H, m), 1.62-1.93 (8H, m), 2.08-2.18 (2H, m), 2.27 and 2.28 (total 9H, s), 2.33-2.39 (3H, m), 2.50-2.60 (1H, m), 2.96-3.05 (1H, m), 3.29 (3H, s), 3.31-3.40 (2H, m), 3.85-3.98 (3H, m), 4.19 (2H, t, J=6Hz), 4.57-4.67 (1H, m), 6.57-6.59 (1H, m), 6.63 (2H, s), 6.78-6.89 (2H, m), 6.96 (1H, d, J=7Hz), 7.09 (1H, t, J=6Hz), 7.15 (1H, s), 7.40-7.46 (1H, m), 8.17 (1H, d, J=6Hz)  
20 25
- 42) 2-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-5-pyridinecarboxamide  
30 NMR (CDCl<sub>3</sub>, δ) : 1.30 (9H, s), 1.35-1.93 (12H, m), 2.10-2.22 (2H, m), 2.28 (9H, s), 2.30-2.40 (3H, m), 2.50-2.62 (1H, m), 2.95-3.08 (1H, m), 3.33  
35

- 149 -

(3H, s), 3.38-3.49 (2H, m), 3.82-3.98 (4H, m),  
4.29 (2H, t, J=6Hz), 4.57-4.67 (1H, m), 6.60-  
6.62 (2H, m), 6.90 (1H, d, J=6Hz), 6.99 (1H, d,  
J=7Hz), 7.09 (1H, t, J=7Hz), 7.44-7.55 (2H, m),  
8.13-8.21 (2H, m), 8.39 (1H, s)

#### Example 14

To an ice bath cooled solution of 4-[2-[(3-aminoprop-  
1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-  
methyloxy]phenyl]benzamide (200 mg) in dichloromethane (10 ml) were added  
triethylamine (36.2 mg) and acetic anhydride (36.5 mg) and  
the mixture was stirred at ambient temperature for 4  
hours. The reaction mixture was washed successively with  
water (10 ml), saturated aqueous sodium hydrogen carbonate  
solution (10 ml) and brine (10 ml), and dried over  
magnesium sulfate. The solvent was evaporated to give  
4-[2-[(3-acetylaminoprop-1-yl)oxy]benzoyl]amino-N-methyl-  
N-[2-[5-(4-methyloxy]phenyl]benzamide (201 mg) as a colorless amorphous.

NMR (CDCl<sub>3</sub>, δ) : 1.51 (2H, m), 1.62-1.86 (4H, m),  
1.93 (3H, s), 2.11 (2H, m), 2.29 (3H, s), 2.30-  
2.40 (6H, m), 3.35 (3H, s), 3.40-3.50 (4H, m),  
3.59 (2H, m), 3.92 (2H, m), 4.18 (2H, t,  
J=7.5Hz), 6.28 (1H, m), 6.75-6.83 (2H, m), 6.94-  
7.17 (4H, m), 7.31 (2H, d, J=8.5Hz), 7.40-7.49  
(3H, m), 8.08 (1H, d, J=7Hz), 9.18 (1H, s)

#### Example 15

To a mixture of 4-[2-[(3-aminoprop-1-  
yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methyloxy]phenyl]benzamide (220 mg) and  
37% aqueous formaldehyde (290 mg) in a mixture of methanol  
(10 ml) and acetic acid (0.2 ml) was added sodium  
cyanoborohydride (44.8 mg) and the mixture was stirred at

- 150 -

ambient temperature for 4 hours. The reaction mixture was diluted with chloroform (20 ml) and the solution was washed successively with saturated aqueous sodium hydrogen carbonate solution (20 ml), water (10 ml) and brine (10 ml). The organic phase was dried over magnesium sulfate and the solvent was evaporated to give 4-[2-[(3-dimethylaminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (215 mg) as a colorless amorphous.

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.55 (2H, m), 1.73 (2H, m), 1.84 (2H, m), 2.11 (2H, m), 2.20 (6H, s), 2.30 (3H, s), 2.32-2.40 (6H, m), 2.46 (2H, t, J=7.5Hz), 3.35 (3H, s), 3.49 (2H, m), 3.62 (2H, m), 4.24 (2H, t, J=7.5Hz), 6.74-6.83 (2H, m), 6.97-7.03 (2H, m), 7.07-7.16 (2H, m), 7.32 (2H, d, J=8.5Hz), 7.42-7.50 (3H, m), 8.22 (2H, d, J=7Hz)

#### Example 16

To a solution of 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (250 mg) was added 4N hydrogen chloride in ethyl acetate (1 ml) and the solution was stirred at ambient temperature for 10 minutes. The white solid was filtered and dried under reduced pressure to give 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride (205 mg) as a white powder.

NMR (D<sub>2</sub>O,  $\delta$ ) : 1.40 (2H, m), 1.59 (2H, m), 1.70 (2H, m), 2.09 (2H, m), 2.42 (2H, t, J=7.5Hz), 2.92 (3H, s), 2.96-3.17 (6H, m), 3.24 (3H, s), 3.41-3.59 (2H, m), 3.69 (1H, m), 3.82 (1H, m), 4.04-4.20 (3H, m), 4.53 (1H, m), 6.72 (1H, d, J=7Hz), 6.81 (1H, t, J=7Hz), 6.93-7.60 (11H, m)

- 151 -

Example 17

The following compounds were obtained according to a similar manner to that of Example 16.

- 5      1) 4-[2-[(3-Acetylaminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride
- NMR (D<sub>2</sub>O, δ) : 1.38 (2H, m), 1.50-1.68 (4H, m), 1.48 (2H, m), 1.81 (3H, s), 2.42 (2H, m), 2.90 (3H, s), 2.97-3.15 (6H, m), 3.24 (3H, s), 3.40-3.61 (4H, m), 3.71-3.92 (2H, m), 4.14 (1H, m), 4.54 (1H, m), 6.62-6.77 (2H, m), 6.79-6.90 (2H, m), 7.00 (1H, m), 7.11 (1H, m), 7.19-7.33 (5H, m), 7.59 (1H, d, J=7Hz)
- 10
- 15      2) 4-[2-[(3-Dimethylaminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- NMR (D<sub>2</sub>O, δ) : 1.51 (2H, m), 1.67 (2H, m), 1.81 (2H, m), 2.22 (2H, m), 2.53 (2H, t, J=7.5Hz), 2.65 (6H, s), 2.82 (3H, s), 3.00-3.17 (2H, m), 3.23 (2H, t, J=7.5Hz), 3.37 (3H, s), 3.89 (1H, m), 4.13 (1H, m), 4.07-4.20 (3H, m), 4.58 (1H, m), 6.92-7.00 (2H, m), 7.11-7.18 (2H, m), 7.26-7.48 (6H, m), 7.54-7.60 (2H, m)
- 20
- 25
- 30      3) 4-[2-[(4-Methylpiperazin-1-yl)carbonylmethoxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- NMR (D<sub>2</sub>O, δ) : 1.36 (2H, m), 1.53 (2H, m), 1.66 (2H, m), 2.98 (6H, s), 2.91-3.25 (10H, m), 3.30 (3H, s), 3.37-3.69 (4H, m), 3.77-3.96 (2H, m), 4.35-4.56 (2H, m), 4.82 (2H, s), 6.75 (1H, d, J=7Hz), 6.84 (1H, t, J=7Hz), 6.92 (1H, d, J=7Hz), 7.03-
- 35

- 152 -

7.15 (2H, m), 7.22 (1H, d, J=7Hz), 7.29 (2H, d, J=8.5Hz), 7.43-7.58 (3H, m), 7.80 (1H, d, J=7Hz).

- 5 4) 4-[2-(3-Piperidinoprop-1-yloxy)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- 10 NMR (D<sub>2</sub>O, δ) : 1.42-1.67 (10H, m), 1.78 (2H, m), 2.20 (2H, m), 2.51 (2H, t, J=7.5Hz), 2.65 (2H, m), 2.94 (3H, s), 2.95-3.21 (6H, m), 3.32 (2H, m), 3.35 (3H, s), 3.57 (2H, m), 3.92-4.04 (2H, m), 4.16-4.25 (4H, m), 6.91-6.99 (2H, m), 7.08-7.17 (2H, m), 7.23-7.47 (6H, m), 7.52-7.60 (2H, m)
- 15 5) 4-[2-[2-(Dimethylamino)eth-1-yloxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- 20 NMR (D<sub>2</sub>O, δ) : 1.52 (2H, m), 1.68 (2H, m), 1.81 (2H, m), 2.52 (2H, t, J=7.5Hz), 2.82 (6H, s), 2.93 (3H, s), 2.97-3.21 (4H, m), 3.37 (3H, s), 3.48-3.62 (2H, m), 3.87 (1H, m), 4.01 (1H, m), 4.24 (1H, m), 4.47 (2H, m), 4.57 (1H, m), 6.92-7.00 (2H, m), 7.13-7.48 (8H, m), 7.52-7.62 (2H, m)
- 25 6) 4-[2-(3-Aminoprop-1-yl)oxy]benzoylamino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylaminoprop-1-yloxy]phenyl]benzamide dihydrochloride
- 30 NMR (D<sub>2</sub>O, δ) : 2.01 (2H, m), 2.17 (2H, m), 2.91 (3H, s), 2.95-3.46 (8H, m), 3.40 (3H, s), 3.54 (2H, m), 4.02-4.16 (4H, m), 4.27 (2H, m), 6.93-7.00 (2H, m), 7.12-7.21 (2H, m), 7.26-7.37 (2H, m), 7.39-7.48 (4H, m), 7.54-7.64 (2H, m)
- 35 7) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-

- 153 -

yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

NMR (D<sub>2</sub>O,  $\delta$ ) : 1.43 (2H, m), 1.60 (2H, m), 1.72 (2H, m), 2.07 (2H, m), 2.18 (3H, s), 2.45 (2H, t, J=7.5Hz), 2.90 (3H, s), 2.92-3.13 (4H, m), 3.30 (3H, s), 3.41-3.63 (4H, m), 3.64 (3H, s), 3.82 (1H, m), 3.92 (1H, m), 4.04-4.61 (3H, m), 4.50 (1H, m), 6.66-6.74 (3H, m), 6.93-7.04 (3H, m), 7.10 (1H, d, J=7Hz), 7.41 (1H, t, J=7Hz), 7.73 (1H, d, J=7Hz), 7.95 (1H, d, J=7Hz)

8) 4-[2-(3-Aminoprop-1-yl)oxy-4-methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.48-1.96 (8H, m), 2.27 (3H, s), 2.32-2.42 (2H, m), 2.78 (3H, s), 3.11-3.22 (2H, m), 3.28 (3H, s), 3.79 (3H, s), 3.80-4.11 (2H, m), 4.22-4.32 (2H, m), 6.58-6.67 (2H, m), 6.79-6.96 (5H, m), 7.87 (1H, d, J=8Hz), 8.69-8.75 (1H, m), 9.41 (1H, br)

9) 4-[2-(3-Aminoprop-1-yl)oxy-3-methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.46-1.92 (6H, m), 2.15-2.57 (4H, m), 2.24 (3H, s), 2.30 (3H, s), 2.62-2.98 (6H, m), 2.80 (3H, s), 3.02-3.29 (4H, m), 3.28 (3H, s), 3.73-4.18 (5H, m), 4.46 (1H, br), 4.62 (1H, br), 6.56-6.68 (2H, m), 6.81-6.96 (3H, m), 7.10 (1H, dd, J=2, 8Hz), 7.30 (1H, d, J=8Hz), 7.66-7.77 (1H, m), 8.28-8.52 (4H, m), 9.65 (1H, br)

10) 4-[2-(3-Acetylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-

- 154 -

yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
hydrochloride

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.92 (6H, m), 1.91 (3H, s),  
1.96-2.25 (2H, m), 2.30 (3H, s), 2.30-2.39 (2H, m),  
2.68 (6H, s), 3.32 (3H, s), 3.35-3.47 (2H, m), 3.76  
(3H, s), 4.26 (2H, br), 4.75 (1H, br), 6.56-7.12 (6H,  
m), 7.47 (1H, br), 8.10 (1H, br), 8.39 (1H, br)

11) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
[4-(3-aminopropionyl)aminobut-1-yl]oxy-4-methylphenyl]-  
N-methylbenzamide dihydrochloride

NMR (CDCl<sub>3</sub>, δ) : 1.59-1.90 (4H, m), 2.04-2.15 (2H, m),  
2.27 (3H, s), 2.30-2.44 (2H, m), 2.87-3.08 (4H, m),  
3.21-3.38 (2H, m), 3.30 (3H, s), 3.75-3.94 (2H, m),  
3.76 (3H, s), 4.21-4.33 (2H, m), 6.55-6.68 (2H, m),  
6.86-7.10 (5H, m), 7.29-7.48 (2H, m), 8.17 (1H,  
br), 8.35 (1H, br)

12) 4-[2-(3-Guanidinoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-(4-dimethylaminopiperidin-1-  
yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
dihydrochloride

NMR (CDCl<sub>3</sub>, δ) : 1.49-1.93 (6H, m), 2.05-2.41 (8H, m),  
2.27 (3H, s), 2.75 (6H, s), 3.08 (2H, br), 3.29  
(3H, s), 3.47 (2H, br), 3.67-4.10 (4H, m), 3.77  
(3H, s), 4.27 (2H, br), 6.56-6.71 (2H, m), 6.81-  
7.09 (5H, m), 7.44 (1H, br), 7.98-8.19 (2H, m),  
8.28-8.45 (1H, m).

13) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[4-  
methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-  
yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.35-1.66 (4H, m), 1.66-1.83 (2H,  
m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.74 and  
2.76 (total 3H, s), 2.80-3.10 (4H, m), 3.18 (3H,



- 155 -

s), 3.28-3.63 (2H, m), 3.68 (3H, s), 3.77-4.18 (3H, m), 4.34-4.52 (1H, m), 6.64 (1H, d, J=9Hz), 6.75-7.12 (6H, m), 7.40 (1H, m), 7.98 (1H, d, J=9Hz), 8.23 (1H, d, J=9Hz)

5

- 14) (S)-4-[2-[(3-Amino-1-methylprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

10 NMR (DMSO-d<sub>6</sub>, δ) : 1.35 (3H, d, J=7Hz), 1.40-1.65 (4H, m), 1.66-1.82 (2H, m), 1.92-2.20 (2H, m), 2.23 (3H, s), 2.38 (2H, t, J=7Hz), 2.64 (3H, s), 2.78-3.43 (11H, m), 3.51-4.07 (7H, m), 4.93-5.09 (1H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.89 (1H, d, J=8Hz), 6.98 (1H, s), 7.04 (1H, d, J=8Hz), 7.12 (1H, dd, J=8, 8Hz), 7.36 (1H, d, J=8Hz), 7.57 (1H, dd, J=8, 8Hz), 7.98-8.35 (4H, m)

15

- 15) 4-(2-Aminobenzenesulfonyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- 20

NMR (DMSO-d<sub>6</sub>, δ) : 1.36-1.45 (2H, m), 1.50-1.59 (2H, m), 1.65-1.73 (2H, m), 2.23 and 2.29 (total 3H, s), 2.34-2.42 (4H, m), 2.77 (3H, d, J=1Hz), 2.92-3.00 (2H, m), 3.11 and 3.13 (total 3H, s), 3.19 (1H, s), 3.36-3.70 (10H, m), 4.03-4.11 (1H, m), 4.40-4.48 (1H, m), 6.44-6.50 (1H, m), 6.60-6.88 (6H, m), 6.94-7.10 (2H, m), 7.27-7.32 (1H, m)

25

ESI-MASS (m/z) : 638 (M+H)

30

- 16) (R)-4-[2-[(4-Aminobut-2-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

35 NMR (DMSO-d<sub>6</sub>, δ) : 1.37 and 1.39 (total 3H, s), 1.40-1.78 (8H, m), 1.94-2.12 (3H, m), 2.23 (3H, s), 2.30-2.40

- 156 -

5 (4H, m), 2.87-2.96 (2H, m), 3.18 (3H, s), 3.32 (3H, s), 3.46-3.58 (2H, m), 3.77 (3H, s), 3.83-3.99 (3H, m), 4.94-5.02 (1H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.88 (1H, d, J=8Hz), 6.98 (1H, s), 7.03 (1H, d, J=8Hz), 7.13 (1H, t, J=8Hz), 7.33 (1H, d, J=9Hz), 7.58 (1H, t, J=8Hz), 7.88-8.02 (2H, br), 8.04 (1H, d, J=9Hz), 8.27 (1H, d, J=8Hz)  
ESI-MASS (m/z) : 674 (M+H)

10 17) (R)-4-[2-[(4-Aminobut-2-yl)oxy]benzoyl]amino-3-methoxy-  
N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-  
yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride  
15 NMR (DMSO-d<sub>6</sub>, δ) : 1.36 and 1.38 (total 3H, s), 1.40-  
1.80 (12H, m), 1.88-2.13 (3H, m), 2.24 (3H, s),  
2.35 (2H, t, J=8Hz), 2.51 (6H, s), 2.89-3.03 (4H,  
m), 3.19 (3H, s), 3.76 (3H, s), 3.83-4.00 (3H, m),  
4.43-4.51 (1H, m), 4.96-5.03 (1H, m), 6.65 (1H, d,  
J=8Hz), 6.83 (1H, s), 6.87-6.92 (1H, m), 6.98 (1H,  
s), 7.03 (1H, d, J=8Hz), 7.14 (1H, t, J=8Hz), 7.34  
20 (1H, d, J=8Hz), 7.58 (1H, t, J=8Hz), 8.04 (1H, d,  
J=8Hz), 8.24-8.30 (1H, m)  
ESI-MASS (m/z) : 702 (M+H)

25 18) (S)-4-[2-[(4-Aminobut-2-yl)oxy]benzoyl]amino-3-methoxy-  
N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-  
yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride  
30 NMR (DMSO-d<sub>6</sub>, δ) : 1.35 and 1.38 (total 3H, s), 1.42-  
1.79 (12H, m), 1.88-2.14 (3H, m), 2.25 (3H, s),  
2.36 (2H, t, J=8Hz), 2.51 (6H, s), 2.89-3.02 (4H,  
m), 3.20 (3H, s), 3.76 (3H, s), 3.84-4.00 (3H, m),  
4.43-4.50 (1H, m), 4.97-5.03 (1H, m), 6.65 (1H, d,  
J=8Hz), 6.82 (1H, s), 6.88-6.92 (1H, m), 6.98 (1H,  
s), 7.02 (1H, d, J=8Hz), 7.15 (1H, t, J=8Hz), 7.34  
35 (1H, d, J=8Hz), 7.58 (1H, t, J=8Hz), 8.03 (1H, d,  
J=8Hz), 8.24-8.30 (1H, m)

- 157 -

ESI-MASS (m/z) : 702 (M+H)

- 19) 4-[2-(4-Aminobut-1-yl)oxybenzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]-phenylbenzamide dihydrochloride

NMR (D<sub>2</sub>O,  $\delta$ ) : 1.35-1.50 (2H, m), 1.56-1.64 (2H, m), 1.68-1.83 (4H, m), 2.47 (2H, t, J=7.5Hz), 2.82-3.12 (5H, m), 2.92 (3H, s), 3.33 (3H, s), 3.43-3.61 (3H, m), 3.81 (1H, m), 3.95 (1H, m), 6.84 (1H, d, J=7Hz), 6.91 (1H, t, J=7Hz), 7.00-7.08 (3H, m), 7.19 (1H, t, J=7Hz), 7.26-7.37 (4H, m), 7.48 (1H, t, J=7Hz), 7.62 (1H, d, J=7Hz)

- 20) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide hydrochloride

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.16 (3H, t, J=7.5Hz), 1.38-1.49 (2H, m), 1.55-1.64 (2H, m), 1.67-1.77 (2H, m), 1.98-2.08 (2H, m), 2.21 (3H, s), 2.31 (2H, t, J=7.5Hz), 2.87-2.97 (2H, m), 3.16 (3H, s), 3.80-3.98 (2H, m), 4.03 (2H, q, J=7.5Hz), 4.19 (2H, t, J=7.5Hz), 6.62 (1H, d, J=7Hz), 6.80 (1H, s), 6.98-7.07 (2H, m), 7.15 (1H, d, J=7Hz), 7.22 (2H, d, J=8Hz), 7.43-7.57 (4H, m), 7.86-8.00 (3H, br)

- 21) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.38-1.49 (2H, m), 1.52-1.62 (2H, m), 1.68-1.78 (2H, m), 1.96-2.09 (2H, m), 2.21 (3H, s), 2.38 (2H, t, J=7.5Hz), 2.73 (3Hx1/2, s), 2.75 (3Hx1/2, s), 2.81-3.07 (4H, m), 3.15 (3H, s), 3.30-3.54 (4H, m), 3.81-4.21 (5H, m), 4.45 (1H, m), 6.65 (1H, d, J=7Hz), 6.81 (1H, s), 6.99-7.08 (2H, m), 7.15 (1H, d, J=7Hz), 7.22 (2H, d, J=8Hz), 7.45-

- 158 -

7.60 (4H, m), 8.04 (2H, br)

- 22) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylprop-1-yl]oxy]-phenylbenzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.90-2.09 (4H, m), 2.48-2.59 (2H, m), 2.72 (3Hx1/2, s), 2.73 (3Hx1/2, s), 2.83-3.10 (4H, m), 3.20 (3H, s), 3.33-3.56 (3H, m), 3.88-4.09 (3H, m), 4.18 (2H, t, J=7.5Hz), 4.47 (1H, m), 4.80 (1H, m), 6.87 (1H, t, J=7Hz), 6.98 (1H, d, J=7Hz), 7.04 (1H, t, J=7Hz), 7.11-7.26 (5H, m), 7.44-7.59 (4H, m), 8.05 (2H, br)

- 23) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-N-methyl-N-[4-methyl-2-(4-methylpiperazin-1-ylcarbonyl)phenylmethoxy]-phenylbenzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.97-2.07 (2H, m), 2.26 (3H, s), 2.75 (3Hx1/2, s), 2.77 (3Hx1/2, s), 2.82-2.95 (2H, m), 3.02-3.14 (2H, m), 3.21 (3H, s), 3.30-3.49 (4H, m), 3.97-4.21 (4H, m), 5.09 (1H, d, J=14Hz), 5.20 (1H, d, J=14Hz), 6.70 (1H, d, J=7Hz), 6.93 (1H, s), 7.02-7.25 (5H, m), 7.43-7.57 (8H, m), 7.92-8.04 (3H, br)

- 24) 4-[2-(3-Hydroxyprop-1-yl)oxybenzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]-phenylbenzamide hydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.39-1.50 (2H, m), 1.52-1.62 (2H, m), 1.69-1.79 (2H, m), 1.84-1.93 (2H, m), 2.40 (2H, t, J=7.5Hz), 2.70 (3Hx1/2, s), 2.72 (3Hx1/2, s), 2.82-3.07 (4H, m), 3.19 (3H, s), 3.28-3.60 (4H, m), 3.80-3.98 (2H, m), 4.10 (1H, m), 4.17 (2H, t, J=7.5Hz), 4.45 (1H, m), 6.85 (1H, t, J=7Hz), 6.98 (1H, d, J=7Hz), 7.03 (1H, t, J=7Hz), 7.13-7.24 (5H, m), 7.43-7.54 (3H, m), 7.62 (1H, d, J=7Hz)

- 159 -

- 25) 4-[2-(4-Hydroxy-1-butyn-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]-phenylbenzamide hydrochloride

5 NMR (DMSO-d<sub>6</sub>, δ) : 1.42-1.52 (2H, m), 1.54-1.64 (2H, m), 1.70-1.82 (2H, m), 2.37-2.47 (6H, m), 2.49 (3H, s), 2.51 (3H, s), 2.84-3.05 (2H, m), 3.32-3.46 (4H, m), 3.84-3.98 (2H, m), 4.08 (1H, m), 4.47 (1H, m), 6.84 (1H, t, J=7Hz), 6.97 (1H, d, J=7Hz), 7.13-7.25 (4H, m), 7.41-7.53 (6H, m)

10

- 26) 4-[2-(4-Aminobut-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]-phenylbenzamide dihydrochloride

15 NMR (DMSO-d<sub>6</sub>, δ) : 1.39-1.62 (8H, m), 1.67-1.80 (2H, m), 2.39 (3H, t, J=7.5Hz), 2.50 (3H, s), 2.63-2.73 (4H, m), 2.81-3.08 (2H, m), 3.18 (3H, s), 3.31-3.42 (4H, m), 3.85-4.00 (2H, m), 4.04 (1H, m), 4.43 (1H, m), 6.84 (1H, t, J=7Hz), 6.99 (1H, d, J=7Hz), 7.11-7.42 (6H, m), 7.50-7.56 (2H, m), 7.75-7.91 (2H, m)

20

- 27) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yl]oxy]phenylbenzamide hydrochloride

25 NMR (DMSO-d<sub>6</sub>, δ) : 1.40-1.51 (2H, m), 1.53-1.62 (2H, m), 1.69-1.80 (2H, m), 1.98 (3H, s), 1.98-2.03 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.71 (3Hx1/2, s), 2.74 (3Hx1/2, s), 2.83-3.05 (2H, m), 3.31-3.50 (3H, m), 3.56 (2H, t, J=7.5Hz), 3.72 (3H, s), 3.81-4.11 (5H, m), 4.32 (2H, t, J=7.5Hz), 4.43 (1H, m), 6.65 (1H, d, J=7Hz), 6.81 (1H, s), 6.87-6.95 (2H, m), 7.05 (1H, d, J=7Hz), 7.11 (1H, t, J=7Hz), 7.26 (1H, d, J=7Hz), 7.54 (1H, t, J=7Hz), 8.03 (1H, d, J=7Hz), 8.28 (1H, d, J=7Hz)

30

35

- 160 -

- 28) 3-Methoxy-4-(2-hydroxybenzoyl)amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]-phenylmethoxy]phenylbenzamide hydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 2.23 (3H, s), 2.75 (3Hx1/2, s),  
2.77 (3Hx1/2, s), 2.97-3.15 (2H, m), 3.21 (3H, s),  
3.24-3.80 (6H, m), 5.06 (1H, d, J=14Hz), 5.19 (1H,  
d, J=14Hz), 6.70 (1H, d, J=7Hz), 6.90-7.01 (3H, m),  
7.10 (1H, d, J=7Hz), 7.22 (2H, d, J=8Hz), 7.41 (1H,  
d, J=7Hz), 7.44-7.55 (7H, m), 7.87 (1H, d, J=7Hz)

- 29) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]phenylbenzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 2.06-2.19 (2H, m), 2.23 (3H, s),  
2.75 (3H, s), 2.87-2.98 (2H, m), 3.02-3.15 (2H, m),  
3.23 (3H, s), 3.32-3.49 (2H, m), 3.65 (3H, s),  
3.71-3.96 (4H, m), 4.29-4.40 (2H, m), 5.04 (1H, d,  
J=14Hz), 5.20 (1H, d, J=14Hz), 6.76 (1H, d, J=7Hz),  
6.88 (1H, d, J=7Hz), 6.90-6.98 (2H, m), 7.09-7.19  
(2H, m), 7.28 (1H, d, J=7Hz), 7.50-7.62 (2H, m),  
7.98-8.15 (4H, m), 8.23 (1H, d, J=7Hz)

- 30) 3-Methoxy-4-[2-(3-aminoprop-1-yl)oxy]phenylmethyl]-amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide trihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.35-1.47 (2H, m), 1.49-1.59 (2H,  
m), 1.64-1.74 (2H, m), 2.00-2.10 (2H, m), 2.22 (3H,  
s), 2.30-2.38 (2H, m), 2.69 (3Hx1/2, s), 2.73  
(3Hx1/2, s), 2.82-3.03 (6H, m), 3.09 (3H, s), 3.29-  
3.41 (2H, m), 3.53 (3H, s), 3.83-4.12 (6H, m), 4.22  
(2H, s), 4.70 (1H, br), 6.21 (1H, d, J=7Hz), 6.58-  
6.66 (2H, m), 6.71-6.99 (5H, m), 7.09 (1H, d,  
J=7Hz), 7.20 (1H, t, J=7Hz), 8.02 (2H, br d)

- 161 -

- 31) 4-(2-Dimethylamino-4-methyl)phenoxyethyl-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.37-1.47 (2H, m), 1.50-1.61 (2H, m), 1.67-1.80 (2H, m), 2.20 (3H, s), 2.29 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.71 (3Hx1/2, s), 2.73 (3Hx1/2, s), 2.80-3.58 (4H, m), 3.03 (6H, s), 3.17 (3H, s), 3.72-4.48 (6H, m), 5.21 (2H, s), 6.62 (1H, d, J=7Hz), 6.78 (1H, s), 6.91 (1H, d, J=7Hz), 7.02 (1H, d, J=7Hz), 7.11 (1H, d, J=7Hz), 7.26 (2H, d, J=8Hz), 7.37 (2H, d, J=8Hz), 7.70 (1H, d, J=7Hz)
- 32) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[4-(4-methylpiperazin-1-yl)carbonyl]-phenyleth-1-yl]phenylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 2.06-2.19 (2H, m), 2.55-3.12 (10H, m), 2.71 (3Hx1/2, s), 2.73 (3Hx1/2, s), 3.18 (3H, s), 3.23-3.48 (2H, m), 3.66 (3H, s), 3.66-3.81 (2H, m), 4.30-4.40 (2H, m), 6.86-6.90 (2H, m), 7.11 (1H, t, J=7Hz), 7.20-7.42 (9H, m), 7.59 (1H, t, J=7Hz), 8.01 (1H, d, J=7Hz), 8.08 (2H, br), 8.27 (1H, d, J=7Hz)
- 33) 4-[2-(3-Aminoprop-1-yl)thiobenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yl]oxy]phenylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.40-1.51 (3H, m), 1.52-1.63 (2H, m), 1.70-1.88 (4H, m), 2.23 (3H, s), 2.40 (3H, t, J=7.5Hz), 2.71 (3Hx1/2, s), 2.72 (3Hx1/2, s), 2.80-2.91 (2H, m), 2.94-3.06 (2H, m), 3.17 (3H, s), 3.32-3.67 (8H, m), 3.60 (3H, s), 3.81-4.10 (3H, m), 4.41 (1H, m), 6.65 (1H, d, J=7Hz), 6.82 (1H, s), 6.86-6.92 (2H, m), 7.02 (1H, d, J=7Hz), 7.27 (1H, t, J=7Hz), 7.41-7.52 (3H, m), 7.71 (1H, d, J=7Hz), 9.37 (1H, s)

- 162 -

- 34) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-(4-dimethylaminopiperidin-1-yl)carbonyl]phenylmethoxy-4-methyl]phenyl-N-methylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.57-1.73 (2H, m), 2.00-2.20 (4H, m), 2.23 (3H, s), 2.70 (3H, s), 2.71 (3H, s), 2.87-3.05 (3H, m), 3.24 (3H, s), 3.33-3.50 (1H, m), 3.66 (3H, s), 3.71-4.05 (4H, m), 4.37 (2H, t, J=7.5Hz), 5.02 (1H, d, J=14Hz), 5.20 (1H, d, J=14Hz), 6.73 (1H, d, J=7Hz), 6.86 (1H, d, J=7Hz), 6.96 (2H, s), 7.10-7.19 (2H, m), 7.29 (1H, d, J=7Hz), 7.43-7.52 (4H, m), 7.58 (1H, t, J=7Hz), 8.00 (1H, d, J=7Hz), 8.03 (1H, d, J=7Hz)
- 35) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonyl-methoxyprop-1-yl]oxy]phenylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.94-2.04 (2H, m), 2.10-2.20 (2H, m), 2.71 (3Hx1/2, s), 2.23 (3Hx1/2, s), 2.84-3.10 (6H, m), 3.21 (3H, s), 3.31-3.50 (2H, m), 3.57-3.81 (4H, m), 3.74 (3H, s), 3.90-4.01 (2H, m), 4.20 (2Hx1/2, s), 4.22 (2Hx1/2, d), 4.35 (2H, t, J=7.5Hz), 6.82-6.97 (3H, m), 7.01 (1H, d, J=7Hz), 7.10-7.28 (4H, m), 7.58 (1H, t, J=7Hz), 8.03 (1H, d, J=7Hz), 8.27 (1H, d, J=7Hz)
- 36) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[(E)-5-(4-dimethylaminopiperidin-1-yl)carbonyl-4-penten-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.36-1.63 (2H, m), 1.84-1.92 (2H, m), 1.97-2.08 (2H, m), 2.10-2.22 (2H, m), 2.22 (3H, s), 2.29-2.43 (2H, m), 2.63 (3H, s), 2.65 (3H, s), 2.70-2.86 (2H, m), 2.88-3.00 (2H, m), 3.14 (3Hx1/2, s), 3.17 (3Hx1/2, s), 3.28-3.42 (2H, m), 3.71 (3H, s), 3.84-4.06 (2H, m), 4.37 (2H, t, J=7.5Hz), 4.51



- 163 -

(1H, m), 6.52 (1H, d, J=15Hz), 6.60 (1H, m), 6.73-7.07 (5H, m), 7.13 (1H, t, J=7Hz), 7.27 (1H, d, J=7Hz), 7.56 (1H, t, J=7Hz), 8.01 (1H, d, J=7Hz), 8.30 (1H, d, J=7Hz)

5

37) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperidin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide hydrochloride

10 NMR (CDCl<sub>3</sub>, δ) : 0.88 (3H, d, J=7.5Hz), 0.90-1.10 (2H, m), 1.34-1.61 (6H, m), 1.70-1.80 (2H, m), 2.10-2.20 (2H, m), 2.23 (3H, s), 2.30 (2H, t, J=7.5Hz), 2.45 (1H, m), 2.85-3.00 (3H, m), 3.18 (3H, s), 3.74 (3H, s), 3.75-4.02 (4H, m), 4.38 (2H, t, J=7.5Hz), 4.78 (1H, m), 6.65 (1H, d, J=7Hz), 6.82 (1H, s), 15 6.88 (1H, d, J=7Hz), 6.98 (1H, s), 7.02 (1H, d, J=7Hz), 7.13 (1H, t, J=7Hz), 7.26 (1H, d, J=7Hz), 7.59 (1H, t, J=7Hz), 8.00 (1H, d, J=7Hz), 8.22 (1H, d, J=7Hz)

20 38) 4-(2,4-Dimethoxybenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide hydrochloride

25 NMR (DMSO-d<sub>6</sub>, δ) : 1.40-1.51 (2H, m), 1.51-1.64 (2H, m), 1.69-1.82 (2H, m), 2.22 (3H, s), 2.38 (2H, t, J=7.5Hz), 2.73 (3H, s), 2.81-3.09 (4H, m), 3.19 (3H, s), 3.25-3.50 (2H, m), 3.76 (6H, sx2), 3.77-4.15 (3H, m), 4.00 (3H, s), 4.44 (1H, m), 6.64 (1H, d, J=7Hz), 6.81 (1H, s), 6.88-6.95 (2H, m), 7.03 (1H, d, J=7Hz), 7.12-7.23 (2H, m), 7.57 (1H, m), 30 8.29 (1H, d, J=7Hz)

39) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(3-aminoprop-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide dihydrochloride

35 NMR (DMSO-d<sub>6</sub>, δ) : 2.00-2.11 (2H, m), 2.13-2.20 (2H,

- 164 -

5 m), 2.25 (3H, s), 2.87-3.00 (4H, m), 3.19 (3H, s), 3.77 (3H, s), 3.89-4.10 (2H, m), 4.36 (2H, t, J=7.5Hz), 6.69 (1H, d, J=7Hz), 6.82 (1H, s), 6.89 (1H, d, J=7Hz), 7.04 (1H, s), 7.05 (1H, d, J=7Hz), 7.15 (1H, d, J=7Hz), 7.38 (1H, d, J=7Hz), 7.56 (1H, t, J=7Hz), 8.01 (1H, d, J=7Hz), 8.28 (1H, d, J=7Hz)

10 40) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(4-aminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide dihydrochloride

15 NMR (DMSO-d<sub>6</sub>, δ) : 1.66-1.85 (4H, m), 2.10-2.20 (2H, m), 2.22 (3H, s), 2.80-3.01 (4H, m), 3.18 (3H, s), 3.75 (3H, s), 3.81-4.03 (2H, m), 4.36 (2H, t, J=7.5Hz), 6.64 (1H, d, J=7Hz), 6.34 (1H, s), 6.90 (1H, d, J=7Hz), 6.96 (1H, s), 7.01 (1H, d, J=7Hz), 7.14 (1H, t, J=7Hz), 7.27 (1H, d, J=7Hz), 7.57 (1H, t, J=7Hz), 8.00 (1H, d, J=7Hz), 8.25 (7H, d, J=7Hz)

20 41) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(4-acetylaminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide hydrochloride

25 NMR (DMSO-d<sub>6</sub>, δ) : 1.49-1.59 (2H, m), 1.67-1.77 (2H, m), 1.80 (3H, s), 2.06-2.20 (2H, m), 2.21 (3H, s), 2.86-3.00 (2H, m), 3.03-3.13 (2H, m), 3.18 (3H, s), 3.74 (3H, s), 3.80-4.02 (2H, m), 4.35 (2H, t, J=7.5Hz), 6.64 (1H, d, J=7Hz), 7.82 (1H, s), 7.88 (1H, d, J=7Hz), 7.96 (1H, s), 7.02 (1H, d, J=7Hz), 7.13 (1H, t, J=7Hz), 7.26 (1H, d, J=7Hz), 7.57 (1H, t, J=7Hz), 8.00 (1H, d, J=7Hz), 8.23 (1H, d, J=7Hz)

30 42) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(4-aminoacetylaminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide dihydrochloride

35 NMR (DMSO-d<sub>6</sub>, δ) : 1.53-1.64 (2H, m), 1.70-1.81 (2H, m), 2.09-2.21 (2H, m), 2.22 (3H, s), 2.86-2.98 (2H,

- 165 -

- 5 m), 3.11-3.23 (2H, m), 3.17 (3H, s), 3.47-3.56 (2H, m), 3.65-4.00 (2H, m), 3.76 (3H, s), 4.38 (2H, t, J=7.5Hz), 6.65 (1H, d, J=7Hz), 6.82 (1H, s), 6.89 (1H, d, J=7Hz), 6.95 (1H, s), 7.03 (1H, d, J=7Hz), 7.12 (1H, t, J=7Hz), 7.25 (1H, d, J=7Hz), 7.56 (1H, t, J=7Hz), 8.00 (1H, d, J=7Hz) 8.22 (1H, d, J=7Hz)
- 43) 3-Methoxy-4-[2-(piperidin-4-yl)oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide dihydrochloride
- 10 NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.49 (2H, m), 1.49-1.61 (2H, m), 1.66-1.76 (2H, m), 1.85-1.97 (2H, m), 2.20 (3H, s), 2.67 (2H, t, J=7.5Hz), 2.73 (3Hx1/2, s), 2.74 (3Hx1/2, s), 2.80-3.13 (6H, m), 3.13 (3H, s), 3.22-
- 15 3.51 (6H, m), 3.60-4.13 (3H, m), 3.74 (3H, s), 4.43 (1H, m), 4.91 (1H, m), 6.65 (1H, d, J=7Hz), 6.81 (1H, s), 6.89 (1H, d, J=7Hz), 6.96 (1H, s), 7.03 (1H, d, J=7Hz), 7.12 (1H, t, J=7Hz), 7.35 (1H, d, J=7Hz), 7.56 (1H, t, J=7Hz), 7.81 (1H, d, J=7Hz),
- 20 8.27 (1H, d, J=7Hz)
- 44) 4-[2-(3-Amino-1-methylprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide dihydrochloride
- 25 NMR (DMSO-d<sub>6</sub>, δ) : 1.35 (3H, d, J=7.5Hz), 1.40-1.63 (4H, m), 1.67-1.80 (2H, m), 1.90-2.18 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.71 (3Hx1/2, s), 2.74 (3Hx1/2, s), 2.80-3.09 (4H, m), 3.18 (3H, s), 3.30-3.52 (4H, m), 3.77 (3H, s), 3.83-4.18 (3H, m),
- 30 4.42 (1H, m), 5.01 (1H, m), 6.64 (1H, d, J=7Hz), 6.81 (1H, s), 6.89 (1H, d, J=7Hz), 6.96 (1H, s), 7.03 (1H, d, J=7Hz), 7.12 (1H, t, J=7Hz), 7.34 (1H, d, J=7Hz), 7.58 (1H, t, J=7Hz), 8.03 (1H, d, J=7Hz), 8.28 (1H, d, J=7Hz)

- 166 -

- 45) 3-Methoxy-4-[2-(pyrid-3-yl)methoxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.36-1.49 (2H, m), 1.49-1.60 (2H, m), 1.66-1.79 (2H, m), 2.20 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.68 (3Hx1/2, s), 2.70 (3Hx1/2, s), 2.80-3.10 (4H, m), 3.16 (3H, s), 3.35 (3H, s), 3.35-3.60 (2H, m), 3.79-4.11 (3H, m), 4.41 (1H, m), 5.58 (2H, s), 6.64 (1H, d, J=7Hz), 6.80-6.90 (3H, m), 7.02 (1H, d, J=7Hz), 7.16 (1H, t, J=7Hz), 7.33 (1H, d, J=7Hz), 7.57 (1H, t, J=7Hz), 7.93-8.00 (2H, m), 8.19 (1H, d, J=7Hz), 8.55 (1H, d, J=7Hz), 8.88 (1H, d, J=6Hz), 9.04 (1H, s)
- 46) 4-[2-(4-Aminobut-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.40-1.50 (2H, m), 1.50-1.61 (2H, m), 1.66-1.79 (4H, m), 1.86-1.95 (2H, m), 2.21 (3H, s), 2.39 (2H, t, J=7.5Hz), 2.73 (3Hx1/2, s), 2.75 (3Hx1/2, s), 2.79-3.10 (4H, m), 3.19 (3H, s), 3.31-3.52 (4H, m), 3.74 (3H, s), 3.82-4.12 (3H, m), 4.30 (2H, t, J=7.5Hz), 4.43 (1H, m), 6.65 (1H, d, J=7Hz), 7.81 (1H, s), 6.89 (1H, d, J=7Hz), 6.97 (1H, s), 7.03 (1H, d, J=7Hz), 7.12 (1H, t, J=7Hz), 7.30 (1H, d, J=7Hz), 7.58 (1H, d, J=7Hz), 8.04 (1H, d, J=7Hz), 8.30 (1H, d, J=7Hz)
- 47) 4-(2-Hydroxy-5-methylbenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide hydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.53-1.96 (6H, m), 2.29 (3H, s), 2.31 (3H, s), 2.33-2.40 (2H, m), 2.79 (3H, s), 3.30 (3H, s), 3.79 (3H, s), 3.80-4.03 (2H, m), 6.63 (2H, br), 6.88-6.98 (4H, m), 7.25 (1H, d, J=8Hz), 8.19

- 167 -

(1H, d, J=8Hz), 8.71 (1H, br)

48) 4-(2-Hydroxy-4-methoxybenzoyl)amino-3-methoxy-N-methyl-  
N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-  
methylphenyl]benzamide hydrochloride

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.92 (6H, m), 2.28 (3H, s),  
2.32-2.45 (2H, m), 2.64-3.05 (4H, m), 2.79 (3H, s),  
3.29 (3H, s), 3.29-3.51 (4H, m), 3.76 (3H, s), 3.80  
(3H, s), 3.81-4.05 (4H, m), 6.43-6.50 (2H, m), 6.61  
(1H, br), 6.85-6.96 (3H, m), 7.36-7.43 (1H, m),  
8.12-8.18 (1H, m), 8.58 (1H, br)

#### Example 18

The following compounds were obtained by separating the  
compounds, which were prepared according to a similar manner  
to Example 4, by using silica gel column chromatography.

1) 4-(2-Benzyloxy)benzoylamino-3-methoxy-N-[(E)-2-(4-  
carboxyphenyl)ethen-1-yl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 3.08 (3H, s), 3.41 (3H, s), 5.19 (2H,  
s), 6.47 (1H, d, J=14Hz), 6.58 (1H, d, J=14Hz), 6.73  
(2H, d, J=6Hz), 6.84 (1H, d, J=7Hz), 6.90-7.10 (5H, m),  
7.20-7.40 (8H, m), 7.71 (2H, d, J=8Hz), 8.26 (1H, d,  
J=7Hz), 8.38 (1H, d, J=7Hz)

2) 4-(2-Benzyloxy)benzoylamino-3-methoxy-N-[(Z)-2-(4-  
carboxyphenyl)ethen-1-yl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 3.09 (3H, s), 3.48 (3H, s), 5.25 (2H,  
s), 6.72-7.42 (15H, m), 7.51-7.64 (3H, m), 8.10  
(2H, d, J=8Hz), 8.22 (1H, d, J=7Hz), 8.33 (1H, d,  
J=7Hz)

#### Example 19

The following compound was obtained according to a  
similar manner to that of Example 4 by using 4-[2-

- 168 -

(acetoxy)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-(5-ethoxycarbonylpent-1-yloxy)phenyl]benzamide as a starting compound.

- 5           4-[2-(Hydroxy)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-(5-carboxypent-1-yloxy)phenyl]benzamide
- NMR (CDCl<sub>3</sub>, δ) : 1.46-1.61 (2H, m), 1.63-1.90 (4H, m),  
                          2.28 (3H, s), 2.39 (2H, t, J=7Hz), 3.33 (3H, s),  
                          3.73-4.00 (5H, m), 6.61 (2H, br s), 6.82-7.11 (5H,  
10           m), 7.35-7.53 (2H, m), 8.16 (1H, d, J=8Hz), 8.75  
                          (1H, br s)

#### Example 20

15           The following compounds were obtained according to a similar manner to that of Example 8.

- 1) 4-[2-(4-Methoxybenzyl)oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]-phenylmethoxy]phenylbenzamide
- 20           NMR (CDCl<sub>3</sub>, δ) : 2.27 (3H, s), 2.31 (3H, s), 2.35-2.53  
                          (4H, m), 3.32 (3H, s), 3.39-3.54 (2H, m), 3.67-3.85  
                          (3H, m), 3.82 (3H, s), 4.95 (1H, d, J=14Hz), 5.06  
                          (1H, d, J=14Hz), 5.12 (2H, s), 6.59-6.67 (2H, m),  
                          6.86-7.02 (5H, m), 7.07-7.21 (4H, m), 7.33-7.52  
25           (7H, m), 8.28 (1H, d, J=7Hz)
- 2) 4-(2-Benzyloxybenzoyl)amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide
- 30           NMR (CDCl<sub>3</sub>, δ) : 1.46-1.60 (2H, m), 1.63-1.92 (4H, m),  
                          2.30 (3H, s), 2.31-2.46 (6H, m), 3.28 (3H, s), 3.35  
                          (3H, s), 3.44-3.54 (2H, m), 3.58-3.69 (2H, m),  
                          3.80-4.04 (2H, m), 5.30 (2H, s), 6.73-7.22 (8H, m),  
                          7.30-7.49 (6H, m), 8.19-8.28 (1H, m), 8.38 (1H, d,  
35           J=9Hz)

- 169 -

- 3) 4-[2-(Benzyloxy)benzoyl]amino-2-chloro-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.65 (2H, m), 1.65-1.97 (4H, m),  
2.30 (3H, s), 2.32-2.48 (6H, m), 3.34 (3H, s),  
3.43-3.56 (2H, m), 3.58-3.70 (2H, m), 3.97 (2H, t, J=7Hz), 5.16 (2H, s), 6.63-6.81 (3H, m), 6.96 (1H, d, J=8Hz), 7.02-7.20 (5H, m), 7.40-7.59 (6H, m), 8.24 (1H, m)

- 4) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-amino-3-methoxy-N-methyl-N-[4-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.42 (9H, s), 1.45-1.82 (8H, m),  
2.10-2.19 (2H, m), 2.30 (3H, s), 2.31-2.41 (6H, m),  
3.27-3.35 (2H, m), 3.43-3.50 (5H, m), 3.60-3.67 (2H, m), 3.82 (3H, s), 3.90 (1H, t, J=7Hz), 4.27 (1H, t, J=7Hz), 4.75-4.82 (1H, br), 6.76 (2H, d, J=8Hz), 6.82 (1H, d, J=8Hz), 6.95-7.04 (3H, m), 7.07-7.13 (1H, m), 7.47 (1H, t, J=8Hz), 8.22 (1H, dd, J=1, 8Hz), 8.42 (1H, d, J=8Hz)

ESI-MASS (m/z) : 746 (M+H)

- 5) 4-[2-[3-(tert-Butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]-amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.45-1.60 (2H, m),  
1.65-1.74 (2H, m), 1.78-1.89 (2H, m), 2.04-2.15 (2H, m), 2.27 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.27-3.39 (2H, m), 3.33 (3H, s), 3.44-3.50 (2H, m), 3.58-3.64 (2H, m), 3.82-4.00 (2H, m), 4.19 (2H, t, J=7.5Hz), 4.86 (1H, br), 6.55-6.62 (2H, m), 6.86 (1H, d, J=7Hz), 6.97 (1H, d, J=7Hz), 7.08 (1H, t, J=7Hz), 7.31 (2H, d, J=8Hz), 7.40-7.53 (3H, m), 8.13 (1H, d, J=7Hz), 9.88 (1H, s)

- 170 -

- 6) 4-(2-Iodobenzoyl)amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.43-1.54 (2H, m), 1.61-1.70 (2H, m),  
1.74-1.86 (2H, m), 2.28 (3H, s), 2.28-2.41 (6H, m),  
3.34 (3H, s), 3.44-3.50 (2H, m), 3.52-3.59 (2H, m),  
3.73-3.99 (2H, m), 6.77-6.84 (2H, m), 7.03 (1H, d,  
J=7Hz), 7.10-7.19 (2H, m), 7.29-7.50 (5H, m), 7.80  
(1H, s), 7.89 (1H, d, J=7Hz)
- 7) 4-(2-Dimethylamino-4-methyl)phenoxyethyl-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.47-1.58 (2H, m), 1.64-1.75 (2H, m),  
1.77-1.88 (2H, m), 2.22 (3H, s), 2.25 (3H, s), 2.28  
(3H, s), 2.31-2.41 (6H, m), 2.72 (6H, s), 3.32 (3H,  
s), 3.43-3.51 (2H, m), 3.58-3.67 (2H, m), 3.79-3.97  
(2H, m), 5.02 (2H, s), 6.49-6.61 (3H, m), 6.71 (1H,  
d, J=7Hz), 7.80-7.85 (2H, m), 7.19 (2H, d, J=8Hz),  
7.28 (2H, d, J=8Hz)
- 8) 4-(2-Benzyloxy)benzoylamino-3-methoxy-N-methyl-N-[(E)-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylethen-1-yl]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.11-2.40 (4H, m), 2.17 (3H, s), 3.11  
(3H, s), 3.18-3.38 (2H, m), 3.44 (3H, s), 3.49-3.68  
(2H, m), 5.27 (2H, s), 6.41 (1H, d, J=14Hz), 6.56  
(1H, d, J=14Hz), 6.70 (2H, d, J=8Hz), 6.88-7.48  
(16H, m), 8.26 (1H, d, J=7Hz), 8.38 (1H, d, J=7Hz)
- 9) 3-Methoxy-4-[2-[3-(tert-butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperidin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 0.93 (3H, d, J=7.5Hz), 0.98-1.14 (2H,  
m), 1.40 (9H, s), 1.42-1.87 (8H, m), 2.07-2.17 (2H,  
m), 2.25 (3H, s), 2.32 (2H, t, J=7.5Hz), 2.50 (1H,



- 171 -

m), 2.97 (1H, m), 3.21-3.32 (2H, m), 3.32 (1H, s),  
3.79 (1H, s), 3.79-4.00 (4H, m), 4.24 (2H, t,  
J=7.5Hz), 4.55 (1H, m), 4.84 (1H, m), 6.59 (1H, d,  
J=7Hz), 6.63 (1H, s), 6.85 (1H, d, J=7Hz), 6.92  
(1H, d, J=7Hz), 6.95-7.13 (3H, m), 7.45 (1H, t,  
J=7Hz), 8.20 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 5  
10  
10) 3-Methoxy-4-[2-[3-(tert-butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-N-[2-[5-[(2S)-carbamoylpyrrolidin-1-yl]carbonylpent-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide

15  
NMR (CDCl<sub>3</sub>, δ) : 1.28-2.20 (12H, m), 1.39 (9H, s),  
2.27 (3H, s), 3.19-3.25 (2H, m), 3.21 (3H, s),  
3.25-3.61 (2H, m), 3.78 (3H, s), 3.81-4.03 (2H, m),  
4.16-4.29 (2H, m), 4.57 (1H, m), 6.55-6.68 (2H, m),  
6.80-7.13 (5H, m), 7.44 (1H, t, J=7Hz), 8.20 (1H,  
d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 20  
11) 3-Methoxy-4-[2-[1-(tert-butoxycarbonyl)piperidin-4-yl]-oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

25  
NMR (CDCl<sub>3</sub>, δ) : 1.41-1.59 (2H, m), 1.46 (9H, s),  
1.69-1.94 (6H, m), 2.00-2.13 (2H, m), 2.26 (3H, s),  
2.29 (3H, s), 2.33-2.41 (8H, m), 2.96-3.17 (2H, m),  
3.31 (3H, s), 3.45-3.51 (2H, m), 3.59-3.67 (2H, m),  
3.74 (3H, s), 3.80-4.01 (2H, m), 4.68 (1H, m),  
6.58-6.63 (2H, m), 6.85 (1H, d, J=7Hz), 6.90 (1H,  
d, J=7Hz), 6.99-7.11 (2H, m), 7.35-7.61 (2H, m),  
8.19 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 30  
12) 3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)amino-1-methylprop-1-yl]oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

35  
NMR (CDCl<sub>3</sub>, δ) : 1.30 (9H, s), 1.31 (3H, d, J=7.5Hz),

- 172 -

5 1.45-2.10 (8H, m), 2.27 (3H, s), 2.29 (3H, s),  
2.32-2.43 (6H, m), 3.20-3.30 (2H, m), 3.32 (3H, s),  
3.45-3.50 (2H, m), 3.60-3.66 (2H, m), 3.79 (3H, s),  
3.82-4.00 (2H, m), 4.72 (1H, m), 6.60 (1H, d,  
J=7Hz), 6.64 (1H, s), 6.81-6.93 (2H, m), 7.00-7.11  
(3H, m), 7.43 (1H, t, J=7Hz), 8.21 (1H, d, J=7Hz),  
8.42 (1H, d, J=7Hz)

10 13) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-(5-aminocarbonylpent-1-  
yl)oxy-4-methylphenyl]benzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.42 (9H, s), 1.50-1.92 (6H, m),  
2.12-2.26 (2H, m), 2.29 (2H, t, J=5Hz), 2.30 (3H,  
s), 3.30 (2H, q, J=5Hz), 3.35 (3H, s), 3.77 (3H,  
s), 3.80-4.02 (2H, m), 4.25 (2H, t, J=5Hz), 6.61-  
6.70 (2H, m), 6.93-7.15 (6H, m), 7.41-7.51 (1H, m),  
8.20 (1H, d, J=7Hz), 8.42 (1H, d, J=7Hz)

20 14) 4-[2-[3-(tert-Butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-[5-[4-(tert-  
butoxycarbonyl)piperazin-1-yl]carbonylpent-1-yl]oxy-4-  
methylphenyl]benzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.49 (9H, s), 1.50-1.90  
(6H, m), 2.12-2.23 (2H, m), 2.30 (3H, s), 2.39 (2H,  
t, J=5Hz), 3.30 (2H, q, J=5Hz), 3.33 (3H, s), 3.35-  
3.42 (4H, m), 3.44 (4H, s), 3.55-3.62 (2H, m), 3.80  
(3H, s), 3.85-4.06 (2H, m), 4.24 (2H, t, J=5Hz),  
4.93 (1H, br), 6.57-6.66 (2H, m), 6.85-7.13 (6H,  
m), 7.44-7.52 (1H, m), 8.20 (1H, d, J=7Hz), 8.41  
30 (1H, d, J=7Hz)

35 15) 4-[2-[3-(tert-Butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-[5-morpholin-4-  
yl]carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.41 (9H, s), 1.50-1.88 (6H, m),

- 173 -

2.10-2.21 (2H, m), 2.30 (3H, s), 2.36 (2H, t, J=5Hz), 3.30 (2H, q, J=5Hz), 3.34 (3H, s), 3.47 (2H, t, J=4Hz), 3.58-3.70 (6H, m), 3.79 (3H, s), 3.84-4.03 (2H, m), 4.25 (2H, t, J=5Hz), 4.89 (1H, br), 6.56-6.68 (2H, m), 6.84-7.16 (6H, m), 7.41-7.51 (2H, m), 8.20 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

5

10

- 16) 4-[2-[3-(tert-Butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[5-(4-methylhomopiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.41 (9H, s), 1.46-1.97 (8H, m), 2.09-2.21 (2H, m), 2.29 (3H, s), 2.32 (2H, t, J=5Hz), 2.33 (3H, s), 2.52-2.66 (4H, m), 3.30 (2H, q, J=5Hz), 3.33 (3H, s), 3.50-3.69 (4H, m), 3.79 (3H, s), 3.84-4.03 (2H, m), 4.24 (2H, t, J=5Hz), 4.94 (1H, br), 6.56-6.67 (2H, m), 6.82-7.12 (6H, m), 7.40-7.49 (1H, m), 8.20 (1H, d, J=7Hz), 8.41 (1H, d, J=8Hz)

15

20

- 17) 4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[5-(2-dimethylaminoeth-1-yl)aminocarbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.42-1.59 (4H, m), 1.67-1.90 (4H, m), 1.97-2.32 (4H, m), 2.28 (3H, s), 2.34 (6H, s), 2.56 (2H, br), 3.25-3.42 (4H, m), 3.32 (2H, s), 3.50 (1H, s), 3.78-4.01 (2H, m), 3.80 (3H, s), 4.25 (2H, t, J=6Hz), 4.91 (1H, br), 6.52-6.76 (3H, m), 6.87-7.13 (7H, m), 7.45 (1H, m), 8.19 (1H, d, J=8Hz), 8.41 (1H, br)

25

30

- 18) 4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[5-[N-(3-dimethylamino-prop-1-yl)-N-methylcarbonylpent-1-yl]oxy-4-methylphenyl]benzamide

35

- 174 -

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.41 (9H, s), 1.50-1.96 (6H, m),  
2.11-2.25 (2H, m), 2.27 (3H, s), 2.30-2.43 (2H, m),  
2.50 (6H, s), 2.91 and 3.02 (total 3H, s, rotamer),  
3.08 and 3.32 (total 2H, q, rotamer, J=5Hz), 3.33  
(3H, s), 3.43 (2H, t, J=5Hz), 3.79 (3H, s), 3.83-  
4.02 (2H, m), 4.25 (2H, t, J=5Hz), 6.57-6.68 (2H,  
m), 6.82-7.13 (6H, m), 7.42-7.50 (1H, m), 8.21 (1H,  
d, J=8Hz), 8.41 (1H, d, J=8Hz)

- 5  
10 19) 4-[2-(3-tert-Butoxycarbonylamino-*prop*-1-yl)oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-[5-[bis(2-hydroxyeth-1-  
yl)amino]carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

15 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 1.55-1.89 (6H, m),  
2.11-2.20 (2H, m), 2.28 (3H, s), 2.40-2.56 (2H, m),  
3.29 (2H, t, J=5Hz), 3.40-3.57 (4H, m), 3.68-4.02  
(6H, m), 4.26 (2H, t, J=5Hz), 6.60-6.68 (2H, m),  
6.90-7.15 (6H, m), 7.42-7.51 (1H, m), 8.19 (1H, d,  
J=8Hz), 8.40 (1H, d, J=8Hz)

- 20 20) 4-[2-(3-tert-Butoxycarbonylamino-*prop*-1-yl)oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-[5-(2,2-dimethyl-  
hydrazino)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

25 NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.40 (9H, s), 1.45-1.90 (6H, m),  
2.08-2.20 (2H, m), 2.28 (3H, s), 2.30-2.45 (2H, m),  
2.51 (3H, s), 2.60 (3H, s), 3.29 (2H, t, J=5Hz),  
3.33 (3H, s), 3.75 (3H, s), 3.79-4.02 (2H, m), 4.25  
(2H, t, J=5Hz), 6.57-6.68 (2H, m), 6.80-7.14 (5H,  
m), 7.41-7.50 (1H, m), 8.21 (1H, d, J=8Hz), 8.40-  
8.48 (1H, br)

- 30 21) 4-[2-(3-tert-Butoxycarbonylamino-*prop*-1-yl)oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-[5-(carbamoylmethylamino)-  
carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

35 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.41 (9H, s), 1.50-1.90 (6H, m),  
2.12-2.19 (2H, m), 2.28 (3H, s), 2.53 (2H, t,

- 175 -

J=5Hz), 3.30 (2H, t, J=5Hz), 3.33 (3H, s), 3.80 (3H, s), 3.84-3.99 (2H, m), 4.05 (2H, br), 4.25 (2H, t, J=5Hz), 4.84 (1H, br), 6.58-6.67 (2H, m), 6.72-7.12 (6H, m), 7.42-7.50 (1H, m), 8.18-8.23 (1H, m), 8.41 (1H, d, J=8Hz)

5

22) 4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[5-(carbamoylethylamino)-carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

10

NMR (CDCl<sub>3</sub>, δ) : 1.41 (9H, s), 1.46-1.86 (6H, m), 2.12-2.25 (4H, m), 2.30 (3H, s), 2.41 (2H, t, J=5Hz), 3.30 (1H, q, J=5Hz), 3.37 (3H, s), 3.49 (1H, q, J=5Hz), 3.79 (3H, s), 3.82-4.03 (2H, m), 4.27 (2H, t, J=5Hz), 6.45-6.67 (4H, m), 6.88-7.15 (6H, m), 7.43-7.51 (1H, m), 8.20 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

15

23) 4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[5-(4-diethylamino-piperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]-benzamide

20

NMR (CDCl<sub>3</sub>, δ) : 1.12 (6H, t, J=5Hz), 1.41 (9H, s), 1.42-1.92 (6H, m), 2.10-2.18 (2H, m), 2.27 (3H, s), 2.27-2.69 (9H, m), 3.26 (2H, t, J=5Hz), 3.31 (3H, s), 3.77 (3H, s), 3.87-4.02 (4H, m), 4.23 (2H, t, J=5Hz), 6.54-6.67 (2H, m), 6.72-7.15 (6H, m), 7.42-7.51 (1H, m), 8.19 (1H, d, J=8Hz), 8.42 (1H, d, J=8Hz)

25

24) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-[2-[3-(4-methylpiperazin-1-yl)carbonylpyrid-6-yl]methoxy-4-methylphenyl]-N-methylbenzamide

30

NMR (CDCl<sub>3</sub>, δ) : 1.39 (9H, s), 2.06-2.18 (2H, m), 2.28 (3H, s), 2.31 (3H, s), 2.35-2.51 (4H, m), 3.27 (2H,

35

- 176 -

q, J=5Hz), 3.38-3.49 (2H, m), 3.41 (1H, s), 3.63 (3H, s), 3.68-3.76 (2H, m), 4.21 (2H, t, J=5Hz), 4.97 (1H, d, J=12Hz), 5.14 (1H, d, J=12Hz), 6.58 (1H, s), 6.72 (1H, d, J=8Hz), 6.91-7.11 (7H, m), 7.20-7.25 (1H, m), 7.43 (1H, dd, J=2, 8Hz), 7.68 (1H, d, J=8Hz), 8.19 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz), 8.60 (1H, s)

5

10

- 25) 4-[2-(Benzyloxy)benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-phenyl]benzamide

15

NMR (CDCl<sub>3</sub>, δ) : 1.30-1.42 (2H, m), 1.48-1.58 (2H, m), 1.63-1.93 (6H, m), 2.29 (6H, s), 2.30-2.40 (3H, m), 2.50-2.60 (1H, m), 2.95-3.06 (1H, m), 3.29 (3H, s), 3.38 (3H, s), 3.80-4.00 (4H, m), 4.57-4.70 (1H, m), 5.30 (2H, s), 6.74-7.20 (9H, m), 7.32-7.45 (5H, m), 8.20-8.37 (1H, m), 8.37-8.42 (1H, m)

20

- 26) 4-[(2-Benzyloxy)benzoyl]amino-N-methyl-N-[2-[3-(4-methylpiperazin-1-yl)carbonylprop-1-yl]oxy]-phenylbenzamide

25

NMR (CDCl<sub>3</sub>, δ) : 2.05-2.16 (2H, m), 2.28 (3H, s), 2.32-2.40 (4H, m), 2.50 (2H, t, J=7.5Hz), 3.33 (3H, s), 3.43-3.50 (2H, m), 3.59-3.65 (2H, m), 3.88-4.05 (2H, m), 5.19 (2H, s), 6.77-6.84 (2H, m), 6.95-7.02 (3H, m), 7.09-7.20 (5H, m), 7.39-7.52 (6H, m), 8.27 (1H, d, J=7Hz)

### Example 21

30

The following compounds were obtained according to similar manners to those of Examples 8 and 16.

35

- 1) 4-(6-Hydroxy-2-pyridylcarbonyl)amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

- 177 -

NMR (CDCl<sub>3</sub>, δ) : 1.47-1.58 (2H, m), 1.64-1.73 (2H, m),  
1.78-1.87 (2H, m), 2.27 (3H, s), 2.29 (3H, s),  
2.28-2.41 (8H, m), 3.33 (3H, s), 3.45-3.51 (2H, m),  
3.59-3.68 (6H, m), 3.86-3.94 (1H, br), 6.55-6.61  
(2H, m), 6.86 (1H, d, J=8Hz), 7.30-7.38 (4H, m),  
7.47-7.54 (2H, m), 8.06-8.10 (1H, m)

ESI-MASS (m/z) : 574 (M+H)

2) 4-[2-(Methoxy)benzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide hydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.36-1.66 (4H, m), 1.66-1.83 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=7Hz), 2.74 (3H, s), 2.80-3.10 (3H, m), 3.17 (3H, s), 3.23-3.53 (3H, m), 3.86 (3H, s), 3.79-3.99 (2H, m), 4.00-4.17 (1H, m), 4.37-4.52 (1H, m), 6.64 (1H, d, J=9Hz), 6.79 (1H, s), 6.98-7.09 (2H, m), 7.11-7.28 (3H, m), 7.43-7.64 (4H, m)

#### Example 22

To a solution of 4-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide (327 mg) and pyridine (80.3 mg) in dichloromethane (6 ml) was added dropwise 2-nitrobenzenesulfonyl chloride (150 mg) at ambient temperature and the mixture was stirred at ambient temperature for 5 hours. The resulting mixture was diluted with dichloromethane (10 ml) and the organic layer was washed successively with saturated sodium bicarbonate aqueous solution and brine. Drying, filtering and removal of solvents afforded a crude product. The crude product was chromatographed on silica gel (eluent; 2-4% methanol in chloroform) to give 4-(2-nitrobenzenesulfonyl)-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (460 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.47-1.82 (6H, m), 2.28 (3H, s), 2.31

- 178 -

(3H, s), 2.35-2.42 (6H, m), 3.30 (3H, s), 3.46-3.53 (5H, m), 3.60-3.68 (4H, m), 6.56-6.96 (6H, m), 7.53-7.88 (4H, m)

5     Example 23

      A solution of 4-[2-[2-[3-(phthalimido)prop-1-yl]oxy]phenyl]vinyl-3-methoxybenzoic acid (370 mg) in tetrahydrofuran (20 ml) was treated at ambient temperature with triethylamine (246 mg), N-methyl-4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]aniline (297 mg), and diphenyl phosphorochloridate (326 mg). The reaction mixture was stirred at 80°C for 18 hours. After concentration, the residue was dissolved in chloroform and washed with brine and dried over magnesium sulfate. The crude product was purified by silica gel column chromatography (SiO<sub>2</sub> 30 g, 3% methanol in chloroform) to give 4-[2-[2-[(3-(phthalimido)prop-1-yl)oxy]phenyl]vinyl]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (550 mg).

20       NMR (CDCl<sub>3</sub>, δ) : 1.47-1.95 (8H, m), 2.18-2.44 (12H, m), 3.31 and 3.34 (total 3H, s), 3.42-3.52 (2H, m), 3.57-3.72 (5H, m), 3.82-4.16 (6H, m), 6.30-7.80 (16H, m)

25     Example 24

      The following compounds were obtained according to a similar manner to that of Example 23.

30     1) 4-[N-Methyl-2-[(3-tert-butoxycarbonylamino)prop-1-yl]-oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

35       NMR (CDCl<sub>3</sub>, δ) : 1.40-1.75 (8H, m), 1.44 (9H, s), 1.89-1.97 (2H, m), 2.29 (6H, s), 2.32-2.42 (6H, m), 3.24 (6H, s), 3.26-3.34 (2H, m), 3.44-3.67 (6H, m),



- 179 -

3.77-3.88 (3H, m), 6.48-6.82 (9H, m), 6.90-6.96  
(1H, m), 7.06-7.13 (1H, m)

ESI-MASS (m/z) : 774 (M+H)

- 5           2) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-  
amino-3-methoxy-N-methyl-N-(4-benzyloxyphenyl)benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.42 (9H, s), 2.09-2.20 (2H, m),  
3.28-3.37 (2H, m), 3.48 (3H, s), 3.81 (3H, s),  
4.22-4.33 (2H, m), 4.70-4.78 (1H, br), 5.00 (2H,  
10           s), 6.82-6.88 (3H, m), 6.97-7.13 (6H, m), 7.31-7.48  
(6H, m), 8.23 (1H, d, J=8Hz), 8.44 (1H, d, J=8Hz)

ESI-MASS (m/z) : 640 (M+H)

- 15           3) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-  
amino-3-benzyloxy-N-methyl-N-cyclohexylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.01-1.12 (2H, br), 1.40 (9H, s),  
1.45-1.82 (10H, m), 2.81-3.07 (5H, m), 3.80-3.89  
(2H, m), 4.40-4.49 (1H, m), 5.18 (2H, s), 6.94 (1H,  
20           d, J=8Hz), 7.02 (1H, d, J=8Hz), 7.07-7.15 (2H, m),  
7.35-7.48 (6H, m), 8.27 (1H, d, J=8Hz) 8.68 (1H, d,  
J=8Hz)

ESI-MASS (m/z) : 616 (M+H)

- 25           4) 4-[(2-Benzyloxy)benzoyl]amino-3-chloro-N-methyl-N-[2-[5-  
(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-  
methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.30-1.45 (2H, m), 1.45-1.57 (2H, m),  
1.62-1.93 (6H, m), 2.22-2.40 (12H, m), 2.50-2.63  
(1H, m), 2.95-3.08 (1H, m), 3.31 (3H, s), 3.80-4.00  
30           (4H, m), 4.58-4.70 (1H, m), 5.37 (2H, s), 6.56-6.62  
(2H, m), 6.83-6.88 (1H, m), 7.02-7.13 (3H, m),  
7.36-7.47 (7H, m), 8.27 (1H, d, J=7Hz), 8.42 (1H,  
d, J=7Hz)

- 35           5) 4-[N-[2-[(3-tert-Butoxycarbonylaminoprop-1-

- 180 -

yl)oxy]phenyl]-tert-butoxycarbonylamino]methyl-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.30 and 1.33 (total 9H, s), 1.43 (9H, s), 1.49-1.60 (2H, m), 1.62-1.98 (6H, m), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.42 (6H, m), 3.20-3.29 (2H, m), 3.32 (3H, s), 3.39 (1H, s), 3.46-3.55 (4H, m), 3.62 (2H, br), 3.82 (1H, br), 3.88-4.03 (3H, m), 6.50-6.60 (2H, m), 6.65-7.00 (6H, m), 7.06-7.22 (2H, m)

6) 4-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl)oxy]phenoxy]-methyl-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.37 (9H, s), 1.47-1.57 (2H, m), 1.66-1.73 (2H, m), 1.73-1.88 (2H, m), 1.93-2.02 (2H, m), 2.28 (3H, s), 2.30 (3H, s), 2.32-2.40 (6H, m), 3.32 (3H, m), 3.25-3.38 (2H, m), 3.47-3.50 (2H, m), 3.62-3.67 (2H, m), 3.70 (3H, s), 3.80-3.88 (1H, m), 3.90-3.98 (2H, m), 4.07-4.17 (2H, m), 5.10 (2H, s), 5.50 (1H, br), 6.53-6.60 (2H, m), 6.70-6.90 (7H, m), 7.15-7.20 (1H, m)

7) 3-Benzoyloxy-4-[2-[(3-tert-butoxycarbonylamino)prop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-benzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.45-1.85 (10H, m), 2.28 (3H, s), 2.29 (3H, s), 2.32-2.39 (6H, m), 2.90-2.98 (2H, m), 3.30 (3H, s), 3.47-3.49 (2H, m), 3.60-3.63 (2H, m), 3.77-3.98 (4H, m), 4.97 (2H, s), 6.56-6.60 (2H, m), 6.80 (1H, d, J=7Hz), 6.89-6.97 (2H, m), 7.04-7.12 (2H, m), 7.33-7.45 (6H, m), 8.19 (1H, d, J=6Hz), 8.41 (1H, d, J=7Hz)

- 181 -

8) 2-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl]oxy]benzoyl]-  
amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-  
yl)carbonylpent-1-yloxy]-4-methylphenyl]-5-  
thiophenecarboxamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.37 (9H, s), 1.48-1.62 (2H, m),  
1.62-1.76 (6H, m), 1.97-2.11 (2H, m), 2.17-2.38  
(9H, m), 2.39 (3H, s), 3.31 (3H, s), 3.33-3.65 (6H,  
m), 3.87 (1H, br), 3.94 (1H, br), 4.02 (1H, s),  
10 4.13-4.20 (2H, m), 6.40-6.57 (2H, m), 6.74-6.82  
(2H, m), 6.92-7.14 (3H, m), 7.40-7.52 (1H, m),  
8.10-8.27 (1H, m)

#### Example 25

15 A solution of (S)-4-[2-[1-methyl-3-(phthalimido)prop-1-  
yl]oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-  
methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide  
(1.1 g) in methanol (30 ml) was stirred and treated with 40%  
methylamine in methanol (10 ml). The reaction mixture was  
20 refluxed for 30 minutes. Then the solvent was concentrated  
and purified by silica gel column chromatography (SiO<sub>2</sub> 40 g,  
chloroform/methanol/ammonia = 90/10/0.5) to give (S)-4-[2-  
[(3-amino-1-methylprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-  
methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-  
carbonylpent-1-yloxy]phenyl]benzamide.

25 NMR (CDCl<sub>3</sub>, δ) : 1.42 (3H, d, J=7Hz), 1.46-1.92 (9H,  
m), 1.98-2.16 (1H, m), 2.20-2.45 (12H, m), 2.86  
(2H, t, J=7Hz), 3.32 (3H, s), 3.42-3.53 (2H, m),  
3.57-3.67 (2H, m), 3.79 (3H, s), 3.82-4.03 (2H, m),  
4.73-4.90 (1H, m), 6.51-6.68 (2H, m), 6.79-6.95  
30 (2H, m), 6.98-7.12 (3H, m), 7.37-7.49 (1H, m), 8.21  
(1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

#### Example 26

35 A solution of 4-[2-[(3-tert-butoxycarbonylamino)prop-1-  
yl]oxy]benzoyl]amino-3-methoxy-N-methyl-N-[2-(5-carboxypent-

- 182 -

1-yloxy)-4-methylphenyl]benzamide (3.5 g) in ethyl acetate (30 ml) was treated at ambient temperature with triethylamine (575 mg), N-methylpiperazine (569 mg), and diphenylphosphoryl azide (1.56 g). The reaction mixture was stirred at the same temperature for 17 hours. The reaction mixture was washed with brine and dried over magnesium sulfate. The crude product was purified by silica gel column chromatography (SiO<sub>2</sub> 100 g, 3% methanol in chloroform) to give 4-[2-[(3-tert-butoxycarbonylamino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (2.93 g).

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.42-1.60 (2H, m), 1.62-1.90 (4H, m), 2.06-2.20 (2H, m), 2.22-2.42 (12H, m), 3.21-3.36 (5H, m), 3.42-3.51 (2H, m), 3.56-3.67 (2H, m), 3.77 (3H, s), 3.81-4.02 (2H, m), 4.23 (2H, t, J=7Hz), 4.86 (1H, m), 6.51-6.67 (2H, m), 6.79-6.93 (2H, m), 6.94-7.13 (3H, m), 7.44 (1H, m), 8.20 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

#### Example 27

The following compound was obtained according to a similar manner to that of Example 26.

4-[2-[(3-tert-Butoxycarbonylamino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.41 (9H, s), 1.46-1.95 (8H, m), 2.06-2.42 (16H, m), 2.56 (1H, m), 3.00 (1H, m), 3.22-3.38 (5H, m), 3.79 (3H, s), 3.83-4.03 (3H, m), 4.25 (2H, t, J=7Hz), 4.61 (1H, m), 4.87 (1H, m), 6.52-6.68 (2H, m), 6.79-6.95 (2H, m), 6.96-7.17 (3H, m), 7.46 (1H, m), 8.21 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

#### Example 28

- 183 -

To a solution of 4-[2-(3-tert-butoxycarbonylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-(5-carboxypent-1-yl)oxy-4-methylphenyl]benzamide (300 mg) and N-methylmorpholine (45 mg), in N,N-dimethylformamide (5 ml) was added isobutyl chloroformate (61 mg) at -15°C and the solution was stirred at the same temperature for 5 minutes. N,N,N'-Trimethylethylenediamine (54 mg) was added to the solution and the mixture was stirred at -15°C for 30 minutes, and then at ambient temperature for 1 hour. The mixture was diluted with ethyl acetate (20 ml) and the solution was washed successively with aqueous sodium hydrogen carbonate solution, water (15 ml x 3) and brine. The solution was dried over potassium carbonate and the solvent was removed under reduced pressure. The residue was purified on silica gel column chromatography (SiO<sub>2</sub> 40 g, 1-5% methanol in chloroform) to give 4-[2-(3-tert-butoxycarbonylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[(2-dimethylaminoeth-1-yl)-N-methylaminocarbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide (312 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.44-2.21 (8H, m), 2.25 (3H, s), 2.27 (6H, s), 2.29-2.50 (4H, m), 2.91 (1H, s), 3.00 (2H, s), 3.26-3.51 (4H, m), 3.31 (3H, s), 3.77 (3H, s), 3.81-4.02 (2H, m), 4.22 (2H, t, J=5Hz), 4.88 (1H, br), 6.52-6.68 (2H, m), 6.79-7.11 (5H, m), 7.43 (1H, m), 8.20 (1H, d, J=9Hz), 8.40 (1H, d, J=8Hz)

#### Example 29

The following compounds were obtained according to a similar manner to that of Example 28.

- 1) 4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[5-(3-dimethylaminoprop-1-yl)aminocarbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.42-1.57 (2H, m),

- 184 -

5 1.61-1.85 (6H, m), 2.04-2.35 (8H, m), 2.25 (3H, s),  
2.29 (9H, s), 2.46 (2H, t, J=6Hz), 3.20-3.38 (4H,  
m), 3.30 (3H, s), 3.76 (3H, s), 3.80-4.00 (2H, m),  
4.24 (2H, t, J=5Hz), 4.90 (1H, br), 6.61-6.72 (2H,  
m), 6.84-7.12 (6H, m), 7.43 (1H, d, J=8Hz), 8.20  
(1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

10 2) 4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-[5-(4-oxopiperidin-1-  
yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.50-1.92 (8H, m), 2.15  
(2H, t, J=6Hz), 2.29 (2H, t, J=5Hz), 2.38-2.51 (6H,  
m), 3.30 (2H, t, J=5Hz), 3.32 (3H, s), 3.70-4.05  
15 (6H, m), 3.80 (3H, s), 4.25 (2H, t, J=5Hz), 4.85  
(1H, br), 6.55-6.67 (2H, m), 6.83-7.15 (6H, m),  
7.40-7.51 (1H, m), 8.20 (1H, d, J=8Hz), 8.40 (1H,  
br)

20 3) 4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-  
amino-3-methoxy-N-methyl-N-[2-[5-(4-pyridylamino-  
carbonyl)pent-1-yl]oxy-4-methylphenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.50-1.61 (2H, m),  
1.75-1.93 (4H, m), 2.09-2.20 (2H, m), 2.30 (3H, s),  
2.42 (2H, br), 3.30 (1H, q, J=5Hz), 3.36 (3H, s),  
25 3.70 (3H, s), 3.72-4.00 (2H, m), 4.25 (2H, t,  
J=5Hz), 4.90 (1H, br), 6.60 (1H, br), 6.72 (1H, d,  
J=8Hz), 6.99-7.12 (6H, m), 7.43-7.51 (1H, m), 7.63  
(1H, d, J=8Hz), 8.19 (1H, d, J=8Hz), 8.42 (1H, d,  
J=7Hz), 8.46 (1H, br), 9.22 (1H, br)

### Example 30

35 To a solution of 4-[2-(3-tert-butoxycarbonylaminoprop-1-  
yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-(5-carboxypent-1-  
yl)oxy-4-methylphenyl]benzamide (250 mg) and N-  
methylmorpholine (37 mg) in dichloromethane (5 ml) was added

- 185 -

pivaloyl chloride (45 mg) at -15°C. After being stirred at the same temperature for 5 minutes, to the mixture was added 1-amino-4-methylpiperazine (47 mg) and the mixture was stirred at -15°C for 1 hour and then stirred at ambient temperature for additional 2 hours. The resulting mixture was poured into saturated aqueous sodium hydrogen carbonate solution (20 ml) and the solution was extracted with chloroform (15 ml x 3). The organic layer was washed with brine and dried over magnesium sulfate. The solvent was evaporated and the residue was purified on silica gel column chromatography (SiO<sub>2</sub> 30 g, 1-15% methanol in chloroform) to give 4-[2-(3-*tert*-butoxycarbonylaminoprop-1-yl)oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)aminocarbonylpent-1-yl]oxy-4-methylphenyl]benzamide (208 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.45-1.90 (6H, m), 2.10-2.19 (2H, m), 2.24 (3H, s), 2.25 (3H, s), 2.51 (2H, t, J=5Hz), 2.54-2.91 (8H, m), 3.30 (2H, t, J=5Hz), 3.34 (3H, s), 3.75 (3H, s), 3.80-4.03 (2H, m), 4.24 (2H, t, J=5Hz), 4.78-4.97 (1H, br), 6.53-6.67 (2H, m), 6.73-7.14 (6H, m), 7.40-7.50 (1H, m), 8.21 (1H, d, J=8Hz), 8.45 (1H, d, J=8Hz)

### Example 31

The following compounds were obtained according to a similar manner to that of Example 9.

- 1) 4-[2-(E)-[2-(4-Methylpiperazin-1-yl)carbonylethen-1-yl]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]-phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.59 (2H, m), 1.67-1.76 (2H, m), 1.79-1.87 (2H, m), 2.21 (3H, s), 2.26 (3H, s), 2.31 (3H, s), 2.31-2.44 (10H, m), 3.17-3.25 (2H, m), 3.34 (3H, s), 3.47-3.52 (2H, m), 3.56-3.67 (3H, m),

- 186 -

3.62 (3H, s), 3.82-3.99 (3H, m), 5.71 (1H, m),  
6.60-6.67 (2H, m), 6.86 (1H, d, J=7Hz), 6.92 (1H,  
d, J=7Hz), 6.98-7.03 (2H, m), 7.14 (1H, d, J=7Hz),  
7.43-7.62 (4H, m), 7.85 (1H, d, J=7Hz)

2) 4-[2-[(4-Methylpiperazin-1-yl)carbonylmethoxy]benzoyl]-  
amino-3-methoxy-N-[2-[5-(4-methylpiperazin-1-  
yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.30-1.90 (6H, m), 2.14 (3H, s), 2.26  
(3H, s), 2.35-2.46 (3H, m), 3.34 (3H, s), 3.46-3.55  
(4H, m), 3.59-3.68 (4H, m), 3.72 (3H, s), 3.80-4.01  
(2H, m), 4.90 (2H, s), 6.58-6.68 (2H, m), 6.82-7.06  
(4H, m), 7.13-7.20 (2H, m), 7.46-7.51 (1H, m), 8.19  
(1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)

### Example 32

A solution of 4-(2-iodobenzoyl)amino-N-[2-(4-methoxyphenyl)methoxy]phenyl-N-methylbenzamide (2.30 g) in a mixture of dichloromethane (30 ml) and trifluoroacetic acid (15 ml) was stirred at ambient temperature for 2 hours and the solvent was evaporated in vacuo. The residual oil was dissolved in chloroform (50 ml) and the solution was washed successively with water (50 ml), aqueous sodium hydrogen carbonate (50 ml) and brine (25 ml). The solution was dried over magnesium sulfate and the solvent was evaporated in vacuo to give 4-(2-iodobenzoyl)amino-N-(2-hydroxy)phenyl-N-methylbenzamide (1.20 g).

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 3.20 (3H, s), 6.69 (1H, t, J=7Hz),  
6.82 (1H, d, J=7Hz), 6.98-7.05 (3H, m), 7.40-7.54  
(4H, m), 7.90 (1H, d, J=7Hz), 9.84 (1H, s)

### Example 33

The following compounds were obtained according to a similar manner to that of Example 32.



- 187 -

- 1) 4-(2-Hydroxybenzoyl)amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]-phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 2.28 (3H, s), 2.32 (3H, s), 2.35-2.51 (4H, m), 3.36 (3H, s), 3.59-3.89 (2H, m), 5.02 (2H, s), 6.63-6.72 (2H, m), 6.88 (1H, t, J=7Hz), 7.00 (2H, d, J=8Hz), 7.20-7.46 (9H, m), 7.70 (1H, d, J=7Hz), 8.68 (1H, s)

- 2) 3-Methoxy-4-(2-hydroxybenzoyl)amino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]-phenylmethoxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 2.23 (3H, s), 2.30 (3H, s), 2.33-2.51 (4H, m), 3.37 (3H, s), 3.41-3.56 (2H, m), 3.68 (3H, s), 3.72-3.87 (2H, m), 4.91 (1H, d, J=14Hz), 5.09 (1H, d, J=14Hz), 6.63-6.71 (2H, m), 6.35-6.93 (2H, m), 7.00 (2H, d, J=8Hz), 7.33-7.50 (7H, m), 8.14 (1H, d, J=7Hz), 8.72 (1H, s)

- 3) 4[2-(3-Hydroxyprop-1-yl)thiobenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.44-1.58 (2H, m), 1.61-1.73 (2H, m), 1.77-1.89 (2H, m), 2.28 (3H, s), 2.31-2.40 (6H, m), 3.02 (2H, t, J=7.5Hz), 3.31 (3H, s), 3.42-3.50 (2H, m), 3.56-3.65 (2H, m), 3.67-3.78 (7H, m), 3.81-4.01 (2H, m), 6.58-6.67 (2H, m), 6.81-6.95 (2H, m), 7.03 (1H, s), 7.25 (1H, m), 7.36-7.50 (2H, m), 7.64 (1H, d, J=7Hz), 8.30 (1H, d, J=7Hz), 8.77 (1H, s)

#### Example 34

The following compound was obtained by using 2-nitro-4-(2-benzyloxybenzoyl)amino-N-methyl-N-[2-[5-(4-dimethylamino)piperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide as a starting compound according to a similar manner to that

- 188 -

of Example 10.

2-Amino-4-(2-hydroxybenzoyl)amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.21-2.02 (10H, m), 2.28-2.44 (12H, m), 2.48-2.69 (1H, m), 2.93-3.08 (1H, m), 3.30 (3H, s), 3.80-4.06 (4H, m), 4.68 (1H, br), 4.73 (2H, s), 5.32 (1H, s), 6.53-6.62 (3H, m), 6.78-6.96 (5H, m), 7.33-7.44 (1H, m), 7.78-7.88 (1H, m)

#### Example 35

A mixture of 4-(2-hydroxybenzoyl)amino-3-methoxy-N-(2-benzyloxy-4-methyl)phenyl-N-methylbenzamide (550 mg), 1-(tert-butoxycarbonyl)-4-hydroxypiperidine (223 mg), diethyl azodicarboxylate (193 mg) and triphenylphosphine (291 mg) in tetrahydrofuran (15 ml) was stirred at ambient temperature for 8 hours and the mixture was diluted with ethyl acetate (25 ml). The solution was washed with water and brine, and organic phase was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by silica gel column (30% ethyl acetate in n-hexane) to give 3-methoxy-4-[2-[1-(tert-butoxycarbonyl)piperidin-4-yl]oxybenzoyl]amino-N-(2-benzyloxy-4-methyl)phenyl-N-methylbenzamide (562 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.44 (9H, s), 1.72-1.90 (2H, m), 1.95-2.12 (2H, m), 2.27 (3H, s), 2.95-3.16 (4H, m), 3.37 (3H, s), 3.60 (3H, s), 3.73-4.00 (2H, m), 4.64 (1H, m), 4.88 (1H, d, J=14Hz), 5.08 (1H, d, J=14Hz), 6.65-6.71 (2H, m), 6.86 (1H, d, J=7Hz), 6.95-7.03 (3H, m), 7.09 (1H, t, J=7Hz), 7.25-7.50 (6H, m), 8.18 (1H, d, J=7Hz), 8.35 (1H, d, J=7Hz)

#### Example 36

The following compounds were obtained according to a similar manner to that of Example 35.

- 189 -

- 1) (S)-4-[2-[1-Methyl-3-(phthalimido)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methyl-piperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.43 (3H, d, J=7Hz), 1.47-1.92 (7H, m), 1.98-2.13 (1H, m), 2.20-2.47 (12H, m), 3.32 (3H, s), 3.42-3.53 (2H, m), 3.57-3.67 (2H, m), 3.73-4.05 (7H, m), 4.77 (1H, m), 6.51-6.69 (2H, m), 6.78-7.12 (5H, m), 7.42 (1H, m), 7.57 (4H, s), 8.08-8.24 (2H, m)

- 2) (R)-4-[2-[4-(Phthalimido-1-yl)but-2-yl]oxy]benzoyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methyl-piperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.44 and 1.47 (total 3H, s), 1.52-1.92 (8H, m), 2.02-2.12 (1H, m), 2.28 (3H, s), 2.30 (3H, s), 2.33-2.42 (6H, m), 3.35 (3H, s), 3.47-3.53 (2H, m), 3.60-3.67 (2H, m), 3.80 (3H, s), 3.85-4.00 (2H, br), 3.88 (2H, t, J=8Hz), 4.74-4.82 (1H, br), 6.57-6.69 (2H, m), 6.81-6.95 (2H, m), 6.98-7.09 (3H, m), 7.43 (1H, t, J=8Hz), 7.53-7.60 (4H, br), 8.14 (1H, d, J=8Hz), 8.20 (1H, d, J=8Hz)

ESI-MASS (m/z) : 804 (M+H)

- 3) (R)-4-[2-[4-(Phthalimido-1-yl)but-2-yl]oxy]benzoyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethyl-aminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.42 and 1.45 (total 3H, s), 1.50-1.90 (12H, m), 2.02-2.10 (1H, m), 2.28 (9H, s), 2.32-2.41 (4H, m), 2.52-2.62 (1H, m), 2.97-3.06 (1H, m), 3.35 (3H, s), 3.80 (3H, s), 3.87 (2H, t, J=8Hz), 3.90-3.97 (2H, m), 4.58-4.68 (1H, m), 4.72-4.81 (1H, m), 6.57-6.67 (2H, m), 6.81-6.93 (2H, m), 6.98-7.08 (3H, m), 7.43 (1H, t, J=8Hz), 7.53-7.59 (4H, br s), 8.13 (1H, d, J=8Hz), 8.20 (1H, d,

- 190 -

J=8Hz)

- 4) (S)-4-[2-[[4-(Phthalimido-1-yl)but-2-yl]oxy]benzoyl]-  
amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethyl-  
aminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]-  
benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.42 and 1.44 (total 3H, s), 1.50-  
1.91 (12H, m), 2.02-2.10 (1H, m), 2.29 (9H, s),  
2.32-2.41 (4H, m), 2.52-2.62 (1H, m), 2.95-3.05  
(1H, m), 3.36 (3H, s), 3.80 (3H, s), 3.86 (2H, t,  
J=8Hz), 3.90-3.97 (2H, m), 4.58-4.66 (1H, m), 4.72-  
4.80 (1H, m), 6.57-6.67 (2H, m), 6.81-6.92 (2H, m),  
6.98-7.08 (3H, m), 7.44 (1H, t, J=8Hz), 7.53-7.60  
(4H, br s), 8.13 (1H, d, J=8Hz), 8.21 (1H, d,  
J=8Hz)

ESI-MASS (m/z) : 832 (M+1)

- 5) 3-Methoxy-4-[2-[3-(phthalimido)-1-methylprop-1-  
yl]oxybenzoyl]amino-N-(2-benzyloxy-4-methyl)phenyl-N-  
methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.41 (3H, d, J=7.5Hz), 1.96-2.12 (2H,  
m), 2.24 (3H, s), 2.27-2.42 (2H, m), 3.39 (3H, s),  
3.60-3.69 (2H, m), 3.86 (2H, t, J=7.5Hz), 4.77 (1H,  
m), 4.94 (1H, d, J=14Hz), 5.08 (1H, d, J=14Hz),  
6.66-6.82 (3H, m), 6.95-7.08 (4H, m), 7.20-7.71  
(10H, m), 8.10-8.21 (2H, m)

### Example 37

The following compounds were obtained according to a  
similar manner to that of Example 14.

- 1) 4-[2-(3-Acetylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-  
N-(2-acetoxy-4-methylphenyl)-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.86 (3H, s), 2.10-2.19 (2H, m), 2.30  
(3H, s), 3.41 (2H, q, J=5Hz), 3.72 (3H, s), 4.21

- 191 -

(2H, t, J=5Hz), 5.94 (1H, br), 6.85 (1H, s), 6.90-7.11 (6H, m), 7.42-7.49 (1H, m), 8.10 (1H, d, J=8Hz), 8.42 (1H, d, J=8Hz)

- 5           2) 4-[2-(3-Acetylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide
- 10           NMR (CDCl<sub>3</sub>, δ) : 1.48-1.64 (2H, m), 1.58-1.85 (4H, m), 1.88 (3H, s), 2.12 (2H, t, J=5Hz), 2.29 (6H, s), 2.34-2.42 (2H, m), 2.57 (2H, t, J=5Hz), 3.30 (2H, q, J=5Hz), 3.32 (3H, s), 3.39 (2H, q, J=5Hz), 3.72-3.79 (2H, m), 3.76 (3H, s), 3.83-4.00 (2H, m), 4.20 (2H, t, J=5Hz), 6.33 (1H, br), 6.57-6.67 (2H, m), 6.83-7.10 (6H, m), 7.43 (1H, dd, J=2, 7Hz), 8.10 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)
- 15

Example 38

To an ice bath cooled solution of 4-[2-(3-aminoprop-1-yl)oxybenzoyl]amino-N-[2-(5-carboxypent-1-yl)oxy-4-methylphenyl]-N-methylbenzamide (650 mg) in dichloromethane (20 ml) were added triethylamine (137 mg) and di-tert-butylidicarbonate (296 mg) and the mixture was stirred at ambient temperature overnight. The solution was washed successively with water, 10% hydrochloric acid, saturated aqueous sodium hydrogen carbonate and brine, and the organic phase was dried over magnesium sulfate. The solvent was evaporated in vacuo to give 4-[2-[3-(tert-butoxycarbonyl)-aminoprop-1-yl]oxybenzoyl]amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methylphenyl]-N-methylbenzamide (749 mg).

20

25

30           NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7.5Hz), 1.40 (9H, s), 1.44-1.56 (2H, m), 1.66-1.76 (2H, m), 1.76-1.87 (2H, m), 2.06-2.15 (2H, m), 2.28 (3H, s), 2.34 (2H, t, J=7.5Hz), 3.31 (3H, s), 3.31-3.40 (2H, m), 3.85-3.97 (2H, m), 4.13 (2H, q, J=7.5Hz), 4.21 (2H, t, J=7.5Hz), 4.74 (1H, br), 6.54-6.62 (2H, m), 6.86

35

- 192 -

(1H, d, J=7Hz), 6.98 (1H, d, J=7Hz), 7.09 (1H, d, J=7Hz), 7.32 (2H, d, J=8Hz), 7.41-7.52 (3H, m), 8.11 (1H, d, J=7Hz), 9.87 (1H, s)

5     Example 39

The following compound was obtained according to a similar manner to that of Example 38.

10     3-Methoxy-4-[2-[3-(tert-butoycarbonyl)amino-1-methylprop-1-yl]oxybenzoyl]amino-N-(2-benzyloxy-4-methyl)-phenyl-N-methylbenzamide

15     NMR (CDCl<sub>3</sub>, δ) : 1.37 (9H, s), 1.41 (3H, d, J=7.5Hz), 1.84-2.11 (2H, m), 2.28 (3H, s), 3.20-3.31 (2H, m), 3.40 (3H, s), 3.64 (3H, s), 4.61 (1H, br), 4.72 (1H, m), 4.90 (1H, d, J=14Hz), 5.09 (1H, d, J=14Hz), 6.62-6.70 (2H, m), 6.84 (1H, d, J=7Hz), 6.93-7.12 (4H, m), 7.28-7.72 (6H, m), 8.22 (1H, d, J=7Hz), 8.38 (1H, d, J=7Hz)

20     Example 40

25     A solution of aqueous 4M sulfuric acid (0.5 ml) and 3-(phthalimid-1-yl)propanal (189 mg) in tetrahydrofuran (10 ml) was slowly added to a solution of 4-(2-aminobenzoylamino)-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenylbenzamide (560 mg) in tetrahydrofuran (10 ml) followed by the portionwise addition of sodium borohydride (59.8 mg) at 0°C. The mixture was diluted with 1,4-dioxane (5 ml) and stirred for an additional 1.5 hours at ambient temperature. The mixture was quenched with water (0.5 ml) and concentrated. The residue was 30 partitioned with ethyl acetate and saturated aqueous sodium hydrogen carbonate. The organic extract was washed with brine and dried over sodium sulfate, concentrated, and purified by silica gel column chromatography (SiO<sub>2</sub>, 30 g, 3% 35 methanol in chloroform) to give 3-methoxy-4-[2-[3-

- 193 -

(phthalimido)prop-1-yl]amino]benzoylamino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (200 mg).

5 NMR (CDCl<sub>3</sub>, δ) : 1.44-1.62 (2H, m), 1.63-1.93 (4H, m),  
1.97-2.12 (2H, m), 2.21-2.46 (12H, m), 3.17-3.38  
(5H, m), 3.42-3.56 (2H, m), 3.57-3.69 (2H, m),  
3.70-4.04 (7H, m), 6.51-6.73 (4H, m), 6.78-6.96  
(2H, m), 7.00 (1H, s), 7.20-7.35 (1H, m), 7.40 (1H,  
10 d, J=8Hz), 7.53-7.67 (3H, m), 7.72-7.86 (2H, m),  
8.13 (1H, d, J=8Hz), 8.34 (1H, s)

#### Example 41

A solution of 4-(2-nitrobenzoyl)amino-N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]-N-methylbenzamide  
15 (800 mg), 20% palladium hydroxide (200 mg) in ethanol (20 ml) was stirred under atmospheric pressure of hydrogen at ambient temperature. After 2 hours, the reaction mixture was filtered through a bed of Celite, and the solvent was removed by rotary evaporation and the crude product was purified by  
20 silica gel column chromatography (SiO<sub>2</sub> 30 g, ethyl acetate/hexane = 3/1) to give 4-(2-aminobenzoyl)amino-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]-benzamide (700 mg).

25 NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7Hz), 1.41-1.57 (2H, m), 1.63-1.87 (4H, m), 2.27 (3H, s), 2.33 (2H, t, J=7Hz), 3.32 (3H, s), 3.78-4.00 (2H, m), 4.12 (2H, q, J=7Hz), 5.38-5.56 (2H, m), 6.55-6.64 (2H, m), 6.64-6.76 (2H, m), 6.87 (1H, d, J=9Hz), 7.22 (1H, d, J=9Hz), 7.28-7.50 (5H, m), 7.79 (1H, br s)

#### Example 42

The following compound was obtained according to a similar manner to that of Preparation 4.

35 4-(2-Aminobenzenesulfonyl)amino-3-methoxy-N-methyl-N-[4-

- 194 -

methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.45-1.54 (2H, m), 1.65-1.82 (4H, m),  
2.30 (3H, s), 2.33 (3H, s), 2.35-2.43 (6H, m), 3.29  
(3H, s), 3.46-3.51 (5H, m), 3.60-3.65 (4H, m),  
4.84-4.89 (2H, m), 6.56-6.89 (6H, m), 7.28-7.48  
(4H, m)

ESI-MASS (m/z) : 638 (M+H)

#### Example 43

A solution of 4-[2-(acetyloxy)benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide (400 mg) in methanol (10 ml) was treated with 1N sodium hydroxide solution (3 ml) at ambient temperature. After 6 hours, the reaction mixture was concentrated in vacuo and extracted with the mixture of dichloromethane and diluted hydrochloric acid. The organic phase was washed with brine and dried over sodium sulfate. The crude product was purified by silica gel column chromatography (SiO<sub>2</sub> 30 g, 5% methanol in chloroform) to give 4-[2-(hydroxy)benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide (290 mg).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.27-2.00 (10H, m), 2.21-2.46 (12H, m), 2.56 (1H, m), 3.00 (1H, m), 3.33 (3H, s), 3.80 (3H, s), 3.82-4.05 (4H, m), 4.63 (1H, m), 6.55-6.68 (2H, m), 6.82-7.09 (5H, m), 7.42 (1H, m), 7.55 (1H, m), 8.20 (1H, m).

#### Example 44

The following compounds were obtained according to a similar manner to that of Example 43.

- 1) 4-(2-Hydroxybenzoyl)amino-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yloxy)-4-methylphenyl]benzamide



- 195 -

- 5 NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7Hz), 1.42-1.58 (2H, m), 1.61-1.90 (4H, m), 2.28 (3H, s), 2.33 (2H, t, J=7Hz), 3.32 (3H, s), 3.80 (3H, s), 3.81-4.02 (2H, m), 4.12 (2H, q, J=7Hz), 6.53-6.67 (2H, m), 6.80-6.99 (3H, m), 7.01 (1H, d, J=8Hz), 7.07 (1H, s), 7.42 (1H, dd, J=8, 8Hz), 7.49 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz), 8.72 (1H, s)
- 10 2) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-(2-methylphenyl)benzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.21 (3H, s), 3.40 (3H, s), 3.78 (3H, s), 6.82-7.23 (9H, m), 7.37-7.53 (2H, m), 8.18 (1H, d, J=8Hz), 8.69 (1H, br s)
- 15 3) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.42-1.59 (2H, m), 1.60-1.89 (4H, m), 2.20-2.46 (12H, m), 3.32 (3H, s), 3.42-3.53 (2H, m), 3.57-3.69 (2H, m), 3.71-4.02 (6H, m), 6.51-6.68 (2H, m), 6.79-7.08 (5H, m), 7.40 (1H, m), 7.51 (1H, d, J=8Hz), 8.18 (1H, d, J=8Hz), 8.86 (1H, br s)
- 20 4) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-(2-benzyloxy-4-methylphenyl)-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.30 (3H, s), 3.38 (3H, s), 3.63 (3H, s), 4.89 (1H, d, J=13Hz), 5.08 (1H, d, J=13Hz), 6.62-6.68 (2H, m), 6.82-7.00 (6H, m), 7.28-7.42 (5H, m), 7.47 (1H, d, J=8Hz), 8.13 (1H, d, J=8Hz), 8.79 (1H, s)
- 25 5) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-[2-[4-(2-oxazolin-2-yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 2.28 (3H, s), 3.40 (3H, s), 3.67 (3H, s), 4.06 (2H, t, J=10Hz), 4.41 (2H, t, J=10Hz),
- 30
- 35

- 196 -

4.92 (1H, d, J=12Hz), 5.10 (1H, d, J=12Hz), 6.60 (1H, s), 6.71 (1H, d, J=8Hz), 6.87-7.08 (5H, m), 7.28 (1H, d, J=8Hz), 7.42 (1H, dd, J=2, 8Hz), 7.52 (1H, d, J=8Hz), 8.16 (1H, d, J=8Hz), 8.82 (1H, s)

6) 4-(2-Hydroxybenzoyl)amino-3-methyl-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.48 (2H, br), 1.60-1.81 (4H, m), 2.19 (3H, s), 2.28 (3H, s), 2.30-2.35 (3H, m), 2.38 (3H, s), 2.50 (4H, br), 3.30 (3H, s), 3.52 (2H, br), 3.69 (2H, br), 3.83 (1H, br), 3.92 (1H, br), 6.62 (2H, s), 6.89-6.93 (2H, m), 7.02-7.10 (2H, m), 7.35 (1H, s), 7.40-7.47 (1H, m), 7.63-7.70 (2H, m), 8.52 (1H, br)

#### Example 45

A solution of 4-[2-[3-(tert-butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (542 mg) in 90% trifluoroacetic acid (10 ml) was stirred at ambient temperature for 3 hours and the solvent was evaporated in vacuo. The residue was stirred with chloroform (20 ml) and saturated aqueous sodium hydrogen carbonate (10 ml) and the organic phase was separated. The solution was washed with brine and dried over magnesium sulfate. The solvent was evaporated in vacuo to give 4-[2-(3-aminoprop-1-yl)oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (465 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.47-1.59 (2H, m), 1.67-2.00 (6H, m), 2.06-2.66 (2H, m), 2.35 (3H, s), 2.39 (3H, s), 2.32-2.41 (4H, m), 2.96 (2H, t, J=7.5Hz), 3.31 (3H, s), 3.45-3.50 (2H, m), 3.58-3.65 (2H, m), 3.89-3.99 (2H, m), 4.29 (2H, d, J=7.5Hz), 6.54-6.62 (2H, m),

- 197 -

6.85 (1H, d, J=7Hz), 7.01 (1H, d, J=7Hz), 7.10 (1H, t, J=7Hz), 7.32 (2H, d, J=8Hz), 7.43-7.50 (3H, m), 8.20 (1H, d, J=7Hz)

5      Example 46

The following compounds were obtained according to a similar manner to that of Example 45.

- 10      1) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

15      NMR (CDCl<sub>3</sub>, δ) : 1.40-1.92 (6H, m), 1.98-2.12 (2H, m), 2.19-2.44 (12H, m), 2.90 (2H, t, J=7Hz), 3.32 (3H, s), 3.40-3.53 (2H, m), 3.56-3.68 (2H, m), 3.78 (3H, s), 3.80-4.02 (2H, m), 4.28 (2H, t, J=7Hz), 6.51-6.67 (2H, m), 6.78-6.95 (2H, m), 6.97-7.16 (3H, m), 7.44 (1H, m), 8.21 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

- 20      2) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

25      NMR (CDCl<sub>3</sub>, δ) : 1.26-1.92 (12H, m), 1.98-2.12 (2H, m), 2.27 (9H, s), 2.29-2.42 (3H, m), 2.56 (1H, m), 2.89 (2H, t, J=7Hz), 3.00 (1H, m), 3.32 (3H, s), 3.78 (3H, s), 3.82-4.02 (3H, m), 4.27 (2H, t, J=7Hz), 4.61 (1H, m), 6.52-6.67 (2H, m), 6.79-6.96 (2H, m), 6.97-7.12 (3H, m), 7.43 (1H, m), 8.21 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

- 30      3) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide

35      NMR (CDCl<sub>3</sub>, δ) : 0.95 (3H, d, J=7.5Hz), 1.00-1.14 (2H, m), 1.46-1.90 (8H, m), 2.01-2.12 (2H, m), 2.26 (3H, m)

- 198 -

5 s), 2.34 (2H, t, J=7.5Hz), 2.52 (1H, m), 2.85-3.03 (3H, m), 3.31 (3H, s), 3.79 (3H, s), 3.79-4.00 (4H, m), 4.32 (2H, t, J=7.5Hz), 4.55 (1H, m), 6.58 (1H, d, J=7Hz), 6.62 (1H, s), 6.84 (1H, d, J=7Hz), 6.90 (1H, d, J=7Hz), 7.00-7.11 (3H, m), 7.42 (1H, t, J=7Hz), 8.21 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

10 4) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[5-[(2S)-carbamoylpyrrolidin-1-yl]carbonylpent-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.48-2.20 (12H, m), 2.28 (3H, s), 2.32-2.40 (2H, m), 2.88-3.00 (2H, m), 3.31 (3H, s), 3.33-3.61 (2H, m), 3.80 (3H, s), 3.82-3.99 (2H, m), 4.29 (2H, t, J=7Hz), 4.54 (1H, m), 6.52-6.63 (2H, m), 6.81-7.10 (5H, m), 7.43 (1H, t, J=7Hz), 8.14 (1H, d, J=7Hz), 8.38 (1H, d, J=7Hz)

20 5) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(4-aminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.63-1.94 (4H, m), 1.99-2.18 (2H, m), 2.23 (3H, s), 2.62-3.07 (2H, m), 3.29 (3H, s), 3.29-3.51 (2H, m), 3.75-4.00 (2H, m), 3.76 (3H, s), 4.21 (2H, t, J=7.5Hz), 6.56-6.85 (4H, m), 7.28-7.62 (2H, m), 8.13 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

30 6) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(4-acetylaminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

35 NMR (CDCl<sub>3</sub>, δ) : 1.60-1.86 (4H, m), 2.00 (3H, s), 2.08-2.20 (2H, m), 2.27 (3H, s), 2.93-3.03 (2H, m), 3.30 (3H, s), 3.30-3.50 (2H, m), 3.77 (3H, s), 3.83-3.98 (2H, m), 4.26 (2H, t, J=7.5Hz), 6.53-6.65 (2H, m), 6.86-7.12 (5H, m), 7.42 (1H, t, J=7Hz), 8.12 (1H, d, J=7Hz), 8.37 (1H, d, J=7Hz)

- 199 -

- 7) 3-Methoxy-4-[2-(piperidin-4-yl)oxybenzoyl]amino-N-(2-hydroxy-4-methyl)phenyl-N-methylbenzamide

NMR (DMSO-d<sub>6</sub>, δ) : 1.50-1.62 (2H, m), 1.94-2.05 (2H, m), 2.14 (3H, s), 2.57 (2H, t, J=7.5Hz), 2.91-3.00 (2H, m), 3.16 (3H, s), 3.75 (3H, s), 4.73 (1H, m), 6.48 (1H, d, J=7Hz), 6.64 (1H, s), 7.87 (1H, d, J=7Hz), 7.92 (1H, d, J=7Hz), 7.01 (1H, s), 7.09 (1H, t, J=7Hz), 7.32 (1H, d, J=7Hz), 7.52 (1H, t, J=7Hz), 8.02 (1H, d, J=7Hz), 8.27 (1H, d, J=7Hz)

- 8) 3-Methoxy-4-[2-(piperidin-4-yl)oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.46-1.88 (8H, m), 2.07-2.19 (2H, m), 2.26 (3H, s), 2.29 (3H, s), 2.32-2.41 (6H, m), 2.72 (2H, t, J=7.5Hz), 3.10-3.20 (2H, m), 3.32 (3H, s), 3.45-3.50 (2H, m), 3.60-3.66 (2H, m), 3.80 (3H, s), 3.83-4.00 (2H, m), 4.57 (1H, m), 6.58 (1H, d, J=7Hz), 6.62 (1H, s), 6.82-6.91 (2H, m), 6.98-7.11 (3H, m), 7.43 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 9) 4-[2-(3-Amino-1-methylprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.42 (3H, d, J=7.5Hz), 1.46-1.89 (6H, m), 1.99-2.11 (2H, m), 2.28 (3H, s), 2.30 (3H, s), 2.31-2.42 (6H, m), 2.85 (2H, t, J=7.5Hz), 3.33 (3H, s), 3.45-3.50 (2H, m), 3.59-3.66 (2H, m), 3.80 (3H, s), 3.84-4.01 (2H, m), 4.80 (1H, m), 6.59 (1H, d, J=7Hz), 6.63 (1H, s), 6.82-6.92 (2H, m), 7.01-7.10 (3H, m), 7.44 (1H, t, J=7Hz), 8.22 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 10) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-

- 200 -

methyl-N-[2-(5-aminocarbonylpent-1-yl)oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40-1.59 (2H, m), 1.61-1.90 (4H, m),  
2.11-2.30 (4H, m), 2.35 (3H, s), 3.00 (2H, t,  
J=6Hz), 3.11 (2H, br), 3.29 (3H, s), 3.75 (3H, s),  
3.76-4.02 (2H, m), 4.23 (2H, t, J=5Hz), 6.00 (1H,  
br), 6.50 (1H, br), 6.55-6.71 (2H, m), 6.87-7.12  
(5H, m), 7.42 (1H, dd, J=2, 7Hz), 8.10 (1H, d,  
J=9Hz), 8.36 (1H, d, J=8Hz)

11) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(morpholin-4-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.90 (6H, m), 2.11 (2H, t,  
J=5Hz), 2.26 (3H, s), 2.21-2.52 (6H, m), 2.79-2.90  
(3H, m), 2.96 (2H, t, J=5Hz), 3.31 (3H, s), 3.40-  
3.49 (2H, m), 3.52-3.62 (2H, m), 3.80 (3H, s),  
3.83-4.04 (2H, m), 4.29 (2H, t, J=5Hz), 6.57-6.68  
(2H, m), 6.81-7.12 (6H, m), 7.41-7.50 (1H, m), 8.17  
(1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)

12) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-oxopiperidin-1-yl)carbonylpent-1-yl]oxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.45-2.05 (8H, m), 2.11 (2H, t,  
J=5Hz), 2.28 (3H, s), 2.41-2.52 (2H, m), 2.96 (2H,  
t, J=5Hz), 3.31 (3H, s), 3.70-4.61 (8H, m), 6.52-  
7.55 (8H, m), 8.02-8.46 (3H, m)

13) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-(2-methoxy-4-methylphenyl)-N-methylbenzamide

NMR (DMSO-d<sub>6</sub>, δ) : 1.90-1.98 (2H, m), 2.25 (3H, s),  
2.71 (2H, t, J=6Hz), 3.19 (3H, s), 3.73 (3H, s),  
4.32 (2H, t, J=5Hz), 6.67 (1H, d, J=8Hz), 6.80-6.96  
(2H, m), 7.26 (1H, d, J=8Hz), 7.55 (1H, dd, J=2, 8Hz),

- 201 -

8.03 (1H, d, J=8Hz), 8.29 (1H, d, J=8Hz)

- 14) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
[4-(thiazol-2-yl)phenylmethyl]oxy-4-methylphenyl]-N-  
methylbenzamide

5  
10  
NMR (CDCl<sub>3</sub>, δ) : 2.02-2.10 (2H, m), 2.29 (3H, s), 2.89  
(2H, t, J=5Hz), 3.40 (3H, s), 3.64 (3H, s), 4.25  
(2H, t, J=5Hz), 4.90 (1H, d, J=11Hz), 5.09 (1H, d,  
J=11Hz), 6.62-6.71 (2H, m), 6.88 (1H, d, J=8Hz),  
6.98-7.10 (5H, m), 7.24-7.48 (4H, m), 7.81 (1H, d,  
J=3Hz), 7.95 (1H, d, J=8Hz), 8.20 (1H, d, J=8Hz),  
8.37 (1H, d, J=8Hz)

- 15) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
[4-(oxazol-2-yl)phenylmethyl]oxy-4-methylphenyl]-N-  
methylbenzamide

15  
20  
NMR (CDCl<sub>3</sub>, δ) : 2.00-2.11 (2H, m), 2.29 (3H, s), 2.89  
(2H, t, J=5Hz), 3.40 (3H, s), 3.66 (3H, s), 4.91  
(1H, d, J=12Hz), 5.10 (1H, d, J=12Hz), 6.64 (1H,  
s), 6.70 (1H, d, J=8Hz), 6.87 (1H, d, J=8Hz), 7.00-  
7.12 (4H, m), 7.21 (1H, s), 7.25-7.49 (4H, m), 7.65  
(1H, s), 8.04 (1H, d, J=8Hz), 8.23 (1H, d, J=8Hz),  
8.37 (1H, d, J=8Hz)

- 25) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
[4-(2-oxazolin-2-yl)phenylmethyl]oxymethylphenyl]-N-  
methylbenzamide

30  
35  
NMR (CDCl<sub>3</sub>, δ) : 2.02-2.11 (2H, m), 2.28 (3H, s), 2.90  
(2H, t, J=5Hz), 3.39 (3H, s), 3.67 (3H, s), 4.05  
(2H, t, J=9Hz), 4.29 (2H, t, J=5Hz), 4.41 (2H, t,  
J=5Hz), 4.89 (1H, d, J=12Hz), 5.09 (1H, d, J=12Hz),  
6.63 (1H, s), 6.70 (1H, d, J=8Hz), 6.84 (1H, d,  
J=8Hz), 7.00-7.12 (4H, m), 7.37 (2H, d, J=8Hz),  
7.41 (1H, d, J=8Hz), 7.93 (2H, d, J=5Hz), 8.20 (1H,  
d, J=8Hz), 8.36 (1H, d, J=8Hz)

- 202 -

- 17) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-(pyrimidin-2-yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide

5 NMR (CDCl<sub>3</sub>, δ) : 2.05-2.14 (2H, m), 2.27 (3H, s), 2.89 (2H, t, J=5Hz), 3.38 (3H, s), 3.64 (3H, s), 4.24 (2H, t, J=5Hz), 4.94 (1H, d, J=13Hz), 5.12 (1H, d, J=13Hz), 6.65-6.72 (2H, m), 6.85 (1H, d, J=8Hz), 6.97-7.18 (5H, m), 7.39-7.46 (3H, m), 8.13 (1H, d, J=8Hz), 8.35 (1H, d, J=8Hz), 8.41 (2H, d, J=8Hz), 10 8.24 (2H, d, J=3Hz)

- 18) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(4-cyanophenylmethyl)oxy-4-methylphenyl]-N-methylbenzamide

15 NMR (CDCl<sub>3</sub>, δ) : 2.09-2.20 (2H, m), 2.28 (3H, s), 2.97 (2H, t, J=5Hz), 3.35 (3H, s), 3.65 (3H, s), 4.24 (2H, br), 4.88 (1H, d, J=12Hz), 5.06 (1H, d, J=12Hz), 6.57 (1H, s), 6.67-6.80 (2H, m), 6.95-7.08 (5H, m), 7.35-7.45 (3H, m), 7.62 (2H, d, J=8Hz), 20 8.11 (1H, d, J=8Hz), 8.30 (1H, d, J=8Hz)

- 19) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(2-dimethylaminoeth-1-yl)oxycarbonylpent-1-yl]oxy-4-methylphenyl]benzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.60 (2H, m), 1.67-1.88 (4H, m), 2.05-2.14 (2H, m), 2.27 (9H, s), 2.38 (2H, t, J=6Hz), 2.58 (2H, t, J=5Hz), 2.92 (2H, t, J=5Hz), 3.33 (3H, s), 3.80 (3H, s), 3.86-4.00 (2H, m), 4.19 (2H, t, J=5Hz), 4.30 (2H, t, J=5Hz), 6.57-6.67 (2H, 30 m), 6.87 (1H, dd, J=2, 8Hz), 7.00-7.11 (4H, m), 7.44 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.38 (1H, d, J=9Hz)

- 20) 4-[2-(3-Aminoprop-1-yloxy)benzoyl]amino-3-methoxy-N-(2-hydroxy-4-methylphenyl)-N-methylbenzamide

35



- 203 -

5 NMR (DMSO-d<sub>6</sub>, δ) : 1.92-2.03 (2H, m), 2.16 (3H, s),  
2.75 (2H, t, J=5Hz), 3.20 (3H, s), 3.75 (3H, s), 4.34  
(2H, t, J=5Hz), 6.49 (1H, d, J=8Hz), 6.66 (1H, s),  
6.87 (1H, d, J=8Hz), 6.92 (1H, d, J=8Hz), 7.12 (1H,  
dd, J=7, 8Hz), 7.29 (1H, d, J=8Hz), 7.58 (1H, dd, J=2,  
8Hz), 8.05 (1H, d, J=8Hz), 8.27 (1H, d, J=8Hz)

10 21) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
[4-(1,5-dimethyl-3-cyanopyrrol-2-yl)phenylmethyl]oxy-4-  
methylphenyl]-N-methylbenzamide

15 NMR (CDCl<sub>3</sub>, δ) : 2.00-2.11 (2H, m), 2.14 (3H, s), 2.21  
(3H, s), 2.89 (2H, t, J=5Hz), 3.40 (3H, s), 3.45  
(3H, s), 3.62 (3H, s), 4.27 (2H, t, J=5Hz), 4.89  
(1H, d, J=13Hz), 5.13 (1H, d, J=13Hz), 6.22 (1H,  
s), 6.68-6.75 (2H, m), 6.89 (1H, d, J=8Hz), 7.00-  
7.12 (5H, m), 7.38-7.47 (6H, m), 8.19 (1H, d,  
J=8Hz), 8.38 (1H, d, J=8Hz)

20 22) 4-[2-(3-Aminoprop-1-yloxy)benzoyl]amino-3-methoxy-N-[2-  
[4-(N,N-dimethylureido)but-1-yl]oxy-4-methylphenyl]-N-  
methylbenzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.62-1.88 (4H, m), 1.90-2.15 (2H, m),  
2.27 (3H, s), 2.86-2.94 (2H, m), 2.90 (6H, s),  
3.22-3.35 (2H, m), 3.31 (3H, s), 3.77 (3H, s),  
3.75-3.98 (2H, m), 4.27 (2H, t, J=5Hz), 6.57-6.70  
(2H, m), 6.88-7.11 (6H, m), 7.42 (1H, dd, J=2,  
8Hz), 8.19 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

30 23) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
[3-(4-methylpiperazin-1-yl)carbonylpyrid-6-yl]methoxy-4-  
methylphenyl]-N-methylbenzamide

35 NMR (CDCl<sub>3</sub>, δ) : 2.09-2.20 (2H, m), 2.28 (3H, s), 2.31  
(3H, s), 2.34-2.52 (4H, m), 2.96 (2H, t, J=5Hz),  
3.40 (3H, s), 3.42-3.50 (2H, m), 3.69 (3H, s),  
3.70-3.84 (2H, m), 4.29 (2H, t, J=5Hz), 4.98 (1H,

- 204 -

5 d, J=13Hz), 5.18 (1H, d, J=13Hz), 6.62 (1H, s),  
6.72 (1H, d, J=8Hz), 6.98-7.11 (5H, m), 7.26-7.34  
(1H, m), 7.45 (1H, dd, J=2, 8Hz), 7.73 (1H, d,  
J=8Hz), 8.16 (1H, d, J=8Hz), 8.36 (1H, d, J=8Hz),  
8.63 (1H, s)

24) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
[4-(3-dimethylaminoprop-1-yloxy-carbonyl)aminobut-1-  
yl]oxy-4-methylphenyl]-N-methylbenzamide

10 NMR (CDCl<sub>3</sub>, δ) : 1.62-1.87 (6H, m), 2.02-2.11 (2H, m),  
2.27 (6H, s), 2.41 (2H, t, J=5Hz), 2.91 (2H, t,  
J=5Hz), 3.22 (2H, q, J=5Hz), 3.30 (3H, s), 3.78  
(3H, s), 3.84-3.95 (2H, m), 4.08 (2H, t, J=5Hz),  
15 4.27 (2H, t, J=5Hz), 6.60-6.66 (2H, m), 6.90 (1H,  
d, J=8Hz), 6.99-7.10 (3H, m), 7.44 (1H, dd, J=2,  
8Hz), 8.18 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

25) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-(4-methylhomopiperazin-1-yl)carbonyl]pent-  
1-yl]oxy-4-methylphenyl]benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.50-2.18 (8H, m), 2.30 (3H, s), 2.32  
(2H, t, J=5Hz), 2.33 (3H, s), 2.53-2.70 (4H, m),  
2.93 (2H, t, J=5Hz), 3.35 (3H, s), 3.52-3.72 (4H,  
m), 3.80 (3H, s), 3.82-4.09 (2H, m), 4.31 (2H, t,  
25 J=5Hz), 6.55-6.70 (2H, m), 6.82-7.18 (6H, m), 7.42-  
7.53 (1H, m), 8.20 (1H, d, J=8Hz), 8.41 (1H, d,  
J=8Hz)

26) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-(2-dimethylaminoethyl)aminocarbonyl]pent-  
1-yl]oxy-4-methylphenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.60 (2H, m), 1.66-2.15 (8H, m),  
2.22 (6H, s), 2.26 (3H, s), 2.41 (2H, t, J=5Hz),  
3.22-3.39 (2H, m), 3.31 (3H, s), 3.70-4.00 (2H, m),  
35 3.78 (3H, s), 4.28 (2H, t, J=5Hz), 6.37 (1H, br),

- 205 -

6.59 (2H, br), 6.81-7.13 (6H, m), 7.42 (1H, dd, J=2, 8Hz), 8.18 (1H, d, J=8Hz), 8.36 (1H, d, J=8Hz)

27) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[N-(2-dimethylaminoethyl)-N-methylamino-carbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.44-2.21 (8H, m), 2.25 (3H, s), 2.27 (6H, s), 2.29-2.50 (4H, m), 2.91 (1H, s), 3.00 (2H, s), 3.26-3.51 (4H, m), 3.31 (3H, s), 3.77 (3H, br s), 3.81-4.02 (2H, m), 4.22 (2H, t, J=5Hz), 4.88 (1H, br), 6.52-6.68 (2H, br), 6.79-7.11 (5H, m), 7.43-7.50 (1H, m), 8.20 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)

28) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[N-(3-dimethylaminoprop-1-yl)carbamoyl]pent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.46-1.60 (2H, m), 1.63-1.99 (8H, m), 2.03-2.14 (2H, m), 2.21 (2H, t, J=5Hz), 2.24 (6H, s), 2.29 (3H, s), 2.39 (2H, t, J=5Hz), 2.90 (2H, t, J=6Hz), 3.25-3.37 (2H, m), 3.32 (3H, s), 3.79 (3H, s), 3.81-4.01 (2H, m), 4.30 (2H, t, J=5Hz), 6.61 (2H, br), 6.85-7.14 (6H, m), 7.39-7.50 (1H, m), 8.20 (1H, d, J=8Hz), 8.40 (1H, br)

29) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[N-(3-dimethylaminoprop-1-yl)-N-methylcarbamoyl]pent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.52-1.94 (6H, m), 2.05-2.14 (2H, m), 2.20 (3H, s), 2.21 (3H, s), 2.26 (3H, s), 2.20-2.45 (6H, s), 2.90 (2H, t, J=5Hz), 2.91 and 2.99 (total 3H, s, rotamer), 3.32 (3H, s), 3.40 (2H, t, J=5Hz), 3.80 (3H, s), 4.31 (2H, t, J=5Hz), 6.55-6.67 (2H, m), 7.41-7.49 (2H, m), 8.21 (1H, d, J=8Hz), 8.42 (1H, d, J=8Hz)

- 206 -

- 30) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-hydroxypiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.43-1.95 (6H, m), 2.03-2.51 (8H, m),  
2.29 (3H, s), 2.94 (2H, t, J=5Hz), 2.98-3.22 (4H,  
m), 3.32 (3H, s), 3.46-3.58 (1H, m), 3.79 (3H, s),  
3.80-4.26 (6H, m), 4.28 (2H, t, J=5Hz), 6.56-6.67  
10 (2H, m), 6.81-7.13 (6H, m), 7.36 (1H, dd, J=8,  
8Hz), 8.10-8.20 (1H, m), 8.33-8.49 (1H, m)

- 31) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-aminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.51-2.03 (6H, m), 2.09-2.19 (2H, m),  
2.27 (3H, s), 2.29-2.42 (4H, m), 2.59-2.71 (2H, m),  
2.94 (2H, t, J=5Hz), 2.96-3.11 (3H, m), 3.33 (3H,  
s), 3.78 (3H, s), 3.85-4.02 (2H, m), 4.22 (2H, t,  
J=5Hz), 6.55-6.67 (2H, m), 6.81-7.12 (6H, m), 7.44  
20 (1H, dd, J=8, 8Hz), 8.19 (1H, d, J=8Hz), 8.40 (1H,  
d, J=8Hz)

- 32) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)-aminocarbonylpent-1-yl]oxy-4-methylphenyl]benzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.46-1.89 (6H, m), 1.93-2.05 (4H, m),  
2.25 (6H, s), 2.49 (2H, t, J=5Hz), 2.52-2.62 (2H,  
m), 2.79-2.89 (2H, m), 2.92 (2H, t, J=5Hz), 3.31  
(3H, s), 3.79 (3H, s), 3.80-4.01 (2H, m), 4.28 (2H,  
t, J=5Hz), 6.56-6.64 (2H, m), 6.80-7.12 (6H, m),  
30 7.41-7.50 (1H, m), 8.18 (1H, d, J=8Hz), 8.40 (1H,  
d, J=8Hz)

- 33) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[bis(2-hydroxyethyl)-aminocarbonylpent-1-yl]oxy-4-methylphenyl]benzamide

35

- 207 -

5 NMR (CDCl<sub>3</sub>, δ) : 1.54-1.91 (6H, m), 2.11-2.20 (2H, m),  
2.26 (3H, s), 2.38-2.59 (4H, m), 3.40-3.57 (4H, m),  
3.61-3.97 (6H, m), 4.22 (2H, t, J=5Hz), 6.60-6.68  
(2H, m), 6.88-7.16 (6H, m), 7.44-7.54 (1H, m), 8.12  
(1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

34) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(2,2-dimethylhydrazino)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

10 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.91 (6H, m), 2.06-2.40 (4H, m),  
2.28 (3H, s), 2.51 (3H, s), 2.57 (3H, s), 2.92 (2H,  
t, J=5Hz), 3.32 (3H, s), 3.78 (3H, s), 3.80-4.02  
(2H, m), 4.28 (2H, t, J=5Hz), 6.55-6.68 (2H, m),  
6.80-7.13 (5H, m), 7.46 (1H, dd, J=8Hz), 8.19 (1H,  
15 d, J=8Hz), 8.38 (1H, br)

35) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(carbamoylmethylamino)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.58 (2H, m), 1.68-1.85 (4H, m),  
2.06-2.17 (2H, m), 2.27 (3H, s), 2.94 (2H, t,  
J=5Hz), 3.31 (3H, s), 3.80 (3H, s), 3.81-4.00 (2H,  
m), 3.89 (2H, d, J=5Hz), 4.28 (2H, t, J=5Hz), 5.78  
(1H, br), 6.60-6.74 (3H, m), 6.90-7.13 (6H, m),  
25 7.41-7.49 (1H, m), 8.17 (1H, d, J=8Hz), 8.39 (1H,  
d, J=8Hz)

36) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(2-carbamoylethylamino)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.58 (2H, m), 1.62-1.84 (4H, m),  
2.14 (2H, t, J=5Hz), 2.22 (2H, t, J=5Hz), 2.29 (3H,  
s), 2.40 (2H, t, J=5Hz), 2.98 (2H, br), 3.30 (3H,  
s), 3.40-3.55 (2H, m), 3.78 (3H, s), 3.80-4.01 (2H,  
35 m), 4.27 (2H, t, J=5Hz), 6.58-6.79 (4H, m), 6.88-

- 208 -

7.12 (6H, m), 7.41-7.49 (1H, m), 8.16 (1H, d, J=8Hz), 8.39 (1H, d, J=7Hz)

- 5 37) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-pyridylaminocarbonyl)pent-1-yl]oxy-4-methylphenyl]benzamide

10 NMR (CDCl<sub>3</sub>, δ) : 1.52-1.89 (6H, m), 2.10-2.22 (2H, m), 2.26 (3H, s), 2.45 (2H, br), 2.95 (2H, t, J=5Hz), 3.32 (3H, s), 3.72 (3H, s), 3.82-4.00 (2H, m), 4.27 (2H, t, J=5Hz), 6.57-6.72 (2H, m), 6.90-7.15 (6H, m), 7.46 (1H, dd, J=2, 8Hz), 7.56 (2H, br), 8.12 (1H, d, J=8Hz), 8.35-8.50 (3H, m), 9.46 (1H, br)

- 15 38) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[4-(diethylaminopiperidin-1-yl)carbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.05 (6H, t, J=5Hz), 1.35-1.95 (10H, m), 2.04-2.13 (2H, m), 2.28 (3H, s), 2.36 (2H, t, J=5Hz), 2.54 (4H, q, J=5Hz), 2.56-2.80 (2H, m), 2.91 (2H, t, J=5Hz), 2.93-3.07 (2H, m), 3.33 (3H, s), 3.80 (3H, s), 3.82-4.03 (2H, m), 4.30 (2H, t, J=5Hz), 6.56-6.68 (2H, m), 6.81-7.12 (6H, m), 7.42-7.49 (1H, m), 8.22 (1H, d, J=7Hz), 8.41 (1H, d, J=8Hz)

- 25 39) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[6-(4-methylpiperazin-1-yl)hex-1-yl]oxy-4-methylphenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.58 (2H, m), 1.62-1.84 (4H, m), 2.14 (2H, t, J=5Hz), 2.29 (3H, s), 2.40 (2H, t, J=5Hz), 2.98 (2H, br), 3.30 (3H, s), 3.40-3.55 (2H, m), 3.78 (3H, s), 3.80-4.01 (2H, m), 4.27 (2H, t, J=5Hz), 6.58-6.79 (4H, m), 7.41-7.49 (1H, m), 8.16 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)

35

- 209 -

- 40) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-(2-pyridyl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.97-2.08 (2H, m), 2.26 (3H, s), 2.85 (2H, t, J=5Hz), 3.40 (3H, s), 3.62 (3H, s), 4.26 (2H, t, J=5Hz), 4.96 (1H, d, J=12Hz), 5.14 (1H, d, J=12Hz), 6.54-6.62 (2H, m), 6.40 (1H, d, J=7Hz), 6.98-7.14 (5H, m), 7.39 (1H, d, J=8Hz), 7.39-7.49 (1H, m), 7.70 (2H, s), 7.98 (1H, d, J=8Hz), 8.22 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz), 8.68 (1H, br)

10

- 41) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-[(4-methylpiperazin-1-yl)carbonylamino]but-1-yl]oxy-4-methylphenyl]-N-methylbenzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.62-1.88 (4H, m), 2.30-2.15 (2H, m), 2.28 (6H, s), 2.34-2.42 (4H, m), 2.93 (2H, t, J=5Hz), 3.25-3.48 (6H, m), 3.33 (3H, s), 3.79 (3H, s), 3.79-3.99 (2H, m), 4.30 (2H, t, J=5Hz), 6.58-6.70 (2H, m), 6.90-7.11 (5H, m), 7.45 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

20

- 42) 4-[2-(3-Aminoprop-1-yl)oxybenzoylamino]-3-methoxy-N-[2-[4-[(4-dimethylaminopiperidin-1-yl)carbonylamino]but-1-yl]oxy-4-methylphenyl]-N-methylbenzamide

25 NMR (CDCl<sub>3</sub>, δ) : 1.44-1.98 (8H, m), 2.26 (3H, s), 2.49 (6H, s), 2.66-2.93 (3H, m), 3.05 (2H, t, J=5Hz), 3.25-3.32 (2H, m), 3.29 (3H, s), 3.79 (3H, s), 3.81-3.99 (2H, m), 4.15-4.29 (4H, m), 6.57-6.64 (2H, m), 6.91-7.12 (5H, m), 7.46 (1H, dd, J=2, 8Hz), 8.04 (1H, d, J=8Hz), 8.35 (1H, d, J=8Hz)

30

#### Example 47

The following compound was obtained by using 4-[2-(3-tert-butoxycarbonylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(3-tert-butoxycarbonylaminoprop-1-yl)oxy-4-

35

- 210 -

methyl]phenyl-N-methylbenzamide as a starting compound according to a similar manner to that of Example 45.

4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(3-aminoprop-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.87-1.98 (2H, m), 2.00-2.09 (2H, m), 2.25 (3H, s), 2.83-2.96 (4H, m), 3.30 (3H, s), 3.78 (3H, s), 3.87-4.10 (2H, m), 4.27 (2H, t, J=7.5Hz), 6.57-6.66 (2H, m), 6.90 (1H, m), 7.00-7.10 (3H, m), 7.42 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.39 (1H, d, J=7Hz)

#### Example 48

The following compounds were obtained according to a similar manner to that of Example 47.

1) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-(4-aminoacetylaminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

MASS (m/z) : 592 (M+1)

2) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(piperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.48-1.95 (6H, m), 2.07-2.20 (2H, m), 2.28 (3H, s), 2.32-2.63 (5H, m), 2.75-3.01 (3H, m), 3.21 (3H, s), 3.40-3.64 (4H, m), 3.78 (3H, s), 3.83-4.08 (2H, m), 4.27 (2H, t, J=5Hz), 6.55-6.70 (2H, m), 6.82-7.17 (6H, m), 7.20-7.50 (1H, m), 8.29 (1H, d, J=7Hz), 8.39 (1H, d, J=8Hz)

3) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-(3-aminopropionyl)aminobut-1-yl]oxy-4-methylphenyl]-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.64-1.88 (4H, m), 2.06-2.19 (2H, m),



- 211 -

2.28 (3H, s), 2.32-2.46 (2H, m), 2.90-3.13 (4H, m),  
 3.23-3.44 (2H, m), 3.30 (3H, s), 3.77 (3H, s),  
 3.78-4.01 (2H, m), 4.27 (2H, br), 6.55-6.68 (2H,  
 m), 6.88-7.11 (5H, m), 7.28-7.50 (2H, m), 8.20 (1H,  
 d, J=8Hz) 8.31 (1H, d, J=8Hz)

4) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
 [4-(piperidin-4-yl)carbonylaminobut-1-yl]oxy-4-  
 methylphenyl]-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.60-1.91 (8H, m), 2.09-2.21 (2H, m),  
 2.28 (3H, s), 2.70 (1H, br), 2.97 (2H, t, J=5Hz),  
 3.11-3.40 (8H, m), 3.30 (3H, s), 3.72-3.96 (2H, m),  
 3.78 (3H, s), 4.28 (2H, t, J=5Hz), 6.57-6.65 (2H,  
 m), 6.90-7.08 (4H, m), 7.23-7.28 (2H, m), 7.38-7.49  
 (2H, m), 8.13 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

5) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
 (4-guanidinobut-1-yl)oxy-4-methylphenyl]-N-  
 methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.62-1.80 (4H, m), 2.05-2.14 (2H, m),  
 2.20 (3H, s), 2.55-2.70 (2H, m), 2.94 (2H, t,  
 J=5Hz), 3.31 (3H, s), 3.62-3.73 (2H, m), 3.72 (3H,  
 s), 4.22 (1H, d, J=5Hz), 6.48 (1H, d, J=8Hz), 6.61  
 (1H, s), 6.75 (1H, d, J=8Hz), 6.95-7.09 (5H, m),  
 7.43 (1H, dd, J=2, 8Hz), 8.03 (1H, d, J=8Hz), 8.32  
 (1H, d, J=8Hz)

#### Example 49

A solution of 4-hydroxy-3-methoxy-N-methyl-N-[4-methyl-  
 2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-  
 benzamide (320 mg) in N,N-dimethylformamide (8 ml) was  
 treated with sodium hydride (29.1 mg, 60% w/w in mineral oil)  
 at 0°C. The reaction mixture was stirred at 0°C for 15  
 minutes and then at ambient temperature for 10 minutes.  
 o-Nitrobenzyl bromide (143 mg) was added, and the reaction

- 212 -

mixture was stirred for 2.5 hours. The reaction was quenched with water and the mixture was diluted with ethyl acetate. The organic phase was washed with saturated aqueous sodium hydrogen carbonate, and brine. The organic solution was dried over magnesium sulfate, concentrated, and purified by silica gel column chromatography (SiO<sub>2</sub> 15 g, 3% methanol in dichloromethane) to give 3-methoxy-4-(2-nitrobenzyloxy)-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (210 mg).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.43-1.59 (2H, m), 1.61-1.88 (4H, m), 2.21-2.44 (12H, m), 3.31 (3H, s), 3.42-3.52 (2H, m), 3.56-3.67 (2H, m), 3.71 (3H, s), 3.78-4.00 (2H, m), 5.46 (3H, s), 6.52-6.67 (3H, m), 6.77-6.91 (2H, m), 6.95 (1H, br s), 7.46 (1H, m), 7.64 (1H, m), 7.84 (1H, d, J=8Hz), 8.14 (1H, d, J=8Hz)

#### Example 50

To a solution of 4-[2-(3-aminopropylthio)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (160 mg) in methanol (5 ml) was added a suspension of sodium metaperiodate (50.6 mg) and 5 ml of water. The mixture was stirred for 20 hours at ice-bath temperature and diluted with chloroform. The lower chloroform layer was removed, and the water layer was extracted with chloroform. The combined organic extracts were dried over anhydrous sodium sulfate. The solvent was removed at reduced pressure, and purified by preparative thin layer chromatography (methanol/dichloromethane/ammonia = 10/90/2) to give free amine (70 mg). To a solution of this amine in ethanol (3 ml) was added 1N hydrochloric acid (0.2 ml) and stirred for 5 minutes. The solution was concentrated to give 4-[2-(3-aminopropylsulfinyl)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride.

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.38-1.67 (4H, m), 1.68-1.88 (2H,

- 213 -

5 m), 1.94-2.13 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7Hz), 2.69-3.12 (9H, m), 3.12-3.58 (7H, m), 3.62 (3H, s), 3.80-4.17 (3H, m), 4.43 (1H, m), 6.64 (1H, d, J=8Hz), 6.83 (1H, s), 6.91 (2H, br s), 7.04 (1H, d, J=8Hz), 7.53 (1H, m), 7.68 (1H, dd, J=8, 8Hz), 7.85 (1H, dd, J=8, 8Hz), 7.90-8.19 (3H, s), 9.84 (1H, s)

Example 51

10 To a solution of 3-methoxy-4-[2-[3-(phthalimido)prop-1-yl]thiobenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (150 mg) in dichloromethane (10 ml) was added m-chloroperbenzoic acid (80.3 mg) and the mixture was stirred  
15 at ambient temperature for 2 hours. The solution was washed successively with saturated aqueous sodium hydrogen carbonate, water and brine, and the organic phase was dried over magnesium sulfate. The solvent was evaporated in vacuo and the residue was purified by silica gel column (2%  
20 methanol in chloroform) to give 3-methoxy-4-[2-[3-(phthalimido)prop-1-yl]sulfonylbenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (150 mg).

MASS (m/z) : 839 (M+1)

25

Example 52

A solution of 4-[2-[2-[(3-aminoprop-1-yl)oxy]phenyl]-vinyl]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenyl]benzamide (100 mg), 20% palladium hydroxide (30 mg) in methanol (5 ml)  
30 was stirred under atmospheric pressure of hydrogen at ambient temperature. After 12 hours, the reaction mixture was filtered through a bed of Celite, and the solvent was removed by rotary evaporation and the crude product was purified by  
35 NH-silica gel (chromatorex) column chromatography (SiO<sub>2</sub> 10 g,

- 214 -

1% methanol in chloroform) to give free amine. To the solution of amine (80 mg) in ethanol (3 ml) was added 1N hydrochloric acid (0.25 ml) and stirred for 5 minutes. The solution was evaporated to give 4-[2-[2-[(3-aminoprop-1-yl)oxy]phenyl]ethyl]-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride (70 mg).

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.36-1.65 (4H, m), 1.65-1.82 (2H, m), 1.97-2.13 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.58-3.11 (13H, m), 3.17 (3H, s), 3.26-3.68 (5H, m), 3.72-4.21 (5H, m), 4.42 (1H, m), 6.63 (1H, d, J=8Hz), 6.70-7.05 (8H, m), 7.13 (1H, dd, J=8, 8Hz), 8.00-8.24 (2H, m)

#### Example 53

The following compounds were obtained according to a similar manner to that of Example 10.

- 1) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-amino-3-methoxy-N-methyl-N-(4-hydroxyphenyl)benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.43 (9H, s), 1.60-1.68 (2H, m), 3.16-3.25 (2H, m), 3.49 (3H, s), 3.63 (3H, s), 4.16-4.23 (2H, m), 4.73-4.80 (1H, br), 6.67-6.74 (3H, m), 6.84-7.01 (5H, m), 7.07-7.14 (2H, m), 7.47 (1H, t, J=8Hz), 8.16 (1H, d, J=8Hz), 8.52 (1H, d, J=8Hz)

ESI-MASS (m/z) : 550 (M+H)

- 2) 3-Methoxy-4-[2-[1-(tert-butoxycarbonyl)piperidin-4-yl]oxybenzoyl]amino-N-(2-hydroxy-4-methyl)phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.43 (9H, s), 1.68-2.10 (4H, m), 2.23 (3H, s), 2.96-3.17 (2H, m), 3.36 (3H, s), 3.64-3.98 (5H, m), 4.60 (1H, m), 6.36-7.03 (7H, m), 7.10 (1H, t, J=7Hz), 7.43 (1H, t, J=7Hz), 8.19 (1H, d, J=7Hz)

- 215 -

- 3) 4-[2-(3-Amino-1-methylprop-1-yl)oxybenzoyl]amino-3-methoxy-N-(2-hydroxy-4-methylphenyl)-N-methylbenzamide

NMR (DMSO-d<sub>6</sub>, δ) : 1.33 (3H, d, J=7.5Hz), 1.63-1.76 (1H, m), 1.87-1.98 (1H, m), 2.14 (3H, s), 2.65 (2H, t, J=7.5Hz), 3.18 (3H, s), 3.74 (3H, s), 4.96 (1H, m), 6.47 (1H, d, J=7Hz), 6.63 (1H, s), 6.86 (1H, d, J=7Hz), 7.91 (1H, d, J=7Hz), 7.01 (1H, s), 7.09 (1H, t, J=7Hz), 7.32 (1H, d, J=7Hz), 7.52 (1H, t, J=7Hz), 8.04 (1H, d, J=7Hz), 8.30 (1H, d, J=7Hz)

- 4) 3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)amino-1-methylprop-1-yl]oxybenzoyl]amino-N-(2-hydroxy-4-methylphenyl)-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.36 (3H, d, J=7.5Hz), 1.40 (9H, s), 1.80-2.10 (2H, m), 2.22 (3H, s), 3.16-3.28 (2H, m), 3.35 (3H, s), 3.69 (3H, s), 4.64 (1H, m), 4.79 (1H, br), 6.52 (1H, m), 6.70-6.82 (2H, m), 6.91-7.11 (4H, m), 7.41 (1H, t, J=7Hz), 8.21 (1H, d, J=7Hz), 8.47 (1H, m)

- 5) 4-(2-Hydroxybenzoylamino-3-methoxy-N-(2-hydroxy-4-methylphenyl)-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 2.26 (3H, s), 3.36 (3H, s), 6.56 (1H, m), 6.65-6.86 (4H, m), 6.96-7.08 (2H, m), 7.35-7.44 (2H, m), 8.20 (1H, br), 8.61 (1H, br)

- 6) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-(2-hydroxy-4-methylphenyl)-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.42 (9H, s), 1.68 (2H, br), 1.99 (2H, br), 2.22 (3H, s), 3.19 (2H, br), 3.39 (3H, s), 3.49 (2H, br), 5.03 (1H, br), 6.43-6.72 (6H, m), 7.08 (2H, br), 7.39 (1H, br), 8.21 (1H, d, J=8Hz), 8.45 (1H, br)

Example 54

The following compounds were obtained according to a similar manner to that of Example 12.

- 5        1) 4-[2-[(3-tert-Butoxycarbonylamino-*prop*-1-yl)oxy]benzoyl]-  
amino-3-methoxy-N-methyl-N-[4-(5-ethoxycarbonylpent-1-  
yloxy)phenyl]benzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.21-1.29 (3H, m), 1.40 (9H, s),  
1.42-1.90 (8H, m), 2.09-2.19 (2H, m), 3.27-3.34  
10        (2H, m), 3.47 (3H, s), 3.82 (3H, s), 3.89 (2H, t,  
J=8Hz), 4.08-4.17 (2H, m), 4.26 (2H, t, J=8Hz),  
4.70-4.77 (1H, br), 6.75 (2H, d, J=8Hz), 6.83 (1H,  
d, J=8Hz), 6.94-7.02 (3H, m), 7.07-7.13 (2H, m),  
15        7.46 (1H, t, J=8Hz), 8.21 (1H, d, J=8Hz), 8.42 (1H,  
d, J=8Hz)  
ESI-MASS (m/z) : 692 (M+H)
- 20        2) 4-[2-Benzyloxy]benzoyl]amino-N-[2-(3-ethoxycarbonyl*prop*-  
1-yl)oxy]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7.5Hz), 2.03-2.17 (2H,  
m), 2.50 (2H, t, J=7.5Hz), 3.32 (3H, s), 3.87-4.04  
(2H, m), 4.16 (2H, q, J=7.5Hz), 5.19 (2H, s), 6.78  
(2H, d, J=8Hz), 6.92-7.00 (3H, m), 7.07-7.21 (5H,  
25        m), 7.38-7.53 (6H, m), 8.26 (1H, d, J=7Hz)
- 30        3) 4-(2-Iodobenzoyl)amino-N-[2-(5-ethoxycarbonylpent-1-  
yl)oxy]phenyl-N-methylbenzamide  
NMR (CDCl<sub>3</sub>, δ) : 1.24 (3H, t, J=7.5Hz), 1.42-1.55 (2H,  
m), 1.63-1.72 (2H, m), 1.76-1.88 (2H, m), 2.31 (2H,  
t, J=7.5Hz), 3.31 (3H, s), 3.81-3.99 (2H, m), 4.11  
(2H, q, J=7.5Hz), 6.76-6.83 (2H, m), 7.00 (1H, d,  
J=7Hz), 8.08-7.17 (2H, m), 7.29-7.49 (5H, m), 7.66  
35        (1H, s), 7.88 (1H, d, J=7Hz)
- 4) 3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)amino*prop*-1-yl]-

- 217 -

oxybenzoyl]amino-N-[2-[3-(tert-butoxycarbonyl)aminoprop-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.39 (9H, s), 1.41 (9H, s), 1.93-1.97 (2H, m), 2.07-2.17 (2H, m), 2.26 (3H, s), 3.22-3.32 (4H, m), 3.30 (3H, s), 3.78 (3H, s), 3.82-4.05 (2H, m), 6.60-6.66 (2H, m), 6.86-6.91 (2H, m), 7.00 (1H, d, J=7Hz), 7.03-7.10 (2H, m), 7.43 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.39 (1H, d, J=7Hz)

- 10 5) 3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-N-methyl-N-[4-methyl-2-[4-(phthalimido)but-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.85-1.92 (2H, m), 2.10-2.17 (2H, m), 2.27 (3H, s), 3.22-3.32 (2H, m), 3.28 (3H, s), 3.74-3.81 (2H, m), 3.81 (3H, s), 3.92-4.15 (2H, m), 4.24 (2H, t, J=7.5Hz), 6.57-6.65 (2H, m), 6.83-6.90 (2H, m), 6.97-7.14 (3H, m), 7.24 (1H, t, J=7Hz), 7.69-7.77 (2H, m), 7.82-7.91 (2H, m), 8.21 (1H, d, J=7Hz), 8.40 (1H, d, J=7Hz)

- 20 6) 3-Methoxy-4-[2-[1-(tert-butoxycarbonyl)piperidin-4-yl]oxybenzoyl]amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.25 (3H, t, J=7.5Hz), 1.42-1.91 (6H, m), 1.45 (9H, s), 2.02-2.12 (2H, m), 2.27 (3H, s), 2.27-2.88 (2H, m), 2.97-3.18 (2H, m), 3.32 (3H, s), 3.40 (2H, t, J=7Hz), 3.74 (3H, s), 3.89-4.00 (2H, m), 4.13 (2H, q, J=7.5Hz), 4.66 (1H, m), 6.59 (1H, d, J=7Hz), 6.61 (1H, s, J=7Hz), 6.80-6.92 (2H, m), 6.98-7.12 (3H, m), 7.43 (1H, t, J=7Hz), 8.19 (1H, d, J=7Hz), 8.39 (1H, d, J=7Hz)

- 35 7) 3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)amino-1-methylprop-1-yl]oxybenzoyl]amino-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

- 218 -

5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.24 (3H, t, J=7.5Hz), 1.38 (9H, s),  
1.40 (2H, d, J=7.5Hz), 1.41-2.10 (8H, m), 2.26 (3H, s),  
2.27-2.33 (2H, m), 3.23-3.30 (2H, m), 3.30 (3H, s),  
3.79 (3H, s), 3.83-3.99 (2H, m), 4.12 (2H, q, J=7.5Hz),  
4.62-4.77 (2H, m), 6.58-6.63 (2H, m), 6.82 (1H, t, J=7Hz),  
7.01 (1H, d, J=7Hz), 7.05-7.12 (2H, m), 7.43 (1H, t, J=7Hz),  
8.21 (1H, d, J=7Hz), 8.39 (1H, d, J=7Hz)

10 8) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-  
amino-3-methoxy-N-(2-methoxy-4-methylphenyl)-N-  
methylbenzamide

15 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 2.08-2.20 (2H, m), 2.29  
(3H, s), 3.28 (2H, q, J=5Hz), 3.31 (3H, s), 3.75 (3H, s),  
3.80 (3H, s), 4.25 (2H, t, J=5Hz), 4.74 (1H, br),  
6.59-6.65 (2H, m), 6.89 (1H, d, J=8Hz), 7.00 (1H, d, J=8Hz),  
7.06-7.13 (2H, m), 7.46 (1H, dd, J=2, 8Hz),  
8.21 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

20 9) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-  
amino-3-methoxy-N-[2-[4-(2-pyridyl)phenylmethyl]oxy-4-  
methylphenyl]-N-methylbenzamide

25 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.39 (9H, s), 2.09 (2H, t, J=5Hz),  
2.29 (3H, s), 3.27 (2H, q, J=5Hz), 3.40 (3H, s),  
3.61 (3H, s), 4.21 (2H, t, J=5Hz), 4.82 (1H, br),  
4.97 (1H, d, J=12Hz), 5.14 (1H, d, J=12Hz), 6.55-  
6.74 (2H, m), 6.89-7.12 (7H, m), 7.19-7.24 (1H, m),  
7.39 (1H, d, J=8Hz), 7.41-7.49 (1H, m), 7.70 (2H, s),  
7.99 (1H, d, J=8Hz), 8.21 (1H, d, J=8Hz), 8.40  
30 (1H, d, J=8Hz), 8.67 (1H, d, J=5Hz)

35 10) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-  
amino-3-methoxy-N-[2-[4-(1,5-dimethyl-3-cyanopyrrol-2-  
yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide  
NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 2.03-2.15 (2H, m), 2.13



- 219 -

(3H, s), 2.30 (3H, s), 3.26 (2H, q, J=5Hz), 3.40 (3H, s), 3.46 (3H, s), 3.58 (3H, s), 4.19 (2H, t, J=5Hz), 4.86 (1H, d, J=12Hz), 5.10 (1H, d, J=12Hz), 6.65-6.73 (2H, m), 6.82 (1H, d, J=8Hz), 6.95-7.10 (4H, m), 7.34-7.44 (6H, m), 8.00 (1H, s), 8.19 (1H, d, J=8Hz), 8.36 (1H, d, J=8Hz)

11) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-

yl]oxybenzoyl]amino-3-methoxy-N-[2-[4-(thiazol-2-

yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 2.05-2.16 (2H, m), 3.27 (2H, q, J=5Hz), 3.40 (3H, s), 3.62 (3H, s), 4.20 (2H, t, J=5Hz), 4.76 (1H, br), 4.89 (1H, d, J=12Hz), 5.07 (1H, d, J=12Hz), 6.62-6.72 (2H, m), 6.89 (1H, d, J=8Hz), 6.96-7.11 (4H, m), 7.28 (1H, d, J=3Hz), 7.31 (2H, d, J=8Hz), 7.42 (1H, dd, J=2, 8Hz), 7.81 (1H, d, J=8Hz), 7.93 (2H, d, J=8Hz), 8.00 (1H, s), 8.20 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

12) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-

yl]oxybenzoyl]amino-3-methoxy-N-[2-[4-(oxazol-2-

yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 2.05-2.16 (2H, m), 2.29 (3H, s), 3.27 (2H, q, J=5Hz), 3.40 (3H, s), 3.65 (3H, s), 4.21 (2H, t, J=5Hz), 4.78 (1H, br), 4.90 (1H, d, J=13Hz), 5.10 (1H, d, J=13Hz), 6.64 (1H, s), 6.70 (1H, d, J=8Hz), 6.85 (1H, d, J=8Hz), 6.98-7.17 (5H, m), 7.20 (1H, s), 7.30-7.49 (3H, m), 7.63 (1H, s), 8.03 (1H, d, J=8Hz), 8.22 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

13) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-

yl]oxybenzoyl]amino-3-methoxy-N-[2-[4-(pyrimidin-2-

yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide

- 220 -

- 5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 2.05-2.16 (2H, m), 2.28 (3H, s), 3.28 (2H, q, J=5Hz), 3.40 (3H, s), 3.65 (3H, s), 4.22 (2H, t, J=5Hz), 4.78 (1H, br), 4.95 (1H, d, J=12Hz), 5.14 (1H, d, J=12Hz), 6.65-6.70 (2H, m), 6.88 (1H, d, J=8Hz), 6.96-7.19 (5H, m), 7.38-7.46 (3H, m), 8.21 (1H, d, J=8Hz), 8.35-8.44 (3H, m), 8.74 (1H, d, J=3Hz)
- 10 14) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-[2-(4-cyanophenylmethyl)oxy-4-methylphenyl]-N-methylbenzamide
- 15 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.41 (9H, s), 2.08-2.20 (2H, m), 2.30 (3H, s), 3.30 (2H, q, J=5Hz), 3.40 (3H, s), 3.68 (3H, s), 4.26 (2H, t, J=5Hz), 4.89 (1H, d, J=13Hz), 5.09 (1H, d, J=13Hz), 6.60 (1H, s), 6.73 (1H, d, J=8Hz), 6.98-7.12 (5H, m), 7.39-7.52 (3H, m), 7.68 (1H, d, J=8Hz), 8.20 (1H, d, J=8Hz), 8.34 (1H, d, J=8Hz)
- 20 15) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-[2-(4-phthalimidobut-1-yl)oxy-4-methylphenyl]-N-methylbenzamide
- 25 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 1.72-1.95 (4H, m), 2.08-2.19 (2H, m), 2.29 (3H, s), 3.31 (2H, q, J=5Hz), 3.33 (3H, s), 3.79 (2H, t, J=5Hz), 3.81 (3H, s), 3.84-4.06 (2H, m), 4.25 (2H, t, J=5Hz), 4.82 (1H, br), 6.57 (1H, d, J=8Hz), 6.62 (1H, s), 6.81-6.89 (2H, m), 6.97 (1H, d, J=8Hz), 7.04-7.10 (2H, m), 7.40-7.48 (1H, m), 7.68-7.74 (2H, m), 7.81-7.89 (2H, m), 8.20 (1H, d, J=8Hz), 8.39 (1H, d, J=8Hz)
- 30
- 35 16) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-[2-(3-methoxycarbonylpyrid-6-yl)methoxy-4-methylphenyl]-N-methylbenzamide

- 221 -

NMR (CDCl<sub>3</sub>, δ) : 1.39 (9H, s), 2.07-2.16 (2H, m), 2.27 (3H, s), 3.29 (2H, q, J=5Hz), 3.42 (3H, s), 3.63 (3H, s), 3.89 (3H, s), 4.24 (2H, t, J=5Hz), 4.95 (1H, d, J=12Hz), 5.08 (1H, d, J=12Hz), 6.58 (1H, s), 6.73 (1H, d, J=8Hz), 6.89 (1H, d, J=8Hz), 6.98 (2H, d, J=8Hz), 7.05-7.12 (3H, m), 7.34 (1H, d, J=8Hz), 7.44 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.28 (1H, d, J=8Hz), 8.37 (1H, d, J=8Hz), 9.14 (1H, s)

#### Example 55

The following compound was obtained according to a similar manner to that of Example 35.

4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl-amino-3-methoxy-N-[2-[4-(tert-butoxycarbonylguanidino)but-1-yl]oxy-4-methylphenyl]-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.43 (9H, s), 1.44 (9H, s), 1.52-1.60 (2H, m), 1.65-1.74 (2H, m), 1.92-2.07 (2H, m), 2.21 (3H, s), 3.10-3.25 (4H, m), 3.38 (3H, s), 3.50 (2H, br), 3.66 (3H, br), 3.78-4.05 (2H, m), 6.49 (2H, br), 6.63-6.82 (3H, m), 7.01-7.10 (2H, m), 7.38 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.44 (1H, br)

#### Example 56

The following compounds were obtained according to a similar manner to that of Example 10.

1) 4-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl]oxy]benzoyl]-amino-3-hydroxy-N-methyl-N-cyclohexylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.07-1.17 (2H, m), 1.41 (9H, s), 1.47-1.76 (8H, m), 2.10-2.20 (2H, m), 2.92-3.00 (2H, m), 3.36-3.44 (2H, m), 3.49 (3H, s), 4.19-4.27 (2H, m), 4.98-5.06 (1H, br), 6.87-6.92 (1H, br), 6.98-7.03 (2H, m), 7.12 (1H, t, J=8Hz), 7.47 (1H,

- 222 -

t, J=8Hz), 8.12-8.22 (1H, br), 8.28 (1H, d, J=8Hz),  
9.72-9.80 (1H, br)

ESI-MASS (m/z) : 526 (M+H)

- 5        2) 4-[2-[(3-tert-Butoxycarbonylamino-1-yl)oxy]benzoyl]-  
amino-3-hydroxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-  
yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

10        NMR (CDCl<sub>3</sub>, δ) : 1.42 (9H, s), 1.50-1.90 (8H, m),  
2.20-2.22 (2H, m), 2.27 (3H, s), 2.32 (3H, s),  
2.35-2.53 (6H, m), 3.29 (3H, s), 3.32-3.42 (2H, m),  
3.50-3.66 (3H, m), 3.72 (2H, br), 3.89 (1H, br),  
4.20 (2H, t, J=6Hz), 5.29 (1H, br), 6.54 (1H, s),  
6.67 (1H, d, J=7Hz), 6.72 (1H, br), 6.96-7.10 (4H,  
15        m), 7.40-7.47 (1H, m), 8.10 (1H, br), 8.27 (1H, d,  
J=6Hz)

#### Example 57

20        To a solution of 4-[(2-benzyloxy)benzoyl]amino-3-[(2-  
benzyloxy)benzoyl]oxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-  
yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide (1.2 g) in  
ethanol (20 ml) was added 1N sodium hydroxide solution (10  
ml) and the mixture was stirred at ambient temperature for 2  
hours. The mixture was concentrated in vacuo and the  
25        solution was adjusted to pH 7 with 1N hydrochloric acid. The  
solution was extracted with ethyl acetate (20 ml) and the  
organic layer was washed with brine (20 ml). The organic  
layer was dried over magnesium sulfate and the solution was  
concentrated in vacuo to give 4-[(2-benzyloxy)benzoyl]amino-  
3-hydroxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-  
30        yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide (930 mg)

35        NMR (CDCl<sub>3</sub>, δ) : 1.48-1.59 (2H, m), 1.70 (4H, br),  
2.29-2.42 (13H, m), 3.29 (3H, s), 3.48 (2H, br),  
3.53 (2H, br), 3.80 (1H, br), 3.90 (1H, br), 5.28  
(2H, s), 6.53-6.65 (3H, m), 6.72 (1H, br), 6.90-  
7.12 (4H, m), 7.34-7.37 (3H, m), 7.40-7.49 (4H, m),

- 223 -

8.20-8.27 (1H, m)

Example 58

5 The following compounds were obtained according to a similar manner to that of Example 12.

- 1) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-amino-3-ethoxycarbonylmethoxy-N-methyl-N-cyclohexylbenzamide

10 NMR (CDCl<sub>3</sub>, δ) : 1.29 (3H, t, J=8Hz), 1.41 (9H, s),  
1.45-1.85 (10H, m), 2.07-2.12 (2H, m), 2.86-3.06  
(3H, br), 3.25-3.32 (2H, m), 4.22-4.33 (4H, m),  
4.76 (2H, s), 4.98-5.07 (1H, br), 6.91 (1H, s),  
15 7.01-7.15 (3H, m), 7.48 (1H, t, J=8Hz), 8.23 (1H,  
d, J=8Hz), 8.69 (1H, d, J=8Hz)

ESI-MASS (m/z) : 634 (M+Na)

- 2) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-amino-3-isopropoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.19-1.28 (6H, m), 1.38 (9H, s),  
1.46-1.58 (2H, m), 1.65-1.88 (6H, m), 1.99-2.10  
(2H, m), 2.25 (3H, s), 2.29 (3H, s), 2.32-2.42 (6H,  
m), 3.15-3.23 (2H, m), 3.31 (3H, s), 3.45-3.50 (2H,  
25 m), 3.60-3.64 (2H, m), 3.84-3.97 (2H, m), 4.24-4.36  
(3H, m), 6.56-6.65 (2H, m), 6.85 (1H, d, J=7Hz),  
6.94-7.02 (3H, m), 7.10 (1H, t, J=6Hz), 7.47 (1H,  
t, J=7Hz), 8.15 (1H, d, J=7Hz), 8.41 (1H, d, J=7Hz)

- 30 3) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-3-propoxybenzamide

35 NMR (CDCl<sub>3</sub>, δ) : 0.97 (3H, t, J=7Hz), 1.42 (9H, s),  
1.47-1.58 (2H, m), 1.67-1.88 (8H, m), 1.98-2.10  
(2H, m), 2.27 (3H, s), 2.28 (3H, s), 2.31-2.41 (6H,

- 224 -

5 m), 3.16-3.26 (2H, m), 3.31 (3H, s), 3.45-3.50 (2H, m), 3.58-3.65 (2H, m), 3.84-3.97 (4H, m), 4.26 (2H, t, J=7Hz), 6.58 (1H, d, J=7Hz), 6.64 (1H, s), 6.84 (1H, d, J=6Hz), 6.95 (1H, d, J=7Hz), 6.99-7.03 (2H, m), 7.09 (1H, t, J=7Hz), 7.45 (1H, t, J=7Hz), 8.16 (1H, d, J=7Hz), 8.38 (1H, d, J=7Hz)

10 4) 4-[2-[(3-tert-Butoxycarbonylamino-3-(3-ethoxycarbonylprop-1-yl)oxy)benzoyl]-amino-3-(3-ethoxycarbonylprop-1-yl)oxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

15 NMR (CDCl<sub>3</sub>, δ) : 1.11 and 1.23 (total 3H, t, J=6Hz), 1.40 (9H, s), 1.48-1.60 (2H, m), 1.60-1.75 (4H, m), 1.75-1.88 (2H, m), 1.98-2.10 (4H, m), 2.26 (3H, s), 2.29 (3H, s), 2.32-2.42 (8H, m), 3.18-3.28 (2H, m), 3.30 (3H, s), 3.45-3.50 (2H, m), 3.62 (2H, br), 3.88-4.10 (5H, m), 4.27 (2H, t, J=6Hz), 6.57 (1H, d, J=7Hz), 6.63 (1H, s), 6.82 (1H, d, J=7Hz), 6.87-6.92 (1H, m), 6.98-7.10 (3H, m), 7.42 (1H, t, J=6Hz), 8.10-8.13 (1H, m), 8.37 (1H, d, J=7Hz)

25 5) 4-[2-[(3-tert-Butoxycarbonylamino-3-(3-ethoxycarbonylmethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

30 NMR (CDCl<sub>3</sub>, δ) : 1.29 (3H, t, J=7Hz), 1.39 (9H, s), 1.46-1.90 (8H, m), 2.00-2.10 (2H, m), 2.28 (3H, s), 2.29 (3H, s), 2.30-2.42 (6H, m), 3.18-3.29 (2H, m), 3.30 (3H, s), 3.42-3.50 (2H, m), 3.58-3.65 (2H, m), 3.85-3.97 (2H, m), 4.18-4.29 (4H, m), 4.52 (2H, s), 6.52-6.13 (2H, m), 6.80 (1H, d, J=7Hz), 6.89-6.99 (3H, m), 7.38-7.48 (1H, m), 8.15 (1H, d, J=7Hz), 8.41 (1H, d, J=7Hz)

35 6) 4-[(2-Benzyloxy)benzoyl]amino-3-ethoxyl-N-methyl-N-[2-

- 225 -

[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.08 (3H, t, J=6Hz), 1.45-1.57 (2H, m), 1.60-1.75 (2H, m), 1.77-1.87 (2H, m), 2.25 (3H, s), 2.29 (3H, s), 2.31-2.39 (7H, m), 3.30 (3H, s), 3.46-3.49 (2H, m), 3.60-3.63 (2H, m), 3.70-3.80 (2H, m), 3.82-3.98 (2H, m), 5.34 (2H, s), 6.52-6.60 (2H, m), 6.80-7.10 (5H, m), 7.27-7.38 (6H, m), 8.20-8.22 (1H, m), 8.38-8.43 (1H, m)

Example 59

The following compounds were obtained according to a similar manner to that of Example 4.

- 1) 4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-amino-3-carboxymethoxy-N-methyl-N-cyclohexylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.03-1.17 (2H, m), 1.39 (9H, s), 1.45-1.85 (8H, m), 2.03-2.12 (2H, m), 2.85-2.98 (3H, m), 3.21-3.33 (2H, m), 4.23-4.31 (2H, m), 4.73 (3H, s), 5.08-5.13 (1H, br), 6.98-7.07 (3H, m), 7.10 (1H, t, J=8Hz), 7.48 (1H, t, J=8Hz), 8.18-8.24 (1H, m), 8.56-8.61 (1H, m)

ESI-MASS (m/z) : 606 (M+Na)

- 2) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-carboxymethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.49 (2H, m), 1.49-1.62 (2H, m), 1.67-1.78 (2H, m), 2.02-2.34 (13H, m), 2.78-2.89 (2H, m), 3.38-3.43 (4H, m), 3.58 (3H, s), 3.89-3.96 (2H, m), 4.00-4.18 (2H, m), 4.30 (2H, br), 6.62 (1H, d, J=6Hz), 6.72-6.87 (3H, m), 6.89-6.97 (1H, m), 7.11 (1H, t, J=7Hz), 7.19 (1H, d, J=7Hz), 7.54 (1H, t, J=6Hz), 7.94 (1H, d, J=6Hz),

- 226 -

8.22 (1H, d, J=7Hz)

- 3) 4-[2-[(3-tert-Butoxycarbonylamino-1-yl)oxy]benzoyl]-  
amino-3-carboxymethoxy-N-methyl-N-[2-[5-(4-  
methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-  
methylphenyl]benzamide  
MASS (m/z) : 804 (M+H)

Example 60

To a mixture of 4-(2-iodobenzoyl)amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (1.12 g) and 3-butyne-1-ol (153 mg) in a mixture of tetrahydrofuran (15 ml) and ethylamine (15 ml) were added bis(triphenylphosphine)palladium(II) chloride (23.5 mg) and copper (I) iodide (3.19 mg) and the mixture was refluxed for 8 hours. The solution was diluted with chloroform (50 ml) and the solution was washed with water and brine. The solution was dried over magnesium sulfate and the solvent was evaporated in vacuo to give an oil. The oil was purified by silica gel column (2% methanol in chloroform) to give 4-[2-(4-hydroxy-1-butyn-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (755 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.44-1.57 (2H, m), 1.61-1.86 (4H, m), 2.27 (3H, s), 2.29-2.40 (6H, m), 2.70 (2H, t, J=7.5Hz), 3.33 (3H, s), 3.44-3.49 (2H, m), 3.53-3.60 (2H, m), 3.74 (2H, t, J=7.5Hz), 3.79-3.99 (2H, m), 6.76-6.84 (2H, m), 7.06 (1H, d, J=7Hz), 7.13 (1H, t, J=7Hz), 7.34 (2H, d, J=8Hz), 7.40-7.47 (2H, m), 7.48-7.56 (3H, m), 7.99 (1H, m), 9.19 (1H, s)

Example 61

To an ice cooled solution of 4-[2-(4-hydroxy-1-butyn-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (755 mg) in



- 227 -

dichloromethane (20 ml) were added triethylamine (150 mg) and methanesulfonyl chloride (156 mg), and the mixture was stirred in an ice bath for 2 hours. The solution was washed successively with water, 10% hydrochloric acid, saturated aqueous sodium hydrogen carbonate and brine, and the organic phase was dried over magnesium sulfate. The solvent was evaporated in vacuo to give 4-[2-(4-methanesulfonyloxy-1-butyn-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (789 mg).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.49-1.60 (2H, m), 1.67-1.86 (2H, m), 1.87-1.90 (2H, m), 2.37 (2H, t, J=7.5Hz), 2.68 (3H, s), 2.86-3.06 (6H, m), 2.92 (3H, s), 3.31 (3H, s), 3.77-4.02 (6H, m), 4.32 (2H, t, J=7.5Hz), 6.77-6.87 (2H, m), 7.04 (1H, d, J=7Hz), 7.17 (1H, t, J=7Hz), 7.32 (2H, d, J=8Hz), 7.41-7.53 (5H, m), 7.90 (1H, m), 8.86 (1H, s)

#### Example 62

The following compounds were obtained according to a similar manner to that of Example 61.

1) 4-[2-(4-Methanesulfonyloxybut-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

MASS (m/z) : 693 (M+1)

2) 4-[2-(3-Methanesulfonyloxyprop-1-yl)thiobenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.48-1.60 (2H, m), 1.65-1.74 (2H, m), 1.75-1.86 (2H, m), 1.98-2.07 (2H, m), 2.26 (3H, s), 2.30-2.39 (2H, m), 2.70-2.78 (4H, m), 2.79-3.42 (2H, m), 2.90 (3H, s), 2.95-3.07 (2H, m), 3.26 (3H, s), 3.71 (3H, s), 3.80-4.01 (4H, m), 4.29 (2H, t, J=7.5Hz), 6.56-6.66 (2H, m), 6.82-7.00 (3H, m),

- 228 -

7.30 (1H, m), 7.39-7.47 (2H, m), 7.60 (1H, d, J=7Hz), 8.27 (1H, d, J=7Hz), 8.58 (1H, s)

Example 63

5 A mixture of 4-[2-(4-methanesulfonyloxy-1-butyn-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (800 mg) and  
10 potassium phthalimide (430 mg) in dimethyl sulfoxide (20 ml) was stirred at 60°C for 5 hours, and the solution was diluted with ethyl acetate (60 ml). The solution was washed with water and brine, and dried over magnesium sulfate. The  
15 solvent was evaporated in vacuo to give 4-[2-[4-(phthalimido)-1-butyn-1-yl]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (620 mg).

20 NMR (CDCl<sub>3</sub>, δ) : 1.50-1.61 (2H, m), 1.67-1.92 (6H, m), 2.30 (3H, s), 2.33-2.44 (6H, m), 3.38 (3H, s), 3.48-3.52 (2H, m), 3.60-3.67 (2H, m), 3.84-4.01 (4H, m), 6.78-6.85 (2H, m), 7.02 (1H, d, J=7Hz), 7.09-7.19 (2H, m), 7.30-7.70 (6H, m), 7.70-7.77 (2H, m), 7.81-7.90 (2H, m), 8.18 (1H, m)

Example 64

25 The following compounds were obtained according to a similar manner to that of Example 63.

1) 4-[2-[4-(Phthalimido)but-1-yl]benzoyl]amino-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenyl-N-methylbenzamide

30 MASS (m/z) : 693 (M+1)

2) 3-Methoxy-4-[2-[3-(phthalimido)prop-1-yl]thiobenzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

35 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.59 (2H, m), 1.61-1.74 (2H, m),

- 229 -

1.78-1.87 (2H, m), 1.92-2.03 (2H, m), 2.26 (3H, s),  
2.29 (3H, s), 2.31-2.42 (6H, m), 2.94 (2H, t,  
J=7.5Hz), 3.31 (3H, s), 3.45-3.53 (2H, m), 3.58-  
3.67 (2H, m), 3.69-3.81 (2H, m), 3.73 (3H, s),  
3.84-4.00 (2H, m), 6.55-6.66 (2H, m), 6.80-6.92  
(2H, m), 7.02 (1H, s), 7.27 (1H, m), 7.34-7.44 (2H,  
m), 7.60-7.90 (5H, m), 8.25 (1H, d, J=7Hz), 8.82  
(1H, s)

10      Example 65

To an ice cooled mixture of 4-[2-(4-amino-1-butyn-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (310 mg), nickel chloride hexahydrate (181 mg) in a mixture of tetrahydrofuran (5 ml) and methanol (5 ml) was added sodium borohydride (96.2 mg) in small portions and the mixture was stirred at the same temperature for 2 hours. The mixture was filtered through bed of Celite and the filtrate was evaporated in vacuo. The residue was dissolved in chloroform (20 ml) and washed with water and brine. The organic solution was dried over magnesium sulfate and the solvent was evaporated in vacuo to give a syrup. The residue was purified by silica gel column (chloroform:methanol:ammonia = 100:10:1) to give 4-[2-(4-aminobut-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide (295 mg).

MASS (m/z) : 597 (M+1)

30      Example 66

The following compound was obtained according to a similar manner to that of Example 65.

4-[2-(4-hydroxybut-1-yl)benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

MASS (m/z) : 615 (M+1)

- 230 -

Example 67

A mixture of 4-amino-3-methoxy-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]-phenylbenzamide (200 mg) and salicyl aldehyde (48.6 mg) in methanol (10 ml) was refluxed overnight in the presence of 3Å molecular sieves (100 mg). The solution was filtered and the filtrate was treated with sodium borohydride (15.1 mg) at 5°C for 2 hours. The reaction mixture was diluted with chloroform (20 ml) and the solution was washed with water and brine. The organic solution was dried over magnesium sulfate and the solvent was evaporated in vacuo to give a crude oil. The product was purified by silica gel column (2% methanol in chloroform) to give 3-methoxy-4-(2-hydroxyphenyl)methylamino-N-methyl-N-[4-methyl-2-[4-(4-methylpiperazin-1-yl)carbonyl]phenylmethoxy]phenylbenzamide (152 mg).

NMR (CDCl<sub>3</sub>, δ) : 2.27 (3H, s), 2.32 (3H, s), 2.32-2.59 (4H, m), 3.34 (3H, s), 3.40-3.55 (2H, m), 3.52 (3H, s), 3.75-3.88 (2H, m), 4.25-4.34 (2H, m), 4.63 (1H, br), 4.42 (1H, d, J=14Hz), 5.08 (1H, d, J=14Hz), 6.43 (1H, d, J=7Hz), 6.62 (1H, s), 6.70 (1H, d, J=7Hz), 6.80-6.88 (4H, m), 7.00 (1H, d, J=7Hz), 7.09-7.18 (2H, m), 7.28 (2H, d, J=8Hz), 7.38 (2H, d, J=8Hz)

Example 68

The following compound was obtained according to a similar manner to that of Example 67.

3-Methoxy-4-(2-hydroxyphenyl)methylamino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy]phenylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.41-1.52 (2H, m), 1.60-1.69 (2H, m), 1.70-1.80 (2H, m), 2.25 (3H, s), 2.29 (3H, s), 2.30-2.43 (6H, m), 3.28 (3H, s), 3.34-3.48 (2H, m), 3.55 (3H, s), 3.65-4.00 (2H, m), 4.30 (2H, d,

- 231 -

J=7Hz), 4.62 (1H, br t, J=7Hz), 6.51 (1H, d, J=7Hz), 6.57-6.64 (2H, m), 6.76-6.95 (5H, m), 7.61-7.68 (2H, m)

5      Example 69

To an ice bath cooled solution of 4-(2-dimethylamino-4-methyl)phenoxyethyl-N-[2-(5-ethoxycarbonylpent-1-yl)oxy]phenylbenzamide (860 mg) in N,N-dimethylformamide (15 ml) was added sodium hydride (60% in oil, 71 mg) and the solution was stirred at the same temperature for 30 minutes. Iodomethane (0.121 ml) was added to the solution and the mixture was stirred at ambient temperature for 3 hours. The mixture was diluted with ethyl acetate (50 ml) and the solution was washed with water and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo to give a crude oil. The crude product was purified by silica gel column chromatography (1% methanol in chloroform) to give 4-(2-dimethylamino-4-methyl)-phenoxyethyl-N-[2-(5-ethoxycarbonylpent-1-yl)oxy]phenyl-N-methylbenzamide (632 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.26 (3H, t, J=7.5Hz), 1.42-1.55 (2H, m), 1.63-1.74 (2H, m), 1.76-1.87 (2H, m), 2.20 (3H, s), 2.23 (3H, s), 2.33 (2H, t, J=7.5Hz), 2.72 (6H, s), 3.30 (3H, s), 3.76-3.97 (2H, m), 4.12 (2H, q, J=7.5Hz), 5.02 (2H, s), 6.52-6.60 (3H, m), 6.70 (1H, d, J=7Hz), 6.80-6.88 (2H, m), 7.20 (2H, d, J=8Hz), 7.31 (2H, d, J=8Hz)

30      Example 70

The following compound was obtained by using 3-methoxy-4-[2-[3-(tert-butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-N-[2-(4-aminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide as a starting compound according to a similar manner to that of Example 14.

35

- 232 -

3-Methoxy-4-[2-[3-(tert-butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-N-[2-(4-acetylaminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide

5 NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.65-1.82 (4H, m), 1.76 (3H, s), 2.05 (3H, s), 2.07-2.21 (2H, m), 2.26 (3H, s), 3.22-3.38 (2H, m), 3.38 (3H, s), 3.77 (3H, s), 3.77-3.96 (2H, m), 4.24 (2H, t, J=7.5Hz), 6.53-6.71 (2H, m), 6.93-7.14 (5H, m), 7.25 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.43 (7H, d, J=7Hz)

10

#### Example 71

To a mixture of 3-methoxy-4-[2-[3-(tert-butoxycarbonyl)-aminoprop-1-yl]oxybenzoyl]amino-N-[2-(4-aminobut-1-yl)oxy-4-methyl]phenyl-N-methylbenzamide (365 mg) and N-(tert-butoxycarbonyl)glycine (111 mg) in N,N-dimethylformamide (15 ml) were added N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride (132 mg) and hydroxybenzotriazole (93.2 mg) and the mixture was stirred at ambient temperature overnight. The solution was diluted with ethyl acetate (30 ml) and the solution was washed successively with saturated aqueous sodium hydrogen carbonate, water and brine. The organic phase was dried over magnesium sulfate and the solvent was evaporated in vacuo to give an amorphous. The crude product was purified by silica gel column chromatography (1% methanol in chloroform) to give 3-methoxy-4-[2-[3-tert-butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-N-[2-4-(tert-butoxycarbonylamino)acetylaminobut-1-yl]oxy-4-methyl]phenyl-N-methylbenzamide (320 mg).

15

20

25

30 NMR (CDCl<sub>3</sub>, δ) : 1.39 (9H, s), 1.42 (9H, s), 1.58-1.70 (2H, m), 1.70-1.80 (2H, m), 2.05-2.17 (2H, m), 2.27 (3H, s), 3.20-3.34 (4H, m), 3.30 (3H, s), 3.70-3.95 (4H, m), 3.74 (3H, s), 4.22 (2H, t, J=7.5Hz), 6.56-6.68 (2H, m), 6.88-7.11 (5H, m), 7.45 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz), 8.28 (1H, d, J=7Hz)

35

- 233 -

Example 72

The following compounds were obtained according to a similar manner to that of Example 71.

- 5           1) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-  
              amino-3-methoxy-N-[2-[4-[3-(tert-butoxycarbonyl)-  
              aminopropionylamino]but-1-yl]oxy-4-methylphenyl]-N-  
              methylbenzamide  
              NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.41 (9H, s), 1.60-1.82  
10               (4H, m), 2.10-2.19 (2H, m), 2.29 (3H, s), 2.48 (2H,  
                  br), 3.25-3.42 (6H, m), 3.32 (3H, s), 3.79 (3H, s),  
                  3.80-3.97 (2H, m), 4.25 (2H, t, J=5Hz), 6.59 (1H,  
                  s), 6.67 (1H, d, J=8Hz), 6.94-7.11 (5H, m), 7.45  
                  (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.39 (1H,  
15               d, J=8Hz)
- 2) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-  
              amino-3-methoxy-N-[2-[4-[[1-(tert-butoxycarbonyl)-  
              piperidin-4-yl]carbonylamino]but-1-yl]oxy-4-  
20               methylphenyl]-4-methylbenzamide  
              NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.44 (9H, s), 1.60-1.81  
                  (8H, m), 2.08-2.18 (2H, m), 2.29 (3H, s), 2.70 (1H,  
                  br), 3.30 (2H, q, J=5Hz), 3.32 (3H, s), 3.76 (3H,  
                  s), 3.76-4.15 (6H, m), 4.22 (2H, t, J=5Hz), 6.59  
25               (1H, s), 6.65 (1H, d, J=8Hz), 6.94-7.10 (6H, m),  
                  7.44 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.39  
                  (1H, d, J=8Hz)

Example 73

- 30           To an ice-cooled mixture of 4-[2-[3-(tert-  
              butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-  
              [2-(4-aminobut-1-yl)oxy-4-methylphenyl]-N-methylbenzamide  
              (430 mg) and triethylamine (68 mg) in dichloromethane (10 ml)  
              was added phenyl chlorocarbonate (106 mg) dropwise and the  
35               solution was stirred at the same temperature for 30 minutes.

- 234 -

The resulting mixture was diluted with dichloromethane (10 ml) and the solution was washed successively with 1N hydrochloric acid saturated aqueous sodium hydrogen carbonate and brine. The solvent was dried over magnesium sulfate and removed under reduced pressure to give 4-[2-[3-(tert-butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-[(2-(4-phenoxy-carbonylamino)but-1-yl)oxy-4-methylphenyl]-N-methylbenzamide (471 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.60-1.90 (4H, m), 2.08-2.17 (2H, m), 2.29 (3H, s), 3.27 (2H, q, J=5Hz), 3.31 (2H, t, J=5Hz), 3.36 (3H, s), 3.78 (3H, s), 3.82-4.00 (2H, m), 4.21 (2H, t, J=5Hz), 4.73 (1H, br), 5.38 (1H, br), 6.61-6.68 (2H, m), 6.91-6.99 (4H, m), 7.06-7.20 (5H, m), 7.30-7.38 (2H, m), 7.42 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

#### Example 74

A mixture of 4-[2-[3-(tert-butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-[2-(4-aminobut-1-yl)oxy-4-methylphenyl]-N-methylbenzamide (120 mg) and 3-(dimethylamino)prop-1-yl phenyl carbonate (127 mg) in N,N-dimethylformamide (5 ml) was stirred at 50°C for 8 hours. The reaction mixture was diluted with ethyl acetate (15 ml) and the solution was washed successively with saturated aqueous sodium bicarbonate solution and brine. The solution was dried over potassium carbonate. The solvent was evaporated and the residue was purified on silica gel column chromatography (SiO<sub>2</sub> 20 g, 3-15% methanol in chloroform) to give 4-[2-[3-(tert-butoxycarbonylamino)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-[2-(3-dimethylaminoprop-1-yl)oxycarbonylamino]but-1-yl]oxy-4-methylphenyl]-N-methylbenzamide (64 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.62-1.87 (6H, m), 2.05-2.18 (2H, m), 2.28 (3H, s), 2.30 (6H, s), 2.44 (2H, t, J=5Hz), 3.20-3.32 (4H, m), 3.32 (3H, s),



- 235 -

3.78 (3H, s), 3.80-4.00 (2H, m), 4.12 (2H, t, J=5Hz), 4.24 (2H, t, J=5Hz), 6.59-6.64 (2H, m), 6.88-7.12 (5H, m), 7.44 (1H, dd, J=2, 8Hz), 8.21 (1H, d, J=8Hz), 8.40 (1H, br)

5

Example 75

To a solution of 4-[2-[(3-tert-butoxycarbonylamino-  
prop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-  
oxopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]-  
10 benzamide (192 mg) in methanol (5 ml) was added sodium  
borohydride (19 mg) at ambient temperature and the mixture  
was stirred at the same temperature for 1 hour. The reaction  
was quenched with 0.5N hydrochloric acid (10 ml) and the  
mixture was extracted with chloroform (15 ml x 3). The  
15 organic layer was washed with aqueous sodium hydrogen  
carbonate and brine, and the solution was dried over  
magnesium sulfate. The solvent was evaporated in vacuo to  
give 4-[2-[(3-tert-butoxycarbonylamino-  
prop-1-yl)oxy]benzoyl]-  
20 amino-3-methoxy-N-methyl-N-[2-[5-(4-hydroxypiperidin-1-  
yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide (199 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.39 (9H, s), 1.41-1.99 (10H, m),  
2.05-2.20 (2H, m), 2.27 (3H, s), 2.30-2.51 (2H, m),  
3.01-3.22 (2H, m), 3.30 (3H, s), 3.65-4.14 (7H, m),  
3.76 (3H, s), 4.22 (2H, t, J=5Hz), 6.52-6.67 (2H,  
25 m), 6.78-7.10 (5H, m), 7.38-7.47 (1H, m), 8.19 (1H,  
d, J=7Hz), 8.39 (1H, d, J=7Hz)

Example 76

To a mixture of 4-[2-[(3-tert-butoxycarbonylamino-  
prop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-  
oxopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]-  
30 benzamide (250 mg), ammonium acetate (51 mg) and acetic acid  
(0.5 ml) in methanol (10 ml) was added sodium  
cyanoborohydride (21 mg) at 0°C and the mixture was stirred  
35 at ambient temperature for 12 hours. The mixture was poured

- 236 -

into ice-cooled 1N aqueous sodium hydroxide solution (15 ml) and the solution was extracted with chloroform (15 ml x 3). The organic layer was washed with brine and dried over potassium carbonate. The solvent was evaporated in vacuo and the residue was purified by silica gel column chromatography (SiO<sub>2</sub> 40 g, 5-15% methanol in chloroform) to give 4-[2-[(3-tert-butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-aminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide (91 mg).

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.41 (9H, s), 1.45-2.01 (12H, m), 2.09-2.20 (2H, m), 2.28 (3H, s), 2.23-2.45 (4H, m), 2.56-2.71 (1H, br), 2.93-3.12 (2H, m), 3.25-3.36 (2H, m), 3.32 (3H, s), 3.79 (3H, s), 3.81-4.02 (2H, m), 4.23 (2H, t, J=5Hz), 4.91-4.08 (1H, br), 6.56-6.68 (2H, m), 6.82-7.13 (5H, m), 7.45 (2H, d, J=8Hz), 8.20 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

#### Example 77

The following compound was obtained according to a similar manner to that of Example 76.

4-[2-(3-tert-Butoxycarbonylaminoprop-1-yl)oxybenzoyl]-amino-3-methoxy-N-methyl-N-[2-[6-(4-methylpiperazin-1-yl)hex-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.39 (9H, s), 1.45-1.84 (8H, m), 2.09-2.22 (2H, m), 2.27 (3H, s), 2.28 (3H, s), 2.32-2.59 (8H, m), 3.32 (1H, q, J=5Hz), 3.34 (3H, s), 3.80 (3H, s), 3.82-4.01 (2H, m), 4.28 (2H, t, J=5Hz), 6.56-6.65 (2H, m), 6.82-7.12 (6H, m), 7.43-7.50 (1H, m), 8.20 (1H, d, J=8Hz), 8.38 (1H, d, J=8Hz)

#### Example 78

To a mixture of 4-[2-(3-tert-butoxycarbonylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-carboxypent-1-

- 237 -

yl]oxy-4-methylphenyl]benzamide (250 mg), 2-dimethylamino-ethanol (99 mg) and 4-dimethylaminopyridine (36 mg) in dichloromethane (10 ml) was added N-ethyl-N'-(3-dimethylaminoprop-1-yl)carbodiimide hydrochloride (71 mg) at 0°C and stirred at the same temperature for 7 hours. The mixture was diluted with chloroform (20 ml) and the solution was washed with water (20 ml x 2) and brine. The solution was dried over magnesium sulfate and the solvent was removed under reduced pressure. This residue was purified by silica gel column chromatography (SiO<sub>2</sub> 30 g, 1-10% methanol in chloroform) to give 4-[2-(3-tert-butoxycarbonylamino-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(2-dimethylaminoeth-1-yl)oxycarbonylpent-1-yl]oxy-4-methylphenyl]-benzamide (238 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.45-1.57 (2H, m), 1.65-1.90 (4H, m), 2.10-2.21 (2H, m), 2.28 (9H, s), 2.39 (2H, t, J=5Hz), 2.55 (2H, t, J=5Hz), 3.30 (2H, t, J=5Hz), 3.32 (3H, s), 3.79 (3H, s), 3.82-4.00 (2H, m), 4.18 (2H, t, J=5Hz), 4.24 (2H, t, J=5Hz), 4.75-4.86 (1H, br), 6.54-6.67 (2H, m), 6.81-7.11 (5H, m), 7.41-7.49 (1H, m), 8.20 (1H, d, J=8Hz), 8.49 (1H, d, J=8Hz).

#### Example 79

To a solution of 4-[2-(3-tert-butoxycarbonylamino-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-(5-ethoxycarbonylpent-1-yl)oxy-4-methylphenyl]benzamide (400 mg) in tetrahydrofuran (5 ml) was added lithium aluminum hydride (12 mg) at -23°C and the mixture was stirred at 0°C for 3 hours. The reaction was quenched with slow addition of 0.5N hydrochloric acid (15 ml) and the solution was stirred at ambient temperature for 20 minutes. The solution was extracted with chloroform (15 ml x 3) and the organic layer was washed with aqueous saturated sodium bicarbonate solution and brine. The solution was dried over magnesium sulfate and

- 238 -

the solvent was removed under reduced pressure to give 4-[2-(3-tert-butoxycarbonylamino-1-propyloxybenzoyl)amino-3-methoxy-N-methyl-N-[2-(6-hydroxyhex-1-yl)oxy-4-methylphenyl]benzamide (456 mg).

5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 1.45-2.20 (10H, m),  
2.27 (3H, s), 3.30 (2H, q, J=5Hz), 3.32 (3H, s),  
3.64 (2H, t, J=5Hz), 3.78 (3H, s), 3.81-4.02 (2H,  
m), 4.23 (2H, t, J=5Hz), 6.57-6.63 (2H, m), 6.84-  
10 7.13 (6H, m), 7.41-7.49 (1H, m), 8.20 (1H, d,  
J=7Hz) 8.41 (1H, d, J=7Hz)

#### Example 80

To a solution of oxalyl chloride (95 mg) in dichloromethane (10 ml) was added dimethyl sulfoxide (117 mg) dropwise at -78°C. The mixture was warmed to -15°C and a solution of 4-[2-(3-tert-butoxycarbonylamino-1-propyloxybenzoyl)amino-3-methoxy-N-methyl-N-[2-(6-hydroxyhex-1-yl)oxy-4-methylphenyl]benzamide (450 mg) in dichloromethane (10 ml) was added thereto. After being stirred at the same temperature for 10 minutes, to the reaction mixture was added triethylamine (343 mg) and stirred at the same temperature for 5 minutes. The resulting solution was warmed to ambient temperature and poured into water. The mixture was extracted with chloroform (15 ml x 3) and the organic layer was washed with brine. The solution was dried over magnesium sulfate and the solvent was evaporated to give 4-[2-(3-tert-butoxycarbonylamino-1-propyloxybenzoyl)amino-3-methoxy-N-methyl-N-[2-(5-formylpent-1-yl)oxy-4-methylphenyl]benzamide (546 mg)

30 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 1.50-1.91 (6H, m),  
2.11-2.23 (2H, m), 2.27 (3H, s), 2.50 (2H, t,  
J=5Hz), 3.31 (1H, q, J=5Hz), 3.34 (3H, s), 3.79  
(3H, s), 3.85-4.00 (2H, m), 4.27 (2H, t, J=5Hz),  
6.60-6.68 (2H, m), 6.81-7.12 (6H, m), 7.42-7.51  
35 (1H, m), 8.21 (1H, d, J=7Hz), 8.41 (1H, d, J=7Hz),

- 239 -

9.89 (1H, s)

Example 81

To a solution of 4-[2-[3-(tert-butoxycarbonyl)aminoprop-1-yl]oxybenzoyl]amino-3-methoxy-N-[2-(4-cyanophenylmethyl)-oxy-4-methylphenyl]-N-methylbenzamide (360 mg) in xylene (8 ml) was added trimethyltin azide (218 mg) and the solution was stirred at 120°C for 3 days. The solution was cooled to ambient temperature and 12N hydrochloric acid (10 ml) was added to the solution to decompose tin salt of the tetrazole compound and the excess reagent. Then the solution was adjusted to pH 7 with saturated aqueous sodium hydroxide at 0°C, and the solution was extracted with ethyl acetate (50 ml x 3). The organic layer was washed with brine, and dried over magnesium sulfate. The solvent was evaporated to give a crude product. The crude product was purified by silica gel column chromatography (SiO<sub>2</sub> 30 g, 2-25% methanol in chloroform) to give 4-[2-(3-aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-(tetrazol-5-yl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide (227 mg).

NMR (CDCl<sub>3</sub>, δ) : 2.15 (3H, br s), 2.14-2.26 (2H, m), 3.17 (2H, q, J=5Hz), 3.40 (3H, s), 3.57 (3H, s), 4.20 (2H, t, J=5Hz), 4.95 (1H, d, J=12Hz), 5.22 (1H, d, J=12Hz), 6.55-6.64 (2H, m), 6.80 (1H, s), 6.92-7.08 (6H, m), 7.23 (1H, br), 7.43 (1H, dd, J=2, 8Hz), 7.76 (2H, d, J=8Hz), 8.20 (1H, d, J=8Hz)

Example 82

A mixture of 4-[2-(3-aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide (275 mg) and O-methylisourea (44 mg) in ethanol (5 ml) was refluxed for 3 days. The solvent was evaporated in vacuo and the residue was purified on basic silica gel column chromatography (SiO<sub>2</sub> 17 g, 1-80% methanol in chloroform) to

- 240 -

give 4-[2-(3-guanidinoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide (53 mg).

5 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40-1.97 (6H, m), 2.06-2.20 (2H, m),  
2.27 (6H, s), 2.28 (3H, s), 2.29-2.41 (4H, m), 2.50  
(1H, br), 3.04 (2H, br), 3.30 (3H, s), 3.42 (2H,  
br), 3.76 (3H, s), 3.78 (2H, br), 3.82-4.01 (2H,  
m), 4.25 (2H, br), 6.55-6.68 (2H, m), 6.81-7.09  
10 (5H, m), 7.28 (1H, s), 7.42 (1H, dd, J=2, 8Hz),  
7.99 (1H, d, J=8Hz), 8.29 (1H, br)

#### Example 83

The following compound was obtained according to a  
similar manner to that of Example 6.

15 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-  
amino-3-methoxy-N-[2-(4-aminobut-1-yl)oxy-4-methylphenyl]-N-  
methylbenzamide

20 NMR (CDCl<sub>3</sub>,  $\delta$ ) : 1.40 (9H, s), 1.81-1.99 (4H, m),  
2.05-2.14 (2H, m), 2.24 (3H, s), 3.08 (2H, br),  
3.29 (2H, br), 3.30 (3H, s), 3.70 (3H, s), 3.76-  
3.96 (2H, m), 4.14 (2H, t, J=5Hz), 5.07 (1H, br),  
6.54-6.61 (2H, m), 6.85-7.04 (4H, m), 7.25 (1H, s),  
25 7.37 (1H, dd, J=2, 8Hz), 8.14 (1H, d, J=8Hz), 8.38  
(1H, d, J=8Hz)

#### Example 84

30 To a solution of 4-[2-[3-(tert-butoxycarbonylamino)prop-  
1-yloxy]benzoylamino]-3-methoxy-N-[2-[4-(phenoxycarbonyl-  
amino)but-1-yl]-4-methylphenyl]-N-methylbenzamide (200 mg) in  
N,N-dimethylformamide (5 ml) was added 1-methylpiperazine (88  
μl) and the solution was stirred at 80°C for 7 hours. The  
solution was diluted with ethyl acetate (15 ml) and washed  
successively with water (20 ml x 4) and brine. The solvent  
35 was dried over magnesium sulfate and removed under reduced

- 241 -

pressure. The crude product was purified on silica gel column chromatography (SiO<sub>2</sub> 25 g, chloroform-methanol 2-10%) to give pure 4-[2-[3-(tert-butoxycarbonylamino)prop-1-yl]oxybenzoyl]amino-3-methoxy-N-[2-[4-[(4-methylpiperazin-1-yl)carbonylamino]but-1-yl]oxy-4-methylphenyl]-N-methylbenzamide (124 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.60-1.81 (4H, m), 2.13-2.22 (2H, m), 2.29 (6H, s), 2.39 (4H, br), 3.79 (3H, s), 3.25-3.51 (8H, m), 3.32 (3H, s), 3.75-3.99 (2H, m), 4.26 (2H, t, J=5Hz), 6.57-6.71 (2H, m), 6.92-7.18 (6H, m), 7.48 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

#### Example 85

The following compounds were obtained according to a similar manner to that of Example 84.

- 1) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-[2-[4-[(4-dimethylaminopiperidin-1-yl)carbonylamino]but-1-yl]oxy-4-methylphenyl]-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.65-1.90 (4H, m), 2.29 (3H, s), 2.30 (6H, s), 2.77 (1H, t, J=11Hz), 3.29 (2H, q, J=5Hz), 3.32 (3H, s), 3.78 (3H, s), 3.85-4.11 (6H, m), 4.25 (2H, t, J=5Hz), 6.55-6.70 (2H, m), 6.92-7.13 (5H, m), 7.45 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

- 2) 4-[2-[3-(tert-Butoxycarbonylamino)prop-1-yl]oxybenzoyl]-amino-3-methoxy-N-[2-(4-ureidobut-1-yl)oxy-4-methylphenyl]-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.45-1.80 (4H, m), 2.01-2.11 (2H, m), 2.27 (3H, s), 3.22-3.31 (2H, m), 3.30 (3H, s), 3.65-3.77 (2H, m), 3.71 (3H, s), 4.22 (2H, t, J=5Hz), 5.16 (2H, br), 6.48 (1H, s), 6.71

- 242 -

(1H, d, J=8Hz), 6.90-7.15 (5H, m), 7.41 (1H, dd, J=2, 8Hz), 8.11 (1H, d, J=8Hz), 8.35 (1H, d, J=8Hz)

Example 86

5 To a solution of 4-[2-[3-(tert-butoxycarbonylamino)prop-1-yloxy]benzoylamino]-3-methoxy-N-[2-[4-(phenoxycarbonyl-amino)but-1-yl]-4-methylphenyl]-N-methylbenzamide (150 mg) in N,N-dimethylformamide (5 ml) was added dimethylamine hydrochloride (40 mg) and the mixture was stirred at 80°C for 10 7 hours. The mixture was cooled to ambient temperature and diluted with ethyl acetate (15 ml). The solution was washed with water (15 ml x 5) and brine, and dried over magnesium sulfate. The solvent was removed in vacuo and the residue was purified on silica gel column chromatography (SiO<sub>2</sub> 20 g, 15 chloroform-methanol 1-5%) to give 4-[2-[3-(tert-butoxycarbonylamino)prop-1-yloxy]benzoylamino]-3-methoxy-N-[2-[4-(N,N-dimethylureido)but-1-yloxy]-4-methylphenyl]-N-methylbenzamide (115 mg).

20 NMR (CDCl<sub>3</sub>, δ) : 1.40 (9H, s), 1.60-1.87 (4H, m), 2.06-2.18 (2H, m), 2.28 (3H, s), 2.90 (6H, s), 3.30 (2H, q, J=5Hz), 3.34 (3H, s), 3.79 (3H, s), 3.85-4.02 (2H, m), 4.23 (2H, t, J=5Hz), 6.57-6.64 (2H, m), 6.90-7.10 (5H, m), 7.44 (1H, dd, J=2, 8Hz), 8.20 (1H, d, J=8Hz), 8.41 (1H, d, J=8Hz)

Example 87

30 To a solution of 4-[2-[(3-tert-butoxycarbonylamino)prop-1-yl]oxy]benzoyl]amino-3-carboxymethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-benzamide (128 mg) in methanol (5 ml) was added dropwise trimethylsilyldiazomethane (5 ml, 2.0M n-hexane solution) and stirred at ambient temperature for 30 minutes. The solution was concentrated in vacuo and the residue was purified by preparative thin layer silica gel chromatography 35 (chloroform:methanol:28% aqueous ammonia solution, 50:5:1) to



- 243 -

give 4-[2-[(3-tert-butoxycarbonylamino)prop-1-yl]oxy]benzoyl]-amino-3-methoxycarbonylmethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]-4-methylphenyl]-benzamide (85 mg).

5 NMR (CDCl<sub>3</sub>, δ) : 1.39 (9H, s), 1.45-1.37 (8H, m),  
2.00-2.10 (2H, m), 2.26 (3H, s), 2.29 (3H, s),  
2.30-2.42 (6H, m), 3.18-3.27 (2H, m), 3.30 (3H, s),  
3.45-3.51 (2H, m), 3.63 (2H, br), 3.79 (3H, s),  
3.87-3.96 (2H, m), 4.22-4.29 (2H, m), 4.54 (2H, s),  
10 6.53-6.13 (2H, m), 6.77-6.85 (1H, m), 6.89 (1H,  
br), 6.92-7.02 (2H, m), 7.02-7.10 (1H, m), 7.43-  
7.47 (1H, m), 8.14-8.19 (1H, m), 8.40-8.45 (1H, m)

#### Example 88

15 The following compound was obtained according to a similar manner to that of Example 8.

4-[2-[(3-tert-Butoxycarbonylamino)prop-1-yl]oxy]benzoyl]-amino-3-dimethylaminocarbonylmethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]-4-methylphenyl]-benzamide

20 NMR (CDCl<sub>3</sub>, δ) : 1.39 (9H, s), 1.48-1.58 (2H, m),  
1.63-1.88 (6H, m), 1.97-2.09 (2H, m), 2.28 (3H, s),  
2.30 (3H, s), 2.31-2.42 (6H, m), 2.99 (3H, s), 3.02  
25 (3H, s), 3.17-3.27 (2H, m), 3.32 (3H, s), 3.50 (2H,  
br), 3.63 (2H, br), 3.83-3.97 (2H, m), 4.22-4.29  
(2H, m), 4.67 (2H, s), 6.53-6.63 (2H, m), 6.80-6.90  
(2H, m), 6.96-7.09 (3H, m), 7.93 (1H, t, J=6Hz),  
8.14 (1H, d, J=6Hz), 8.38 (1H, d, J=7Hz)

#### Example 89

To a solution of 4-[2-[(3-tert-butoxycarbonylamino)prop-1-yl]oxy]benzoyl]amino-3-ethoxycarbonylmethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]-4-methylphenyl]benzamide (102 mg) in 7.5N ammonia in methanol

- 244 -

(5 ml) was stirred at ambient temperature for 24 hours. The solution was concentrated in vacuo to give 4-[2-[(3-tert-butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-3-aminocarbonylmethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide (92 mg).

NMR (CDCl<sub>3</sub>, δ) : 1.37 (9H, s), 1.48-1.60 (2H, m), 1.60-1.75 (4H, m), 1.75-1.88 (2H, m), 1.97-2.08 (2H, m), 2.27 (3H, s), 2.28 (3H, s), 2.30-2.41 (6H, m), 3.17-3.27 (2H, m), 3.30 (3H, s), 3.47 (3H, s), 3.52-3.62 (2H, m), 3.90-3.97 (2H, m), 4.16-4.29 (4H, m), 5.85 (1H, br), 6.57 (1H, d, J=7Hz), 6.67 (1H, s), 6.75-6.90 (2H, m), 7.00 (1H, d, J=7Hz), 7.07-7.17 (2H, m), 8.00 (1H, s), 8.18-8.21 (1H, m), 8.25 (1H, d, J=7Hz)

#### Example 90

The following compound was obtained according to a similar manner to that of Example 89.

4-[2-[(3-tert-Butoxycarbonylaminoprop-1-yl)oxy]benzoyl]-amino-3-methylaminocarbonylmethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-benzamide

NMR (CDCl<sub>3</sub>, δ) : 3.37 (9H, s), 1.45-1.77 (6H, m), 1.77-1.88 (2H, m), 1.96-2.08 (2H, m), 2.28 (3H, s), 2.29 (3H, s), 2.29-2.40 (6H, m), 2.82-2.83 (3H, s), 3.18-3.27 (2H, m), 3.30 (3H, s), 3.43-3.50 (3H, m), 3.57 (2H, br), 3.90-3.97 (2H, m), 4.18-4.30 (3H, m), 6.57 (1H, d, J=6Hz), 6.65 (1H, s), 6.76-6.83 (2H, m), 7.00 (1H, d, J=7Hz), 7.06-7.15 (2H, m), 7.45 (1H, t, J=7Hz), 8.16-8.22 (2H, m)

#### Example 91

The following compound was obtained according to similar manners to those of Examples 8 and 16.

- 245 -

4-(2-Aminobenzoyl)amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

5 NMR (DMSO-d<sub>6</sub>, δ) : 1.35-1.66 (4H, m), 1.66-1.82 (2H, m), 2.22 (3H, s), 2.40 (2H, t, J=7Hz), 2.73 (3H, s), 2.77-3.11 (3H, m), 3.17 (3H, s), 3.28-3.56 (3H, m), 3.76-4.17 (3H, m), 4.35-4.52 (1H, m), 6.63 (1H, d, J=9Hz), 6.79 (1H, s), 6.91 (1H, dd, J=9, 9Hz), 6.98-7.11 (2H, m), 7.22 (2H, d, J=9Hz), 7.36 (1H, dd, J=9, 9Hz), 7.54 (2H, d, J=9Hz), 7.69 (1H, d, J=9Hz)

#### Example 92

15 The following compounds were obtained according to similar manners to those of Examples 6 and 16.

1) 4-[2-[(3-Aminoprop-1-yl)amino]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yloxy]phenyl]benzamide trihydrochloride

20 NMR (DMSO-d<sub>6</sub>, δ) : 1.36-1.65 (4H, m), 1.66-1.92 (4H, m), 2.23 (3H, s), 2.38 (2H, t, J=7Hz), 2.68-2.77 (3H, m), 2.77-3.12 (4H, m), 3.18 (3H, s), 3.22 (2H, t, J=7Hz), 3.28-3.56 (3H, m), 3.63 (3H, s), 3.75-4.32 (4H, m), 4.42 (1H, m), 6.58-6.69 (2H, m), 6.78 (1H, d, J=8Hz), 6.83 (1H, s), 6.86-6.96 (2H, m), 7.03 (1H, d, J=8Hz), 7.34 (1H, dd, J=8, 8Hz), 7.61 (1H, d, J=8Hz), 7.67 (1H, d, J=8Hz), 7.91-8.17 (3H, m), 9.23 (1H, s)

30 2) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

35 NMR (DMSO-d<sub>6</sub>, δ) : 1.34-1.66 (4H, m), 1.66-1.83 (2H, m), 2.04-2.24 (2H, m), 2.32-2.46 (2H, m), 2.74 (3H, s), 2.79-3.12 (4H, m), 3.22 (3H, s), 3.29-3.58 (3H,

- 246 -

m), 3.63-4.19 (7H, m), 4.28-4.52 (3H, m), 6.80-7.08 (4H, m), 7.08-7.36 (4H, m), 7.58 (1H, dd, J=9, 9Hz), 8.02 (1H, d, J=9Hz), 8.13 (2H, br s), 8.28 (1H, d, J=9Hz)

5

- 3) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)-carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

10

NMR (DMSO-d<sub>6</sub>, δ) : 1.28-1.82 (8H, m), 1.90-2.51 (11H, m), 2.64 (6H, s), 2.74-3.06 (3H, m), 3.18 (3H, s), 3.22-4.08 (6H, m), 4.29-4.41 (2H, m), 4.51 (1H, m), 6.64 (1H, d, J=8Hz), 6.75-7.20 (5H, m), 7.27 (1H, d, J=8Hz), 7.58 (1H, m), 7.94-8.32 (5H, m)

15

- 4) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-2-chloro-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonyl]pent-1-yloxy]phenyl]benzamide dihydrochloride

20

NMR (DMSO-d<sub>6</sub>, δ) : 1.39-1.68 (4H, m), 1.69-1.90 (2H, m), 1.92-2.12 (2H, m), 2.31-2.50 (2H, m), 2.73 (3H, br s), 2.79-3.10 (4H, m), 3.17-3.61 (7H, m), 3.92-4.26 (5H, m), 4.42 (1H, m), 6.77 (1H, m), 6.92-7.23 (6H, m), 7.34-7.58 (3H, m), 7.81 (1H, s), 7.90-8.14 (3H, m)

25

- 5) 4-[2-(3-Aminoprop-1-yl)oxy-5-methylbenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

30

NMR (CDCl<sub>3</sub>, δ) : 1.50-1.93 (8H, m), 2.28 (3H, s), 2.28-2.36 (2H, m), 2.31 (3H, s), 2.79 (3H, s), 3.09-3.20 (2H, m), 3.29 (3H, s), 3.80 (3H, s), 3.85-4.04 (2H, m), 4.18-4.28 (2H, m), 6.57-6.66 (2H, m), 6.80-6.95 (4H, m), 7.20-7.25 (1H, m), 7.72 (1H, br), 8.51 (1H, br)

35

- 247 -

- 6) 4-[2-(3-Aminoprop-1-yl)oxy-4-chlorobenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

5 NMR (CDCl<sub>3</sub>, δ) : 1.45-1.86 (8H, m), 2.23 (3H, s),  
2.29-2.43 (2H, m), 2.78 (3H, s), 3.05-3.16 (2H, m),  
3.23 (3H, s), 3.78 (3H, s), 3.82-4.03 (2H, m),  
4.18-4.32 (2H, m), 6.54-6.64 (2H, m), 6.78-7.08  
(4H, m), 7.94 (1H, d, J=8Hz), 8.58 (1H, br)

10

- 7) 4-[2-(3-Aminoprop-1-yl)oxy-4-methoxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)-carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

15 NMR (CDCl<sub>3</sub>, δ) : 1.40-1.89 (6H, m), 2.28 (3H, s),  
2.30-2.61 (6H, m), 2.70-3.04 (4H, m), 3.08-3.25  
(2H, m), 3.28 (3H, s), 3.80 (6H, s), 3.82-4.08 (2H,  
m), 4.26 (2H, br), 6.49-6.66 (4H, m), 6.78-7.00  
(3H, m), 7.93-8.02 (1H, m), 8.30 (1H, br), 8.52  
20 (2H, br)

- 8) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

25 NMR (DMSO-d<sub>6</sub>, δ) : 1.42 (2H, br), 1.53 (2H, br), 1.74  
(2H, br), 2.03 (2H, br), 2.13-2.20 (2H, m), 2.30-  
2.38 (2H, m), 2.66 (3H, s), 2.67 (3H, s), 2.94 (4H,  
br), 3.20 (3H, s), 3.28-3.40 (2H, m), 3.73 (3H, s),  
3.82-4.08 (4H, m), 4.33-4.40 (2H, m), 4.47-4.57  
30 (1H, m), 6.82-7.00 (4H, m), 7.10-7.29 (4H, m),  
7.53-7.60 (1H, m), 8.00 (1H, d, J=7Hz), 8.22-8.30  
(1H, m)

- 9) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methyl-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-

35

- 248 -

yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.40-1.53 (2H, m), 1.53-1.65 (2H, m), 1.66-1.82 (2H, m), 2.01-2.13 (2H, m), 2.18 (3H, s), 2.23 (3H, s), 2.36-2.46 (2H, m), 2.73-2.74 (3H, s), 2.78-3.08 (6H, m), 3.18 (3H, s), 4.27 (2H, br), 4.40-4.50 (1H, m), 6.65 (1H, d, J=6Hz), 6.82 (1H, s), 6.98-7.13 (3H, m), 7.17-7.30 (2H, m), 7.45-7.57 (2H, m), 7.22 (1H, d, J=6Hz), 9.67 (1H, s)

- 10) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-ethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride
- NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.23 (3H, t, J=6Hz), 1.38-1.50 (2H, m), 1.50-1.65 (2H, m), 1.65-1.82 (2H, m), 2.05-2.17 (2H, m), 2.21 (3H, s), 2.32-2.43 (2H, m), 2.70-2.73 (3H, m), 2.80-3.08 (7H, m), 3.18 (3H, s), 3.22-3.55 (6H, m), 3.92-4.15 (2H, m), 4.32-4.48 (4H, m), 6.63 (1H, d, J=7Hz), 6.83 (1H, s), 6.89-6.92 (2H, m), 7.02 (1H, d, J=7Hz), 7.13 (1H, t, J=6Hz), 7.29 (1H, d, J=7Hz), 7.58 (1H, t, J=7Hz), 7.99 (1H, d, J=7Hz), 8.18-8.27 (1H, m)

### Example 93

The following compounds were obtained according to similar manners to those of Examples 1 and 16.

- 1) 4-[2-(Dimethylamino)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride
- NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.36-1.65 (4H, m), 1.67-1.82 (2H, m), 2.22 (3H, s), 2.38 (2H, t, J=7Hz), 2.64-3.14 (12H, m), 3.18 (3H, s), 3.28-3.42 (2H, m), 3.50 (1H, m), 3.73 (3H, s), 3.79-4.14 (3H, m), 4.42 (1H, m), 6.64 (1H, d, J=8Hz), 6.82 (1H, s), 6.83-6.97 (2H, m), 7.02 (1H, d, J=8Hz), 7.35 (1H, m), 7.52-

- 249 -

7.67 (2H, m), 8.07 (1H, d, J=8Hz), 8.14 (1H, m)

- 2) 4-[2-(Dimethylaminosulfonyl)benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

5 NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.64 (4H, m), 1.67-1.82 (2H, m), 2.23 (3H, s), 2.38 (2H, t, J=7Hz), 2.69 (6H, s), 2.74 (3H, s), 2.80-3.12 (4H, m), 3.18 (3H, s), 3.23-3.52 (2H, m), 3.59 (3H, s), 3.81-4.16 (3H, m), 10 4.44 (1H, m), 6.66 (1H, d, J=9Hz), 6.77-6.96 (3H, m), 7.02 (1H, d, J=9Hz), 7.51 (1H, m), 7.60-7.92 (4H, m)

- 3) 3-Methoxy-4-[2-(morpholinosulfonyl)benzoyl]amino-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

15 NMR (DMSO-d<sub>6</sub>, δ) : 1.37-1.65 (4H, m), 1.66-1.83 (2H, m), 2.23 (3H, s), 2.32-2.44 (2H, m), 2.73 (3H, s), 2.81-3.10 (6H, m), 3.18 (3H, s), 3.25-3.71 (11H, m), 20 3.80-4.20 (3H, m), 4.42 (1H, m), 6.66 (1H, d, J=8Hz), 6.76-6.96 (3H, m), 7.02 (1H, d, J=8Hz), 7.53 (1H, d, J=8Hz), 7.62-7.93 (4H, m), 8.31 (1H, s)

- 25 4) 4-[2-(Isoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.39 (6H, d, J=7Hz), 1.38-1.66 (4H, m), 1.67-1.83 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=7Hz), 2.76 (3H, s), 2.82-3.11 (4H, m), 3.18 (3H, s), 3.74 (3H, s), 3.79-4.18 (5H, m), 4.36-4.52 (1H, m), 4.98 (1H, m), 6.65 (1H, d, J=8Hz), 6.73-7.17 (5H, m), 7.30 (1H, d, J=8Hz), 7.54 (1H, dd, J=8, 8Hz), 8.04 (1H, d, J=8Hz), 8.31 (1H, d, J=8Hz)

35

- 250 -

Example 94

The following compound was obtained according to similar manners to those of Examples 16 and 30.

5           4-[2-[2-[(3-Aminoprop-1-yl)oxy]phenyl]vinyl]-3-methoxy-  
N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-  
yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

10           NMR (DMSO-d<sub>6</sub>, δ) : 1.33-1.64 (4H, m), 1.64-1.83 (2H,  
m), 1.95-2.17 (2H, m), 2.22 (3H, s), 2.39 (2H, t,  
J=7Hz), 2.72 (3H, s), 2.78-3.10 (6H, m), 3.15 and  
3.16 (total 3H, s), 3.28-3.60 (2H, m), 3.64 (3H,  
s), 3.80-4.20 (5H, m), 4.42 (1H, m), 6.44-7.60  
(12H, m), 8.00-8.26 (2H, m)

15           Example 95

The following compounds were obtained according to similar manners to those of Examples 1 and 43.

20           1) 4-(2-Hydroxybenzoyl)amino-3-methoxy-N-[2-[4-(4-  
dimethylaminopiperidin-1-yl)carbonyl-4-  
methyl]phenylmethoxy]phenyl-N-methylbenzamide  
MASS (m/z) : 637 (M+1)

25           2) 4-(2-Hydroxy)benzoylamino-3-methoxy-N-methyl-N-[2-[3-(4-  
methylpiperazin-1-yl)carbonylmethoxyprop-1-yl]oxy]-  
phenylbenzamide

30           NMR (CDCl<sub>3</sub>, δ) : 2.05-2.16 (2H, m), 2.28 (3H, s),  
2.33-2.40 (4H, m), 3.35 (3H, s), 3.40-3.45 (2H, m),  
3.57-3.63 (2H, m), 3.69 (2H, t, J=7.5Hz), 3.78 (3H,  
s), 3.94-4.11 (2H, m), 4.12 (2H, s), 6.79-7.04 (7H,  
m), 7.18 (1H, t, J=7Hz), 7.42 (1H, t, J=7Hz), 7.50  
(1H, d, J=7Hz), 8.20 (1H, d, J=7Hz), 8.81 (1H, s)

35           3) 4-(2-Hydroxy)benzoyl-3-methoxy-N-[2-[(E)-5-(4-  
dimethylaminopiperidin-1-yl)carbonyl-4-penten-1-yl]oxy-



- 251 -

## 4-methylphenyl-N-methylbenzamide

NMR (CDCl<sub>3</sub>, δ) : 1.33-1.53 (2H, m), 1.84-2.05 (4H, m),  
2.27 (3H, s), 2.33 (3H, s), 2.40 (3H, s), 2.30-4.13  
(11H, m), 3.32 (3H, s), 4.67 (1H, m), 6.30 (1H, d,  
J=15Hz), 6.55-6.66 (2H, m), 6.78-7.56 (8H, m), 8.18  
(1H, m)

Example 96

The following compound was obtained according to similar  
manners to those of Examples 4, 16 and 45.

4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-(3-  
carboxyprop-1-yl)oxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-  
yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.52 (2H, m), 1.52-1.65 (2H,  
m), 1.67-1.93 (4H, m), 2.05-2.16 (2H, m), 2.01 (3H,  
s), 2.29-2.43 (5H, m), 2.73 (3H, s), 3.22-3.56 (4H,  
m), 3.82-4.14 (5H, m), 4.30-4.47 (3H, m), 8.63 (1H,  
d, J=7Hz), 8.81 (1H, s), 8.88-8.92 (2H, m), 7.03  
(1H, d, J=7Hz), 7.13 (1H, t, J=7Hz), 7.27 (1H, d,  
J=7Hz), 7.56 (1H, t, J=6Hz), 7.96 (1H, d, J=6Hz),  
8.22 (1H, d, J=7Hz)

Example 97

The following compound was obtained according to similar  
manners to those of Preparation 4 and Example 16.

4-(2-Aminobenzyloxy)-3-methoxy-N-methyl-N-[4-methyl-2-  
[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-  
benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.35-1.64 (4H, m), 1.64-1.81 (2H,  
m), 2.23 (3H, s), 2.39 (2H, t, J=7Hz), 2.75 (3H,  
s), 2.80-3.09 (2H, m), 3.16 (3H, s), 3.27-3.50 (2H,  
m), 3.57 (3H, s), 3.73-4.15 (5H, m), 4.43 (1H, m),

- 252 -

5.08 (2H, s), 6.64 (1H, d, J=8Hz), 6.76-7.42 (9H, m)

Example 98

5 The following compound was obtained according to similar manners to those of Examples 14 and 16.

4-[2-(3-Acetylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide hydrochloride

10 NMR (DMSO-d<sub>6</sub>, δ) : 1.36-1.50 (2H, m), 1.50-1.64 (2H, m), 1.67-1.84 (2H, m), 1.92-2.06 (2H, m), 2.22 (3H, s), 2.32-2.44 (2H, m), 2.50 (3H, s), 2.74 and 2.75 (total 3H, s), 2.81-3.08 (3H, m), 3.19 (3H, s), 15 3.30-3.54 (3H, m), 3.70 (3H, s), 3.79-4.16 (3H, m), 4.20-4.30 (2H, m), 6.64 (1H, d, J=8Hz), 6.81 (1H, s), 6.83-6.97 (2H, m), 7.03 (1H, d, J=8Hz), 7.12 (1H, dd, J=8, 8Hz), 7.25 (1H, d, J=8Hz), 7.51-7.61 (1H, m), 7.92-8.08 (2H, m), 8.28 (1H, d, J=8Hz)

Example 99

The following compound was obtained according to similar manners to those of Examples 15 and 26.

25 4-[2-(3-Dimethylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

30 NMR (CDCl<sub>3</sub>, δ) : 1.46-1.87 (6H, m), 2.26 (3H, s), 2.37 (2H, t, J=5Hz), 2.50 (2H, br), 2.76 (6H, s), 2.77 (6H, s), 3.02-3.30 (3H, m), 3.29 (3H, s), 3.79 (3H, s), 3.80-4.04 (2H, m), 4.33 (2H, br), 6.54-6.62 (2H, m), 6.72-7.13 (5H, m), 8.05 (1H, d, J=8Hz), 8.37 (1H, d, J=8Hz), 9.85 (1H, br)

35

- 253 -

Example 100

The following compound was obtained according to similar manners to those of Examples 8 and 45.

5           4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-(5-dimethylaminocarbonyl)pent-1-yloxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.51-2.19 (10H, m), 2.27 (3H, s),  
2.35 (2H, t, J=6Hz), 2.92 (3H, s), 3.00 (3H, s),  
10           3.32 (3H, s), 3.77 (3H, s), 3.80-4.08 (2H, m), 4.29  
(2H, t, J=4Hz), 6.55-6.76 (2H, m), 6.83-7.20 (5H,  
m), 7.46 (1H, br), 8.21 (1H, d, J=8Hz), 8.40 (1H,  
d, J=8Hz)

15           Example 101

The following compound was obtained according to similar manners to those of Examples 16 and 41.

20           4-(2-Aminobenzoyl)amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]-benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.66 (4H, m), 1.68-1.83 (2H,  
m), 2.24 (3H, s), 2.34-2.44 (2H, m), 2.76 (3H, s),  
2.80-3.09 (3H, m), 3.19 (3H, s), 3.30-3.53 (3H, m),  
25           3.64 (3H, s), 3.80-4.51 (4H, m), 6.60-6.76 (2H, m),  
6.79-6.97 (4H, m), 7.05 (1H, d, J=9Hz), 7.26 (1H,  
dd, J=9, 9Hz), 7.58-7.72 (2H, m), 9.19 (1H, br s)

30           Example 102

To a solution of 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (7.35 g) in ethanol (230 ml) was added 0.5M sulfuric acid in ethanol (22.3 ml) at 80°C. The mixture was stirred for 24 hours at  
35           ambient temperature. The precipitate was filtered through a

- 254 -

glass funnel followed by rinsing with ethanol. The resulting white, crystalline solid was dried over air for 7 days to give 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide sulfate (5.2 g).

NMR (DMSO-d<sub>6</sub>, δ) : 1.35-1.63 (4H, m), 1.65-1.81 (2H, m), 2.04-2.40 (14H, m), 2.96 (2H, t, J=7Hz), 3.03-4.06 (12H, m), 4.35 (2H, t, J=7Hz), 6.64 (1H, d, J=8Hz), 6.83 (1H, s), 6.89 (1H, d, J=8Hz), 6.98 (1H, s), 7.02 (1H, d, J=8Hz), 7.13 (1H, dd, J=8, 8Hz), 7.26 (1H, d, J=8Hz), 7.59 (1H, dd, J=8, 8Hz), 8.01 (1H, d, J=8Hz), 8.23 (1H, d, J=8Hz).

#### Example 103

To a solution of 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]-amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide (10.7 g) in ethanol (155 ml) was added a solution of L-(+)-tartaric acid (2.43 g) in ethanol (60 ml) at 80°C. The solution was stirred at ambient temperature for 1 hour. The solvent was removed at reduced pressure and resulting solid was dissolved in distilled water (1 l) and the solution was filtered through micro filter and the filtrate was lyophilized to give 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide tartrate (5.2 g).

NMR (DMSO-d<sub>6</sub>, δ) : 1.34-1.62 (4H, m), 1.66-1.81 (2H, m), 2.03-2.38 (14H, m), 2.96 (2H, t, J=7Hz), 3.18 (3H, s), 3.37-3.48 (4H, m), 3.74 (3H, s), 3.80-4.04 (4H, m), 4.33 (2H, t, J=7Hz), 6.64 (1H, d, J=8Hz), 6.83 (1H, s), 6.89 (1H, d, J=8Hz), 6.97 (1H, s), 7.02 (1H, d, J=8Hz), 7.13 (1H, dd, J=8, 8Hz), 7.26 (1H, d, J=8Hz), 7.58 (1H, dd, J=8, 8Hz), 8.02 (1H, d, J=8Hz), 8.25 (1H, d, J=8Hz).

- 255 -

Example 104

The following compounds were obtained according to similar manners to those of Examples 16 and 45.

- 5           1) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-(2-methylphenyl)benzamide hydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 2.06-2.32 (5H, m), 2.87-3.05 (2H, m), 3.26 (3H, s), 3.72 (3H, s), 4.35 (2H, t, J=7Hz), 6.84-6.98 (2H, m), 7.08-7.36 (6H, m), 7.58 (1H, dd, J=8, 8Hz), 7.89-8.16 (4H, m), 8.26 (1H, d, J=8Hz)  
10
- 15           2) 4-[3-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.78 (12H, m), 1.98-2.07 (4H, m), 2.24 (3H, s), 2.36 (2H, t, J=8Hz), 2.43-2.54 (1H, m), 2.67 (3H, s), 2.69 (3H, s), 2.92-3.01 (2H, m), 3.19 (3H, s), 3.64 (3H, s), 3.88-4.03 (1H, m), 4.13 (2H, t, J=8Hz), 4.48-4.57 (1H, m), 6.65 (1H, d, J=8Hz), 6.82 (1H, s), 6.88-6.93 (2H, m), 7.03 (1H, d, J=8Hz), 7.17 (1H, d, J=8Hz), 7.38-7.52 (3H, m), 7.62 (1H, d, J=8Hz), 7.92-8.01 (2H, br), 9.33 (1H, s)  
20  
25           ESI-MASS (m/z) : 688 (M+H)
- 30           3) 4-[N-Methyl-2-[(3-aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-methyl-2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.32-1.65 (8H, m), 2.27 (3H, s), 2.33-2.40 (2H, m), 2.77 (3H, s), 2.86-3.02 (5H, m), 3.12 (3H, s), 3.33-3.70 (13H, m), 4.00-4.10 (1H, m), 4.40-4.50 (1H, m), 6.58-6.78 (6H, m), 6.84-7.00 (3H, m), 7.20 (1H, t, J=8Hz), 7.89-7.97 (2H, br s)  
35           ESI-MASS (m/z) : 674

- 256 -

- 4) 4-[2-[3-Aminoprop-1-yl]oxy]benzoyl]amino-3-methoxy-N-methyl-N-[4-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]phenyl]benzamide dihydrochloride

5 NMR (DMSO-d<sub>6</sub>, δ) : 1.37-1.73 (8H, m), 2.12-2.20 (2H, m), 2.37 (2H, t, J=8Hz), 2.72-2.79 (4H, m), 2.89-3.01 (4H, m), 3.29-3.40 (4H, m), 3.80 (3H, s), 3.89 (2H, t, J=8Hz), 3.98-4.04 (1H, m), 4.34-4.41 (3H, m), 6.80-6.86 (3H, m), 7.04-7.19 (4H, m), 7.29 (1H, t, J=8Hz), 7.59 (1H, t, J=8Hz), 7.95-8.06 (4H, m), 8.27 (1H, d, J=8Hz)

10 ESI-MASS (m/z) : 646 (M+H)

- 5) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylhomopiperazin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

15 NMR (CDCl<sub>3</sub>, δ) : 1.47-1.89 (8H, m), 2.27 (3H, s), 2.30-2.46 (4H, m), 2.77-2.96 (2H, m), 3.15-3.63 (11H, m), 3.30 (3H, s), 3.76-4.04 (5H, m), 4.15-4.40 (2H, m), 6.60 (2H, br), 6.78-7.11 (5H, m), 7.43 (1H, br), 7.98-8.05 (1H, m), 8.29-8.37 (1H, m), 8.52 (2H, br)

- 6) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(2-dimethylaminoethyl)aminocarbonyl]pent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

25 NMR (CDCl<sub>3</sub>, δ) : 1.38-1.87 (6H, m), 2.06-2.45 (4H, m), 2.22 (3H, s), 2.25-2.44 (2H, m), 2.76 (3H, s), 2.80 (3H, s), 3.07-3.22 (2H, m), 3.24 (3H, s), 3.54 (2H, br), 3.77-3.95 (2H, m), 3.80 (3H, s), 4.24 (2H, br), 6.57-6.62 (2H, m), 6.80-7.08 (4H, m), 7.39-7.47 (1H, m), 7.97 (1H, d, J=8Hz), 8.20-8.38 (2H, m)

- 7) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[N-(2-dimethylaminoethyl)-N-methylamino-

35

- 257 -

carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
dihydrochloride

NMR (CDCl<sub>3</sub>, δ) : 1.37-1.82 (6H, m), 2.22 (3H, s),  
2.29-2.47 (4H, m), 2.85 (6H, s), 3.02 (3H, s),  
3.08-3.33 (6H, m), 3.26 (3H, s), 3.58-3.95 (4H, m),  
3.83 (3H, s), 4.28 (3H, br), 6.55-6.65 (2H, m),  
6.82-7.06 (5H, m), 7.39-7.47 (1H, m), 8.03 (1H, d,  
J=8Hz), 8.33 (1H, br)

8) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-[N-(3-dimethylaminoprop-1-yl)carbamoyl]-  
pent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.37-1.99 (8H, m), 2.23 (3H, s),  
2.25-2.44 (4H, m), 2.76 (6H, s), 3.05-3.41 (6H, m),  
3.22 (3H, s), 3.78-3.94 (2H, m), 4.22 (2H, br),  
6.56 (2H, br), 6.81-7.04 (5H, m), 7.39 (1H, br),  
8.00 (1H, br), 8.29 (1H, br), 8.56 (3H, br)

9) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-[N-(3-dimethylaminoprop-1-yl)-N-  
methylcarbamoyl]pent-1-yl]oxy-4-methylphenyl]benzamide  
dihydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.33-1.99 (8H, m), 2.26 (3H, s),  
2.26-2.47 (4H, m), 2.78 (6H, s), 2.96 (3H, s),  
3.05-3.39 (6H, m), 3.26 (3H, s), 3.79-3.99 (2H, m),  
3.78 (3H, s), 4.30 (2H, br), 6.62 (2H, m), 6.83-  
7.08 (5H, m), 7.45 (1H, br), 8.01 (1H, br), 8.35  
(1H, br), 8.64 (2H, br)

10) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-(4-hydroxypiperidin-1-yl)carbonylpent-1-  
yl]oxy-4-methylphenyl]benzamide hydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.32-2.06 (10H, m), 2.23 (3H, s),  
2.25-2.40 (4H, m), 2.99-3.07 (2H, m), 3.23 (3H, s),  
3.43-4.00 (7H, m), 4.23 (2H, br), 6.52-6.63 (2H,

- 258 -

m), 6.81-7.12 (4H, m), 7.38-7.49 (1H, m), 7.97 (1H, br), 8.30 (1H, br)

- 5 11) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-aminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

10 NMR (CDCl<sub>3</sub>, δ) : 1.40-1.85 (12H, m), 2.24 (3H, s), 2.28-2.45 (2H), 2.87-3.11 (7H, m), 3.25 (3H, s), 3.84-4.00 (2H, m), 3.79 (3H, s), 4.25 (2H, br), 6.54-6.63 (2H, m), 6.95-7.09 (4H, m), 7.43 (1H, br), 8.04 (1H, br), 8.41 (1H, br)

- 15 12) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)aminocarbonylpent-1-yl]oxy-4-methylphenyl]benzamide trihydrochloride

20 NMR (CDCl<sub>3</sub>, δ) : 1.32-1.80 (6H, m), 2.04-2.15 (2H, m), 2.26 (3H, s), 2.90-3.36 (10H, m), 3.24 (3H, s), 3.76 (3H, s), 3.85-4.02 (2H, m), 4.26 (2H, br), 6.54-6.63 (2H, m), 6.75-7.09 (4H, m), 7.40-7.49 (1H, m), 8.00 (1H, d, J=8Hz), 8.39 (1H, br), 8.62 (1H, br)

- 25 13) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-[bis(2-hydroxyeth-1-yl)amino]carbonylpent-1-yl]oxy-4-methylphenyl]benzamide hydrochloride

30 NMR (CDCl<sub>3</sub>, δ) : 1.50-1.88 (6H, m), 2.05-2.54 (4H, m), 2.28 (3H, s), 3.03 (2H, br), 3.30 (3H, s), 3.41-3.69 (8H, m), 3.78 (3H, s), 3.82-4.00 (2H, m), 4.23 (2H, br), 6.59-6.69 (2H, m), 6.81-7.22 (4H, m), 7.46 (1H, br), 8.09 (1H, br), 8.38 (1H, br)

- 35 14) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(2,2-dimethylhydrazino)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride

NMR (CDCl<sub>3</sub>, δ) : 1.36-1.82 (6H, m), 2.22 (3H, s),



- 259 -

2.26-2.39 (4H, m), 2.88-3.11 (2H, m), 3.11 (6H, s),  
3.32 (3H, s), 3.70-3.94 (2H, m), 3.77 (3H, s), 4.21  
(2H, br), 6.52-6.61 (2H, m), 6.80-7.14 (5H, m),  
7.42 (1H, br), 7.97 (1H, br), 8.25 (3H, br)

5

- 15) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-(carbamoylmethylamino)carbonylpent-1-  
yl]oxy-4-methylphenyl]benzamide hydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.20-1.68 (6H, m), 2.08-2.41 (7H, m),  
2.97-3.35 (5H, m), 3.29-4.27 (9H), 6.38-7.04 (6H,  
m), 7.90-8.29 (6H, m)

10

- 16) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-(2-carbamoylethylamino)carbonylpent-1-  
yl]oxy-4-methylphenyl]benzamide hydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.36-1.81 (6H, m), 2.06-2.40 (6H, m),  
2.23 (3H, s), 3.13 (2H, br), 3.22 (3H, s), 3.32  
(2H, br), 3.55-3.93 (2H, m), 3.78 (3H, s), 4.22  
(2H, br), 6.53-6.63 (2H, m), 6.81-7.04 (5H, m),  
7.39 (1H, br), 7.77 (1H, br), 7.99 (1H, br), 8.28-  
8.47 (3H, m)

15

20

- 17) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-(4-pyridylaminocarbonyl)pent-1-yl]oxy-4-  
methylphenyl]benzamide dihydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.25-1.83 (6H, m), 2.10-2.49 (4H, m),  
2.22 (3H, s), 2.90-3.37 (2H, m), 3.23 (3H, s),  
3.68-3.95 (2H, m), 3.76 (3H, s), 4.21 (2H, br),  
6.51-6.63 (2H, m), 6.66-7.04 (6H, m), 7.88-8.51  
(7H, m)

25

30

- 18) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-  
methyl-N-[2-[5-[4-(diethylaminopiperidin-1-  
yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide  
dihydrochloride

35

- 260 -

NMR (CDCl<sub>3</sub>, δ) : 1.38 (6H, t, J=8Hz), 1.45-1.90 (10H, m), 1.93-2.08 (2H, m), 2.28 (3H, s), 2.30-2.48 (2H, m), 2.92-3.23 (5H, m), 3.25-3.36 (4H, m), 3.29 (3H, s), 3.69 (3H, s), 3.75-4.08 (3H, m), 4.28 (2H, br), 6.54-6.65 (2H, m), 6.81-7.08 (5H, m), 7.45 (1H, br), 7.93 (1H, br), 8.36 (1H, br)

- 5  
10  
15
- 19) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[6-(4-methylpiperazin-1-yl)hex-1-yl]oxy-4-methylphenyl]benzamide trihydrochloride
- NMR (CDCl<sub>3</sub>, δ) : 1.36-1.94 (8H, m), 2.21 (3H, s), 2.25-2.42 (2H, m), 2.90-3.39 (6H, m), 3.10 (3H, s), 3.19 (3H, s), 3.58-4.04 (6H, m), 3.82 (3H, s), 4.18 (1H, br), 6.46-6.63 (2H, m), 6.74-6.98 (4H, m), 7.38 (1H, br), 7.97 (1H, br), 8.28 (1H, br), 8.45 (2H, br)

- 20  
25
- 20) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-(2-pyridyl)phenylmethyl]oxy-4-methylphenyl]-N-methylbenzamide dihydrochloride
- NMR (CDCl<sub>3</sub>, δ) : 2.29 (3H, s), 2.39 (2H, br), 3.17 (2H, br), 3.37 (3H, s), 3.44 (3H, br), 4.12-4.30 (2H, m), 4.73 (1H, br), 5.07 (1H, br), 6.61 (1H, br), 6.70-6.79 (2H, m), 6.94-7.03 (2H, m), 7.12 (1H, d, J=8Hz), 7.38-7.47 (3H, m), 7.89-8.23 (5H, m), 8.73 (3H, br), 8.90 (1H, br)

- 30  
35
- 21) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-[4-[(4-methylpiperazin-1-yl)carbonylamino]but-1-yl]oxy-4-methylphenyl]-N-methylbenzamide dihydrochloride
- NMR (CDCl<sub>3</sub>, δ) : 1.62-2.04 (4H, m), 2.23 (3H, s), 2.27-2.40 (2H, m), 2.74 (3H, s), 3.03-3.14 (2H, m), 3.22 (3H, s), 3.35-3.51 (4H, m), 3.78 (3H, s), 3.85-3.96 (2H, m), 4.26 (2H, br), 6.57-6.64 (2H, m), 6.67-7.09 (5H, m), 7.42 (1H, m), 7.96 (1H, d),

- 261 -

J=8Hz), 8.30 (1H, d, J=8Hz), 8.60 (3H, br)

22) 4-[2-(3-Aminoprop-1-yl)oxybenzoylamino]-3-methoxy-N-[2-  
[4-[(4-dimethylaminopiperidin-1-yl)carbonylamino]but-1-  
yl]oxy-4-methylphenyl]-N-methylbenzamide dihydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.58-2.12 (10H, m), 2.27 (3H, s),  
2.30-2.48 (2H, m), 2.57-2.81 (8H, m), 3.05-3.31  
(7H, m), 3.27 (3H, s), 3.75-3.99 (5H, m), 4.27 (1H,  
br), 6.57-6.63 (2H, m), 6.85-7.09 (5H, m), 7.44  
(2H, br), 7.96 (1H, br), 8.34 (1H, br), 8.75 (1H,  
br)

23) 4-[2-(3-Aminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-[2-  
(4-ureidobut-1-yl)oxy-4-methylphenyl]-N-methylbenzamide  
hydrochloride  
NMR (CDCl<sub>3</sub>, δ) : 1.42-1.81 (4H, m), 2.00-2.15 (2H, m),  
2.25 (3H, s), 2.88 (2H, t, J=5Hz), 2.92 (2H, br),  
3.30 (3H, s), 3.63-3.80 (2H, m), 3.71 (3H, s), 4.21  
(2H, t, J=5Hz), 6.51 (1H, s), 6.71 (1H, d, J=8Hz),  
6.85-7.12 (5H, m), 7.44 (1H, dd, J=2, 8Hz), 8.12  
(1H, d, J=8Hz), 8.36 (1H, d, J=8Hz)

24) 4-[2-[3-Aminoprop-1-yl)oxy]benzoyl]amino-3-chloro-N-  
methyl-N-[2-[5-(4-dimethylaminopiperidin-1-  
yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide  
dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.34-1.50 (2H, m), 1.50-1.62 (2H,  
m), 1.65-1.80 (2H, m), 1.98-2.17 (4H, m), 2.22 (3H,  
s), 2.30-2.40 (2H, m), 2.66 (3H, s), 2.67 (3H, s),  
2.85-3.05 (3H, m), 3.17 (3H, s), 3.33 (1H, br),  
3.80-4.07 (3H, m), 4.33-4.42 (2H, m), 4.47-4.57  
(1H, m), 6.68 (1H, d, J=7Hz), 6.82 (1H, s), 7.08-  
7.23 (3H, m), 7.29 (1H, d, J=7Hz), 7.41 (1H, s),  
7.68 (1H, t, J=6Hz), 7.92 (1H, d, J=7Hz), 8.09 (1H,  
d, J=7Hz)

- 262 -

- 25) 3-(3-Aminoprop-1-yl)oxy-4-[2-[3-aminoprop-1-yl)oxy]-benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide trihydrochloride

5 NMR (DMSO-d<sub>6</sub>, δ) : 1.37-1.50 (2H, m), 1.50-1.62 (2H, m), 1.67-1.80 (2H, m), 1.97-2.19 (4H, m), 2.22 (3H, s), 2.30-2.41 (2H, m), 2.57 (1H, s), 2.92 (6H, br), 3.17 (3H, s), 3.68 (1H, br), 3.93 (2H, br), 4.10 (2H, br), 4.40 (2H, br), 6.66 (1H, d, J=6Hz), 6.78-6.87 (2H, m), 6.95-7.04 (2H, m), 7.12 (1H, t, J=6Hz), 7.29 (1H, d, J=7Hz), 7.57 (1H, t, J=6Hz), 7.93 (1H, d, J=6Hz), 8.14 (1H, d, J=7Hz)

- 15 26) 2-Amino-4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide trihydrochloride

20 NMR (DMSO-d<sub>6</sub>, δ) : 1.41-1.51 (2H, m), 1.51-1.66 (1H, m), 1.74-1.84 (1H, m), 1.98-2.12 (4H, m), 2.30-2.40 (2H, m), 2.67 (3H, s), 2.68 (3H, s), 2.89-3.06 (4H, m), 3.16 (3H, s), 3.33 (2H, br), 3.96-4.10 (4H, m), 4.13-4.20 (2H, m), 4.47-4.58 (1H, m), 6.60 (1H, d, J=7Hz), 6.78 (2H, s), 6.85 (1H, s), 6.97-7.07 (2H, m), 7.13 (1H, d, J=7Hz), 7.27 (1H, s), 7.43-7.56 (2H, m)

- 30 27) 2-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-5-pyridinecarboxamide trihydrochloride

35 NMR (DMSO-d<sub>6</sub>, δ) : 1.32-1.80 (8H, m), 1.97-2.20 (4H, m), 2.22 (3H, s), 2.27-2.40 (3H, m), 2.65 (3H, s), 2.67 (3H, s), 2.92-3.10 (4H, m), 3.19 (3H, s), 3.33 (1H, br), 3.80-4.07 (3H, m), 4.22-4.29 (2H, m), 6.69 (1H, d, J=7Hz), 6.82 (1H, s), 7.07-7.14 (2H, m), 7.20 (1H, d, J=7Hz), 7.56 (1H, t, J=6Hz), 7.66

- 263 -

(1H, d, J=6Hz), 7.78 (1H, d, J=7Hz), 8.00-8.04 (1H, m), 8.23 (1H, s)

- 28) 4-[N-[2-[(3-Aminoprop-1-yl)oxy]phenyl]amino]methyl-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide trihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.35-1.49 (2H, m), 1.49-1.62 (2H, m), 1.62-1.79 (2H, m), 2.01-2.16 (2H, m), 2.23 (3H, s), 2.34-2.40 (2H, m), 2.71 and 2.72 (total 3H, s), 2.76-3.12 (8H, m), 3.17 (3H, s), 3.27-3.41 (2H, m), 3.41-3.54 (4H, m), 3.70-3.81 (1H, m), 3.89-3.98 (1H, m), 4.02-4.08 (3H, m), 4.25 (2H, s), 4.39-4.45 (1H, m), 6.60-6.80 (6H, m), 6.93 (2H, s), 6.98 (1H, d, J=7Hz), 7.10 (1H, d, J=7Hz)

- 29) 4-[2-[(3-Aminoprop-1-yl)oxy]phenyl]oxymethyl-3-methoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.39-1.50 (2H, m), 1.50-1.63 (2H, m), 1.65-1.82 (2H, m), 1.97-2.10 (2H, m), 2.21 (3H, s), 2.35-2.41 (2H, m), 2.71 and 2.72 (total 3H, s), 2.78-3.10 (7H, m), 3.18 (3H, s), 3.29-3.41 (2H, m), 3.41-3.67 (4H, m), 3.82 (1H, br), 3.89-4.00 (1H, m), 4.00-4.12 (3H, m), 4.38-4.48 (1H, m), 4.57 and 4.93 (total 2H, s), 6.61 (1H, d, J=7Hz), 6.69-6.97 (6H, m), 6.97-7.07 (2H, m), 7.20-7.25 (1H, m)

- 30) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-benzyloxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.37-1.50 (2H, m), 1.50-1.63 (2H, m), 1.63-1.79 (2H, m), 1.79-1.91 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=6Hz), 2.60-2.77 (5H, m), 2.79-3.10 (4H, m), 3.15 (3H, s), 3.30-3.67 (3H, m),

- 264 -

5 3.77-4.12 (5H, m), 4.37-4.49 (1H, m), 5.06 (2H, s),  
6.62 (1H, d, J=6Hz), 6.82 (1H, s), 6.90 (1H, d,  
J=7Hz), 6.97 (1H, d, J=7Hz), 7.03 (1H, s), 7.12  
(1H, t, J=7Hz), 7.22 (1H, d, J=7Hz), 7.30-7.46 (5H,  
m), 7.54 (1H, t, J=6Hz), 7.97 (1H, d, J=7Hz), 8.23  
(1H, d, J=7Hz)

10 31) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-hydroxy-N-  
methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-  
yloxy]-4-methylphenyl]benzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.43 (2H, br), 1.49-1.62 (2H, m),  
1.63-1.82 (2H, m), 2.00-2.40 (16H, m), 2.90-2.97  
(2H, m), 3.14 (3H, s), 3.30-3.50 (5H, m), 3.89 (2H,  
15 br), 4.20-4.38 (2H, m), 6.50-6.68 (2H, m), 6.80  
(1H, s), 6.87-6.99 (2H, m), 7.12 (1H, t, J=6Hz),  
7.22 (1H, d, J=6Hz), 7.49-7.60 (1H, m), 7.97-8.18  
(2H, m)

20 32) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-  
ethoxycarbonylmethoxy-N-methyl-N-[2-[5-(4-  
methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-  
methylphenyl]benzamide dihydrochloride  
NMR (DMSO-d<sub>6</sub>, δ) : 1.09 and 1.22 (total 3H, t, J=6Hz),  
25 1.37-1.51 (2H, m), 1.51-1.66 (2H, m), 1.67-1.80  
(2H, m), 2.05-2.18 (2H, m), 2.23 (3H, s), 2.38 (2H,  
t, J=6Hz), 2.73-2.74 (3H, m), 2.90-3.10 (5H, m),  
3.17 (3H, s), 3.30-3.58 (2H, m), 3.80-4.00 (2H, m),  
4.00-4.20 (3H, m), 4.32-4.50 (3H, m), 4.80 (2H, s),  
30 6.62 (1H, d, J=6Hz), 6.82 (1H, s), 6.89-6.92 (2H,  
m), 7.01 (1H, d, J=7Hz), 7.15 (1H, t, J=6Hz), 7.27  
(1H, d, J=7Hz), 7.58 (1H, t, J=6Hz), 8.00 (1H, d,  
J=6Hz), 8.27 (1H, d, J=7Hz)

35 33) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-  
methoxycarbonylmethoxy-N-methyl-N-[2-[5-(4-

- 265 -

methyloperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.35-1.50 (2H, m), 1.50-1.63 (2H, m), 1.63-1.90 (2H, m), 2.00-2.14 (2H, m), 2.21 (3H, s), 2.25-2.43 (2H, m), 2.71 (3H, s), 2.77-3.05 (5H, m), 3.15 (3H, s), 3.18-3.57 (6H, m), 3.70 (3H, s), 3.73-4.12 (3H, m), 4.12-4.49 (3H, m), 4.80 (2H, s), 6.63 (1H, d, J=7Hz), 6.70-7.20 (5H, m), 7.27 (1H, d, J=7Hz), 7.57 (1H, t, J=7Hz), 7.93-8.10 (1H, m), 8.23 (1H, d, J=6Hz)

34) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-dimethylaminocarbonylmethoxy-N-methyl-N-[2-[5-(4-methyloperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.37-1.49 (2H, m), 1.50-1.62 (2H, m), 1.63-1.79 (2H, m), 1.98-2.10 (2H, m), 2.21 (3H, s), 2.32-2.43 (2H, m), 2.71 (3H, s), 2.86 (3H, s), 2.98 (3H, s), 2.82-3.05 (5H, m), 3.15 (3H, s), 3.90 (2H, br), 4.02-4.12 (2H, m), 4.28-4.38 (2H, m), 4.38-4.48 (1H, m), 4.83 (2H, s), 6.62 (1H, d, J=7Hz), 6.80 (1H, s), 6.82-6.92 (2H, m), 7.00 (1H, d, J=7Hz), 7.12 (1H, t, J=7Hz), 7.23 (1H, d, J=7Hz), 7.55 (1H, t, J=7Hz), 8.20 (1H, d, J=7Hz)

35) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-methylaminocarbonylmethoxy-N-methyl-N-[2-[5-(4-methyloperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.51 (1H, m), 1.51-1.65 (2H, m), 1.68-1.80 (2H, m), 2.00-2.23 (2H, m), 2.22 (3H, s), 2.34-2.40 (3H, m), 2.50 (3H, s), 2.58 (2H, br), 2.62 (3H, s), 2.63 (3H, s), 2.90 (4H, br), 3.15 (3H, s), 3.88-3.97 (2H, m), 4.26-4.33 (2H, m), 4.37-4.54 (2H, m), 6.62 (1H, d, J=7Hz), 6.82 (2H,

- 266 -

s), 6.88 (1H, d, J=7Hz), 6.97 (1H, d, J=7Hz), 7.12 (1H, t, J=7Hz), 7.22 (1H, d, J=7Hz), 7.57 (1H, t, J=7Hz), 7.90 (1H, d, J=7Hz), 8.12-8.25 (2H, m)

- 5        36) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-aminocarbonylmethoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]benzamide dihydrochloride

10        NMR (DMSO-d<sub>6</sub>, δ) : 1.38-1.52 (2H, m), 1.52-1.67 (2H, m), 1.68-1.83 (2H, m), 2.00-2.15 (2H, m), 2.23 (3H, s), 2.39 (2H, t, J=6Hz), 2.62 and 2.63 (total 3H, s), 2.72 and 2.73 (total 3H, s), 2.80-3.10 (6H, m), 3.15 (3H, s), 3.87-3.98 (2H, m), 4.03-4.13 (1H, m), 6.27-6.37 (1H, m), 6.37-6.56 (2H, m), 6.62 (1H, d, J=7Hz), 6.82 (2H, s), 6.90 (1H, d, J=7Hz), 6.98 (1H, d, J=6Hz), 7.12 (1H, t, J=7Hz), 7.26 (1H, d, J=7Hz), 7.57 (1H, t, J=6Hz), 7.92 (1H, d, J=7Hz), 8.13-8.30 (2H, m)

- 20        37) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-3-propoxybenzamide dihydrochloride

25        NMR (DMSO-d<sub>6</sub>, δ) : 0.89 (3H, t, J=6Hz), 1.37-1.50 (2H, m), 1.50-1.68 (4H, m), 1.68-1.80 (2H, m), 2.02-2.18 (2H, m), 2.20 (3H, s), 2.38 (2H, t, J=6Hz), 2.47 (3H, s), 2.75-3.12 (5H, m), 3.17 (3H, s), 3.30-3.42 (2H, m), 3.42-3.56 (1H, m), 3.80-4.00 (4H, m), 4.00-4.13 (1H, m), 4.32-4.50 (4H, m), 6.61 (1H, d, J=7Hz), 6.82 (1H, s), 6.88 (1H, s), 6.94 (1H, d, J=7Hz), 7.02 (1H, d, J=7Hz), 7.13 (1H, t, J=7Hz), 7.29 (1H, d, J=7Hz), 7.56 (1H, t, J=7Hz), 7.97 (1H, d, J=7Hz), 8.22 (1H, d, J=7Hz)

- 35        38) 4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-isopropoxy-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-



- 267 -

yloxy]-4-methylphenyl]benzamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.10-1.27 (6H, m), 1.37-1.50 (2H, m), 1.50-1.64 (2H, m), 1.67-1.82 (2H, m), 2.03-2.07 (2H, m), 2.22 (3H, s), 2.39 (2H, t, J=6Hz), 2.72 and 2.73 (total 3H, s), 2.78-3.12 (6H, m), 3.17 (3H, s), 3.30-3.43 (2H, m), 3.43-3.60 (1H, m), 3.80-4.02 (2H, m), 4.02-4.13 (1H, m), 4.23-4.50 (4H, m), 6.64 (1H, d, J=7Hz), 6.81-6.90 (2H, m), 6.98 (1H, d, J=7Hz), 7.03 (1H, d, J=7Hz), 7.13 (1H, t, J=6Hz), 7.32 (1H, d, J=7Hz), 7.56 (1H, t, J=6Hz), 7.94 (1H, d, J=6Hz), 8.22 (1H, d, J=7Hz)

39) 2-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-N-methyl-N-[2-[5-(4-methylpiperazin-1-yl)carbonylpent-1-yloxy]-4-methylphenyl]-5-thiophenecarboxamide dihydrochloride

NMR (DMSO-d<sub>6</sub>, δ) : 1.20-1.38 (2H, m), 1.38-1.52 (2H, m), 1.53-1.70 (2H, m), 1.98-2.10 (2H, m), 2.22-2.32 (2H, m), 2.33 (3H, s), 2.69-2.72 (3H, m), 2.76-3.07 (5H, m), 3.16 (3H, s), 3.27-3.54 (3H, m), 3.78-4.09 (3H, m), 4.10-4.20 (2H, m), 4.33-4.47 (2H, m), 6.15 (1H, br), 6.55 (1H, d, J=5Hz), 6.81 (1H, d, J=7Hz), 6.97 (1H, s), 7.07 (1H, t, J=6Hz), 7.13-7.20 (2H, m), 7.44-7.60 (2H, m)

#### Example 105

To a solution of 4-[2-[(3-tert-butoxycarbonylaminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-(4-hydroxyphenyl)benzamide (50 mg) in chloroform (3.0 ml) was added a solution of 4N hydrogen chloride in ethyl acetate (1.0 ml) and the mixture was stirred at ambient temperature for 2 hours. The resulting mixture was evaporated in vacuo and the residue was solidified with diethyl ether. Diethyl ether was removed in vacuo to give 4-[2-[(3-aminoprop-1-yl)oxy]benzoyl]amino-3-methoxy-N-methyl-N-(4-hydroxyphenyl)-benzamide hydrochloride (40 mg).

- 268 -

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 2.11-2.21 (2H, m), 2.96 (2H, q, J=8Hz), 3.30 (3H, s), 3.78 (3H, s), 4.37 (2H, t, J=8Hz), 6.66 (2H, d, J=8Hz), 6.88 (1H, d, J=8Hz), 6.97 (1H, s), 6.99 (2H, d, J=8Hz), 7.15 (1H, t, J=8Hz), 7.27 (1H, d, J=8Hz), 7.55-7.62 (1H, m), 7.97-8.05 (3H, m), 8.28 (1H, d, J=8Hz), 9.54-9.59 (1H, br s)

ESI-MASS (m/z) : 450 (M+H)

Example 106

The following compound was obtained according to a similar manner to that of Example 105.

4-[2-[(3-Aminoprop-1-yl)oxy]benzoyl]amino-3-carboxymethoxy-N-methyl-N-cyclohexylbenzamide hydrochloride

NMR (DMSO-d<sub>6</sub>,  $\delta$ ) : 1.02-1.10 (2H, m), 1.46-1.80 (8H, m), 2.08-2.12 (2H, m), 2.80 (3H, s), 2.92-2.99 (2H, m), 3.30-3.47 (2H, br), 4.39 (2H, t, J=7Hz), 4.96 (2H, s), 6.98-7.04 (2H, br s), 7.18 (1H, t, J=8Hz), 7.30 (1H, d, J=8Hz), 7.60 (1H, t, J=8Hz), 7.95-8.05 (3H, br), 8.07 (1H, d, J=8Hz), 8.51 (1H, d, J=8Hz)

ESI-MASS (m/z) : 484 (M+H)

Example 107

1) A solution of 4-[2-[3-(9-fluorenylmethyl)oxycarbonylaminoprop-1-yl]thiobenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide (110 mg) in a mixture of N,N-dimethylformamide and piperidine (4:1, 5 ml) was stirred at ambient temperature for 30 minutes and the resulting solution was diluted with ethyl acetate (20 ml). The solution was washed with water (10 ml x 3) and brine, and the solution was dried over potassium carbonate. The solvent was evaporated and the residue was purified on basic silica gel column chromatography (SiO<sub>2</sub> 30 g, 1-15% methanol in chloroform) to

- 269 -

give 4-[2-(3-aminoprop-1-yl)thiobenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide.

5 NMR (CDCl<sub>3</sub>, δ) : 1.36-1.92 (12H, m), 2.29 (6H, s),  
2.30 (3H, s), 2.36 (2H, t, J=5Hz), 2.59 (1H, t,  
J=11Hz), 2.77 (2H, t, J=5Hz), 2.99 (2H, t, J=5Hz),  
3.32 (3H, s), 3.75 (3H, s), 3.85-4.03 (4H, m),  
6.57-6.66 (2H, m), 6.84-6.90 (1H, d, J=8Hz), 7.02  
10 (1H, s), 7.39-7.48 (3H, m), 7.65 (1H, d, J=8Hz),  
8.30 (1H, d, J=8Hz), 8.80 (1H, s)

2) To a solution of the obtained compound in ethanol (5 ml) was added 1N hydrochloric acid (0.15 ml). The volatile solvent was removed by evaporation and the residue was  
15 lyophilized to give 4-[2-(3-aminoprop-1-yl)thiobenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)-carbonylpent-1-yl]oxy-4-methylphenyl]benzamide dihydrochloride (45 mg).

20 NMR (CDCl<sub>3</sub>, δ) : 1.44-1.92 (6H, m), 2.02-2.16 (2H, m),  
2.28 (3H, s), 2.30-2.41 (2H, m), 2.73 (6H, br),  
2.99-3.14 (2H, m), 3.27-3.33 (1H, m), 3.31 (3H, s),  
3.62-3.79 (4H, m), 3.71 (3H, s), 3.82-4.10 (2H, m),  
6.55-6.67 (2H, m), 6.83-7.02 (5H, m), 7.35-7.52  
25 (2H, m), 8.23 (1H, br), 8.54 (2H, br)

#### Example 108

The following compound was obtained according to a similar manner to that of Example 15.

30 4-2-(3-Dimethylaminoprop-1-yl)oxybenzoyl]amino-3-methoxy-N-methyl-N-[2-[5-(4-dimethylaminopiperidin-1-yl)carbonylpent-1-yl]oxy-4-methylphenyl]benzamide

NMR (CDCl<sub>3</sub>, δ) : 1.49-1.60 (2H, m), 1.66-1.95 (4H, m),  
2.21 (6H, s), 2.27 (6H, s), 2.35-2.48 (4H, m), 2.58  
35 (2H, t, J=11Hz), 3.32 (2H, t, J=11Hz), 3.33 (3H,

- 270 -

s), 3.80 (3H, s), 3.82-4.00 (2H, m), 4.25 (2H, t, J=5Hz), 4.64 (1H, br), 6.55-6.64 (2H, m), 6.85 (1H, d, J=8Hz), 6.89 (1H, d, J=8Hz), 7.00-7.11 (3H, m), 7.26 (1H, s), 7.40-7.48 (1H, m), 8.21 (1H, d, J=8Hz), 8.40 (1H, d, J=8Hz)

5

10

15

20

25

30

35


- 271 -

## C L A I M S

1. A compound of the formula :

5

10



15

wherein

20

25

30

35

$R^1$  is aryl, cyclo(lower)alkyl or a heterocyclic group,  
 each of which may be substituted with substituent(s)  
 selected from the group consisting of halogen;  
 hydroxy; nitro; amino; acyl; substituted acyl;  
 acyl(lower)alkylsulfinyl; acyl(lower)alkylsulfonyl;  
 acyloxy; lower alkylamino(lower)alkylcarbamoyloxy;  
 aryl; cyano; a heterocyclic group;  
 lower alkenyl optionally substituted with acyl,  
 substituted acyl, aryl or acyl-substituted aryl;  
 lower alkynyl optionally substituted with amino,  
 acylamino or substituted acylamino;  
 lower alkyl optionally substituted with halogen,  
 amino, lower alkylamino, acylamino, substituted  
 acylamino, hydroxy, acyloxy, acyl(lower)alkanoyloxy,  
 acyl, substituted acyl, acyl(lower)alkoxyimino, aryl  
 or acyl-substituted aryl;  
 lower alkylthio optionally substituted with acyl or  
 substituted acyl;  
 alkoxy optionally substituted with aryl, substituted  
 aryl, hydroxy, acyloxy, amino, lower alkylamino,

- 272 -

protected amino, a heterocyclic group, acyl-substituted pyridyl, substituted acyl-substituted pyridyl, halogen, acyl(lower)alkylamino, N-protected-acyl(lower)alkylamino, N-acyl(lower)alkyl-N-lower alkylamino, acyl, substituted acyl, acylamino, substituted acylamino, lower alkylhydrazinocarbonylamino, hydroxyimino, acyl(lower)alkoxyimino, substituted acyl(lower)alkoxyimino, acyl(lower)alkoxy, guanidino or N-protected guanidino; and lower alkenyloxy optionally substituted with acyl or substituted acyl;

5  $R^2$  is hydrogen; lower alkyl optionally substituted with hydroxy, aryl or acyl; or cyclo(lower)alkyl;

10  $R^3$  is hydrogen; halogen; hydroxy; acyloxy; substituted acyloxy; lower alkyl optionally substituted with hydroxy or lower alkoxy; lower alkoxy optionally substituted with aryl, amino, protected amino, acyl, hydroxy, cyano or lower alkylthio; nitro; amino; acyl; substituted acyl; or cyclo(lower)alkyloxy;

15  $R^4$  is hydroxy; halogen; nitro; amino; protected amino; lower alkylamino; acyloxy; amino(lower)alkylamino; N-protected amino(lower)alkylamino; lower alkoxy optionally substituted with hydroxy, aryl, substituted aryl, acyl, substituted acyl, amino, lower alkylamino, acylamino, substituted acylamino, protected amino, a heterocyclic group or guanidino; lower alkylthio optionally substituted with acyl, substituted acyl, amino, lower alkylamino, acylamino, substituted acylamino, protected amino, a heterocyclic group, hydroxy, lower alkylsulfonyloxy, arylsulfonyloxy, ar(lower)alkoxy or substituted ar(lower)alkoxy; lower alkyl substituted with acyl, substituted acyl, amino, lower alkylamino, acylamino, substituted acylamino, protected amino, a

20  
25  
30  
35

- 273 -

heterocyclic group, hydroxy, lower alkylsulfonyloxy  
 or arylsulfonyloxy; lower alkenyl optionally  
 substituted with acyl; lower alkynyl optionally  
 substituted with hydroxy, amino, protected amino,  
 5 lower alkylsulfonyloxy or arylsulfonyloxy;  
 amino(lower)alkylsulfonyl; N-protected  
 amino(lower)alkylsulfonyl; lower alkylaminosulfonyl;  
 a heterocyclicsulfonyl; amino(lower)alkylsulfinyl;  
 N-protected amino(lower)alkylsulfinyl; piperidyloxy;  
 10 or N-protected piperidyloxy;  
 $R^5$  is hydrogen, lower alkyl, lower alkoxy or halogen;  
 A is a single bond, O or NH;  
 E is lower alkylene, lower alkenylene,  $-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ ,  $-\overset{\text{O}}{\underset{\text{O}}{\text{S}}}-$ , or  
 15 a group of the formula :



in which G is lower alkylene and J is O or  $-\overset{\text{R}^6}{\underset{|}{\text{N}}}-$   
 20 (wherein  $R^6$  is hydrogen or N-protective group);  
 X is  $-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{N}-$  or S; and  
 Y is CH or N;  
 and pharmaceutically acceptable salts thereof.

25 2. A compound according to claim 1, wherein  
 $R^1$  is aryl which may be substituted with lower alkoxy  
 optionally substituted with acylamino or acyl;  
 $R^2$  is lower alkyl;  
 $R^3$  is hydrogen, lower alkyl or lower alkoxy;  
 30  $R^4$  is hydroxy, or lower alkoxy, lower alkylthio or  
 lower alkyl, each of which may be substituted  
 with hydroxy, aryl, substituted aryl, acyl,  
 amino, lower alkylamino, acylamino, protected  
 amino or a heterocyclic group;  
 35  $R^5$  is hydrogen, lower alkyl, lower alkoxy or halogen;

- 274 -

A is NH;  
 E is  $\text{-}\overset{\text{O}}{\parallel}\text{C-}$ ;  
 X is  $\text{-CH=CH-}$ ; and  
 Y is CH.

5

3. A compound according to claim 2, wherein  
 $R^1$  is phenyl or tolyl, each of which is substituted  
 with lower alkoxy substituted with acyl;  
 $R^3$  is lower alkoxy or lower alkyl; and  
 $R^4$  is lower alkoxy, lower alkylthio or lower alkyl,  
 each of which is substituted with amino or  
 hydroxy.

10

4. A compound according to claim 3, wherein  
 $R^1$  is phenyl or tolyl, each of which is substituted  
 with lower alkoxy substituted with N-(lower  
 alkyl)piperazinylcarbonyl;  
 $R^3$  is lower alkoxy;  
 $R^4$  is lower alkoxy substituted with amino; and  
 $R^5$  is hydrogen.


15

20

5. A process for preparing the formula :

25

30



(I)

wherein

$R^1$  is aryl, cyclo(lower)alkyl or a heterocyclic group,  
 each of which may be substituted with substituent(s)

35



- 275 -

selected from the group consisting of halogen;  
hydroxy; nitro; amino; acyl; substituted acyl;  
acyl(lower)alkylsulfinyl; acyl(lower)alkylsulfonyl;  
acyloxy; lower alkylamino(lower)alkylcarbamoxyloxy;  
5 aryl; cyano; a heterocyclic group;  
lower alkenyl optionally substituted with acyl,  
substituted acyl, aryl or acyl-substituted aryl;  
lower alkynyl optionally substituted with amino,  
acylamino or substituted acylamino;  
10 lower alkyl optionally substituted with halogen,  
amino, lower alkylamino, acylamino, substituted  
acylamino, hydroxy, acyloxy, acyl(lower)alkanoyloxy,  
acyl, substituted acyl, acyl(lower)alkoxyimino, aryl  
or acyl-substituted aryl;  
15 lower alkylthio optionally substituted with acyl or  
substituted acyl;  
alkoxy optionally substituted with aryl, substituted  
aryl, hydroxy, acyloxy, amino, lower alkylamino,  
protected amino, a heterocyclic group, acyl-  
20 substituted pyridyl, substituted acyl-substituted  
pyridyl, halogen, acyl(lower)alkylamino, N-protected-  
acyl(lower)alkylamino, N-acyl(lower)alkyl-N-lower  
alkylamino, acyl, substituted acyl, acylamino,  
substituted acylamino, lower  
25 alkylhydrazinocarbonylamino, hydroxyimino,  
acyl(lower)alkoxyimino, substituted  
acyl(lower)alkoxyimino, acyl(lower)alkoxy, guanidino  
or N-protected guanidino; and  
lower alkenyloxy optionally substituted with acyl or  
30 substituted acyl;  
R<sup>2</sup> is hydrogen; lower alkyl optionally substituted with  
hydroxy, aryl or acyl; or cyclo(lower)alkyl;  
R<sup>3</sup> is hydrogen; halogen; hydroxy; acyloxy; substituted  
acyloxy; lower alkyl optionally substituted with  
35 hydroxy or lower alkoxy; lower alkoxy optionally

- 276 -

substituted with aryl, amino, protected amino, acyl, hydroxy, cyano or lower alkylthio; nitro; amino; acyl; substituted acyl; or cyclo(lower)alkyloxy;

5  $R^4$  is hydroxy; halogen; nitro; amino; protected amino; lower alkylamino; acyloxy; amino(lower)alkylamino; N-protected amino(lower)alkylamino; lower alkoxy optionally substituted with hydroxy, aryl, substituted aryl, acyl, substituted acyl, amino, lower alkylamino, acylamino, substituted acylamino, 10 protected amino, a heterocyclic group or guanidino; lower alkylthio optionally substituted with acyl, substituted acyl, amino, lower alkylamino, acylamino, substituted acylamino, protected amino, a heterocyclic group, hydroxy, lower alkylsulfonyloxy, 15 arylsulfonyloxy, ar(lower)alkoxy or substituted ar(lower)alkoxy; or lower alkyl substituted with acyl, substituted acyl, amino, lower alkylamino, acylamino, substituted acylamino, protected amino, a heterocyclic group, hydroxy, lower alkylsulfonyloxy or arylsulfonyloxy; lower alkenyl optionally 20 substituted with acyl; lower alkynyl optionally substituted with hydroxy, amino, protected amino, lower alkylsulfonyloxy or arylsulfonyloxy; amino(lower)alkylsulfonyl; N-protected amino(lower)alkylsulfonyl, lower alkylaminosulfonyl; 25 a heterocyclicsulfonyl; amino(lower)alkylsulfinyl; N-protected amino(lower)alkylsulfinyl; piperidyloxy; or N-protected piperidyloxy;

30  $R^5$  is hydrogen, lower alkyl, lower alkoxy or halogen; A is a single bond, O or NH; E is lower alkylene, lower alkenylene,  $\begin{smallmatrix} \text{O} \\ \parallel \\ -\text{C}- \end{smallmatrix}$ ,  $\begin{smallmatrix} \text{O} \\ \parallel \\ -\text{S}- \\ \parallel \\ \text{O} \end{smallmatrix}$ , or a group of the formula :

-G-J-

- 277 -

in which G is lower alkylene and J is O or  $\begin{array}{c} R^6 \\ | \\ -N- \end{array}$


(wherein  $R^6$  is hydrogen or N-protective group);

X is  $-CH=CH-$ ,  $-CH=N-$  or S; and


Y is CH or N;

or pharmaceutically acceptable salts thereof,  
which comprises,

1) reacting a compound of the formula :




or its salt with a compound of the formula :



or its reactive derivative at the carboxy group or  
the sulfo group, or a salt thereof to provide a  
compound of the formula :

- 278 -

5



10

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , X and Y are each as defined  
 above, and

15

$E_a$  is  $-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ ,  $-\overset{\text{O}}{\underset{\text{O}}{\text{S}}}-$ , or


2) reacting a compound of the formula :

20



or its salt with a compound of the formula :

25




30

or its reactive derivative at the carboxy group  
 or a salt thereof to provide a compound of the  
 formula :

35

- 279 -

5




10

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined  
 above, or

15

3) subjecting a compound of the formula :


20



25

or its salt to deesterification reaction to provide a  
 compound of the formula :

30



35

- 280 -

or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_a^1$  is aryl, haloaryl, cyclo(lower)alkyl or


5 a heterocyclic group, each of which is substituted  
with esterified carboxy; lower alkenyl substituted  
with esterified carboxy or esterified carboxy-  
substituted aryl; lower alkyl substituted with  
esterified carboxy, esterified  
10 carboxy(lower)alkanoyloxy or esterified  
carboxy(lower)alkoxyimino;  
lower alkylthio substituted with esterified carboxy;  
alkoxy substituted with esterified carboxy-  
substituted aryl, esterified carboxy-substituted  
pyridyl, esterified carboxy(lower)alkylamino,  
15 N-protected-esterified carboxy(lower)alkylamino,  
N-esterified carboxy(lower)alkyl-N-lower alkylamino,  
esterified carboxy or esterified  
carboxy(lower)alkoxyimino; or lower alkenyloxy  
substituted with esterified carboxy; and

20  $R_D^1$  is aryl, haloaryl, cyclo(lower)alkyl or  
a heterocyclic group, each of which is substituted  
with carboxy; lower alkenyl substituted with carboxy;  
lower alkyl substituted with carboxy or carboxy-  
substituted aryl, carboxy(lower)alkanoyloxy or  
25 carboxy(lower)alkoxyimino;  
lower alkylthio substituted with carboxy;  
alkoxy substituted with carboxy-substituted aryl,  
carboxy-substituted pyridyl,  
carboxy(lower)alkylamino, N-protected-  
30 carboxy(lower)alkylamino, N-carboxy(lower)alkyl-N-  
lower alkylamino, carboxy or  
carboxy(lower)alkoxyimino; or lower alkenyloxy  
substituted with carboxy; or

35 4) subjecting a compound of the formula :

- 281 -


5



10

or its salt to deesterification reaction to provide a compound of the formula :

15



20

25

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined  
 above,

30

$R_a^4$  is lower alkoxy substituted with esterified  
 carboxy; lower alkylthio substituted with  
 esterified carboxy; lower alkyl substituted with  
 esterified carboxy; or lower alkenyl substituted  
 with esterified carboxy; and


35

$R_b^4$  is lower alkoxy substituted with carboxy;  
 lower alkylthio substituted with carboxy;  
 lower alkyl substituted with carboxy; or


- 282 -

lower alkenyl substituted with carboxy; or

5) subjecting a compound of the formula :



15 or its salt to elimination reaction of the N-protective group to provide a compound of the formula :




25 or its salt, in the above formulas,  
 30  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R^4$  is protected amino; N-protected piperidyloxy;  
 N-protected amino(lower)alkylamino; lower alkoxy  
 substituted with protected amino;  
 lower alkylthio substituted with protected amino;  
 lower alkyl substituted with protected amino; lower  
 35 alkynyl substituted with protected amino; or  
 N-protected amino(lower)alkylsulfonyl; and




- 283 -

$R_D^4$  is amino; piperidyloxy; amino(lower)alkylamino;  
 lower alkoxy substituted with amino; lower  
 alkylthio substituted with amino; lower alkyl  
 substituted with amino; lower alkynyl substituted  
 with amino; or amino(lower)alkylsulfonyl; or

6) reacting a compound of the formula :



or its reactive derivative at the carboxy group or a salt  
 thereof with an amine or its salt to provide a compound of  
 the formula :



or its salt, in the above formulas,  
 $R_D^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $A$ ,  $E$ ,  $X$  and  $Y$  are each as defined  
 above, and


- 284 -

$R_C^1$  is aryl, haloaryl, cyclo(lower)alkyl or  
a heterocyclic group, each of which is substituted  
with substituted or unsubstituted N-containing  
heterocycliccarbonyl; carbamoyl; substituted or  
5 unsubstituted lower alkylcarbamoyl; lower alkenyl  
substituted with substituted or unsubstituted  
N-containing heterocycliccarbonyl, carbamoyl,  
substituted or unsubstituted lower alkylcarbamoyl or  
N-containing heterocycliccarbonyl-substituted aryl;  
10 lower alkyl substituted with substituted or  
unsubstituted N-containing heterocycliccarbonyl,  
carbamoyl, substituted or unsubstituted lower  
alkylcarbamoyl, substituted or unsubstituted  
N-containing heterocycliccarbonyl(lower)alkanoyloxy,  
15 carbamoyl(lower)alkanoyloxy, substituted or  
unsubstituted lower alkylcarbamoyl(lower)alkanoyloxy,  
substituted or unsubstituted N-containing  
heterocycliccarbonyl(lower)alkoxyimino,  
carbamoyl(lower)alkoxyimino or substituted or  
20 unsubstituted lower alkylcarbamoyl(lower)alkoxyimino;  
lower alkylthio substituted with substituted or  
unsubstituted N-containing heterocycliccarbonyl,  
carbamoyl or substituted or unsubstituted lower  
alkylcarbamoyl; alkoxy substituted with substituted  
25 or unsubstituted N-containing heterocycliccarbonyl-  
substituted aryl, carbamoyl-substituted aryl,  
substituted or unsubstituted lower alkylcarbamoyl-  
substituted aryl, substituted or unsubstituted N-  
containing heterocycliccarbonyl-substituted pyridyl,  
30 carbamoyl-substituted pyridyl, substituted or  
unsubstituted lower alkylcarbamoyl-substituted  
pyridyl, substituted or unsubstituted N-containing  
heterocycliccarbonyl(lower)alkylamino,  
carbamoyl(lower)alkylamino, substituted or  
35 unsubstituted lower alkylcarbamoyl(lower)alkylamino,

- 285 -

N-protected-(substituted or unsubstituted  
 N-containing heterocyclic)carbonyl(lower)alkylamino,  
 N-protected-carbamoyl(lower)alkylamino, N-protected-  
 substituted or unsubstituted lower alkylcarbamoyl-  
 (lower)alkylamino, N-(substituted or unsubstituted  
 N-containing heterocyclic)carbonyl(lower)alkyl-N-  
 lower alkylamino, N-carbamoyl(lower)alkyl-N-lower  
 alkylamino, substituted or unsubstituted N-lower  
 alkylcarbamoyl-N-lower alkylamino, substituted or  
 unsubstituted N-containing heterocycliccarbonyl,  
 carbamoyl, substituted or unsubstituted lower  
 alkylcarbamoyl, substituted or unsubstituted  
 N-containing heterocycliccarbonyl(lower)alkoxyimino,  
 carbamoyl(lower)alkoxyimino or substituted or  
 unsubstituted lower alkylcarbamoyl(lower)alkoxyimino;  
 or lower alkenyloxy substituted with substituted or  
 unsubstituted N-containing heterocycliccarbonyl,  
 carbamoyl or substituted or unsubstituted lower  
 alkylcarbamoyl; or


7) reacting a compound of the formula :



or its reactive derivative at the carboxy group  
 or a salt thereof with an amine or its salt to  
 provide a compound of the formula :

- 286 -

5



10

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined  
 above, and


15

$R^4_e$  is lower alkoxy, lower alkylthio, lower alkyl, or  
 lower alkenyl each of which is substituted with  
 substituted or unsubstituted N-containing  
 heterocycliccarbonyl, carbamoyl, or substituted or  
 unsubstituted lower alkylcarbamoyl; or

20

8) subjecting a compound of the formula :

25



30


35

or its salt to debenzylation reaction to provide a

- 287 -

compound of the formula :

5



10

15

or its salt, in the above formulas,

 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above, $R^4_f$  is methoxy substituted with aryl or substituted aryl;

or lower alkylthio which is substituted with

methoxy substituted with aryl or substituted aryl;


20

and

 $R^4_g$  is hydroxy; or lower alkylthio substituted with  
hydroxy; or

9) reacting a compound of the formula :

25



30


35

- 288 -

or its salt with a compound of the formula :



or its salt to provide a compound of the formula :



or its salt, in the above formulas,

$R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,

$R_{ga}^4$  is hydroxy;

$R^7$  is lower alkyl optionally substituted with hydroxy, aryl, substituted aryl, acyl, amino, lower alkylamino, acylamino, protected amino or a heterocyclic group; or N-protected piperidyl;


$Z^1$  is hydroxy; or acid residue; and

$R_h^4$  is lower alkoxy substituted with hydroxy, aryl, substituted aryl, acyl, amino, lower alkylamino, acylamino, protected amino or a heterocyclic group; or N-protected piperidyloxy; or

10) reacting a compound of the formula :

- 289 -

5




10

15

or its salt with an acylating agent to provide a compound of the formula :

20



25

30

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_{da}^4$  is lower alkoxy substituted with amino; lower  
 alkylthio substituted with amino; or lower alkyl  
 substituted with amino; and


35

$R_i^4$  is lower alkoxy substituted with acylamino or


- 290 -

substituted acylamino; lower alkylthio substituted  
with acylamino or substituted acylamino; or lower  
alkyl substituted with acylamino or substituted  
acylamino; or

11) reacting a compound of the formula :



or its salt with lower alkanal or N-protected  
amino(lower)alkanal in the presence of a reducing agent  
to provide a compound of the formula :





- 291 -


or its salt, in the above formulas,

$R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined  
above,


$R_{db}^4$  is amino; lower alkoxy substituted with amino; lower  
alkylthio substituted with amino; or lower alkyl  
substituted with amino; and

$R_j^4$  is lower alkoxy substituted with lower alkylamino;  
lower alkylthio substituted with lower alkylamino;  
lower alkyl substituted with lower alkylamino;  
lower alkylamino; or N-protected  
amino(lower)alkylamino; or

12) subjecting a compound of the formula :




or its salt to reduction to provide a compound of the  
formula :



- 292 -

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 or


- 5 13) subjecting a compound of the formula :



15

or its salt to deacylation reaction to provide a  
 compound of the formula :

20



30


or its salt, in the above formulas;  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 and  
 $R^4_k$  is acyloxy, or

35

- 14) reacting a compound of the formula :

- 293 -

5




(VII)

10

or its salt with a compound of the formula :

15




(VIII)

20

or its salt to provide a compound of the formula :

25



(Is)

30

35


- 294 -

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , X and Y are each as defined above,  
 $Z^2$  is acid residue, and  
 $E_b$  is lower alkylene, or

5

15) reacting a compound of the formula :

10




15

or its salt with an oxidizing agent to provide a  
 compound of the formula :

20

25



30

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R^4$  is lower alkylthio substituted with amino or

35


- 295 -

protected amino; and

$R^4$  is lower alkylsulfinyl substituted with amino or  
protected amino, or lower alkylsulfonyl substituted  
with amino or protected amino; or

5


16) subjecting a compound of the formula :



15

or its salt to catalytic reduction to provide a compound  
of the formula :

20



30

or its salt, in the above formulas,


$R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A,  $E_b$ , X and Y are each as defined  
above, and

35

$E_c$  is lower alkenylene, or


- 296 -

17) subjecting a compound of the formula :



15

or its salt to debenzoylation reaction to provide a compound of the formula :



30

or its salt, in the above formulas,  
 35  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $A$ ,  $E$ ,  $X$  and  $Y$  are each as defined above,  
 $R^1$  is aryl which is substituted with methoxy substituted


- 297 -

with aryl or substituted aryl,  
 $R_e^1$  is aryl which is substituted with hydroxy, or

18) reacting a compound of the formula :

5


10



15


or its salt with a compound of the formula :

20



or its salt to provide a compound of the formula :

25



30

35

- 298 -


or its salt, in the above formulas,  
 $R_e^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined  
 above,

$Z^3$  is hydroxy, or acid residue,


$R^8$  is lower alkyl optionally substituted with acyl,  
 acylamino, protected amino, aryl, substituted aryl,  
 acyl-substituted pyridyl, or N-protected guanidino;  
 and

$R_f^1$  is aryl which is substituted with lower alkoxy  
 optionally substituted with acyl, acylamino,  
 protected amino, aryl, substituted aryl, acyl-  
 substituted pyridyl or N-protected guanidino; or

19) subjecting a compound of the formula :



or its salt to elimination reaction of the hydroxy  
 protective group to provide a compound of the formula :





- 299 -


or its salt, in the above formulas,

$R^1$ ,  $R^2$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,

and

$R_a^3$  is methoxy substituted with aryl; acyloxy; or  
substituted acyloxy; or


20) reacting a compound of the formula :



or its salt with a compound of the formula :



or its salt to provide a compound of the formula :



or its salt, in the above formulas,

$R^1$ ,  $R^2$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,


- 300 -

$Z^4$  is acid residue,


$R^9$  is lower alkyl optionally substituted with esterified carboxy, and

$R_D^3$  is lower alkoxy optionally substituted with esterified carboxy, or

21) subjecting a compound of the formula :



or its salt to deesterification reaction to provide a compound of the formula :



or its salt, in the above formulas,


$R^1$ ,  $R^2$ ,  $R^4$ ,  $R^5$ , A, E, X, and Y are each as defined above,

$R_C^3$  is lower alkoxy substituted with esterified carboxy, and


- 301 -

$R_d^3$  is lower alkoxy substituted with carboxy, or

22) reacting a compound of the formula :



15 or its salt with an alkyne compound in the presence of a palladium compound, a copper compound to provide a compound of the formula :



or its salt, in the above formulas,


30  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  $R_n^4$  is halogen, and

$R_o^4$  is lower alkynyl optionally substituted with hydroxy, amino, protected amino, lower alkylsulfonyloxy or arylsulfonyloxy, or

35

- 302 -


23) reacting a compound of the formula :



or its salt with a compound of the formula :



to provide a compound of the formula :



or its salt, in the above formulas,  
 30  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_p^4$  is lower alkylthio, lower alkyl or lower alkynyl,  
 each of which is substituted with hydroxy,  
 $Z^5$  is halogen,  
 $R^{10}$  is lower alkylsulfonyl or arylsulfonyl, and  
 35  $R_q^4$  is lower alkylthio, lower alkyl or lower alkynyl,


- 303 -

each of which is substituted with lower  
alkylsulfonyloxy or arylsulfonyloxy, or

24) reacting a compound of the formula :

5


10



15

or its salt with alkali metal phthalimide to provide a  
compound of the formula :

20



25

or its salt, in the above formulas,  
R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, A, E, X and Y are each as defined  
above, and  
R<sub>F</sub><sup>4</sup> is lower alkylthio, lower alkyl or lower alkynyl,  
each of which is substituted with phthalimido, or


30

25) reacting a compound of the formula :

35

- 304 -

5




(I-6)

10

or its salt with a reducing agent to provide a compound of the formula :

15



(I-10)

20

25

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined  
 above, and

30


$R_S^4$  is lower alkyl optionally substituted with hydroxy,  
 amino, protected amino, lower alkylsulfonyloxy or  
 arylsulfonyloxy, or

26) reacting a compound of the formula :

35

- 305 -


5



10

or its salt with a compound of the formula :


15



20

or its salt to provide a compound of the formula :

25



30

35

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , X and Y are each as defined

- 306 -


above, and

 $E_d$  is a single bond or lower alkylene, or

27) reacting a compound of the formula :

5

10



15


or its salt with a compound of the formula :



20

in the presence of a base to provide a compound of the formula :

25



30

or its salt, in the above formulas,  
 $R^1$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $G$ ,  $X$  and  $Y$  are each as defined above,  
 $Z^6$  is acid residue, and

35




- 307 -

$R_a^2$  is lower alkyl optionally substituted with aryl or acyl, or

28) reacting a compound of the formula :

5


10



15

or its salt with an acylating agent to provide a compound of the formula :

20



25

30

or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_g^1$  is aryl which is substituted with lower alkoxy  
substituted with amino, and  
 $R_h^1$  is aryl which is substituted with lower alkoxy  
substituted with acylamino or substituted

35


- 308 -

acylamino, or

29) reacting a compound of the formula :

5

10




(I-16a)

15

or its salt with a reducing agent to provide a compound  
of the formula :

20



(I-17)

25

30

or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R^1_i$  is aryl which is substituted with lower alkoxy  
substituted with oxopiperidylcarbonyl, and  
 $R^1_j$  is aryl which is substituted with lower alkoxy

35


- 309 -

substituted with hydroxypiperidylcarbonyl, or

30) reacting a compound of the formula :

5

10




15

or its salt with an amine compound or its salt in the  
 presence of a reducing agent to provide a compound of  
 the formula :

20

25



30

or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_K^1$  is aryl which is substituted with lower alkoxy  
 substituted with formyl or oxopiperidylcarbonyl,


35

- 310 -


and

$R_l^1$  is aryl which is substituted with lower alkoxy  
substituted with aminopiperidylcarbonyl or N-lower  
alkylpiperazinyl, or

31) reacting a compound of the formula :



or its reactive derivative at the carboxy group or a  
salt thereof with lower alkylamino(lower)alkanol to  
provide a compound of the formula :




or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_m^1$  is aryl which is substituted with lower alkoxy  
substituted with carboxy, and  
 $R_n^1$  is aryl which is substituted with lower alkoxy  
substituted with lower


- 311 -

alkylamino(lower)alkoxycarbonyl, or

32) reacting a compound of the formula :



15 or its salt with a reducing agent to provide a compound of the formula :




or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R^1_O$  is aryl which is substituted with lower alkoxy  
 30 substituted with esterified carboxy, and  
 $R^1_P$  is aryl which is substituted with lower alkoxy  
 substituted with hydroxy, or

33) subjecting a compound of the formula :

35

- 312 -

5




(I-22)

10

or its salt to oxidation reaction to provide a compound of the formula :

15



(I-23)

20

25

or its salt, in the above formulas,  
 $R_p^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined  
 above, and


$R_q^1$  is aryl which is substituted with lower alkoxy  
 substituted with formyl, or

30 34) reacting a compound of the formula :

35

- 313 -


5



10

or its salt with an azide compound to provide a compound of the formula :

15



20

25

or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_I^1$  is aryl which is substituted with lower alkoxy  
 substituted with cyano-substituted aryl, and  
 $R_S^1$  is aryl which is substituted with lower alkoxy  
 substituted with tetrazolyl-substituted aryl, or


30

35) reacting a compound of the formula :

35

- 314 -


5



10

or its salt with an isourea compound to provide a compound of the formula :

15



20

25

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_t^4$  is lower alkoxy substituted with amino, and  
 $R_u^4$  is lower alkoxy substituted with guanidino, or

30


36) subjecting a compound of the formula :

35



- 315 -

5




10

or its salt to elimination reaction of the N-protective group to provide a compound of the formula :

15

20




25

or its salt, in the above formulas,  
 $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R^1_{\text{C}}$  is aryl which is substituted with lower alkoxy  
substituted with protected amino, N-protected  
amino(lower)alkanoylamino, N-protected  
piperazinylcarbonyl or N-protected guanidino; and  
 $R^1_{\text{U}}$  is aryl which is substituted with lower alkoxy  
substituted with amino, amino(lower)alkanoylamino,  
piperazinylcarbonyl or guanidino; or


35

- 316 -

37) reacting a compound of the formula :




or its salt with N-lower alkylpiperazine,  
 15 dimethylaminopiperidine, ammonia or N,N-  
 dimethylformamide to provide a compound of the formula :



or its salt, in the above formulas,  
 30  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined above,  
 $R_V^1$  is aryl which is substituted with lower alkoxy  
 substituted with phenoxy carbonylamino, and  
 $R_W^1$  is aryl which is substituted with lower alkoxy  
 35 substituted with N-lower  
 alkylpiperazinyl carbonylamino,  
 dimethylaminopiperidyl carbonylamino, carbamoylamino  
 or dimethylcarbamoylamino, or


- 317 -

38) reacting a compound of the formula :



10

or its reactive derivative at the carboxy group or a  
salt thereof with a hydroxy compound or a diazo compound  
15 to provide a compound of the formula :




25

or its salt, in the above formulas,  
R<sup>1</sup>, R<sup>2</sup>, R<sub>C</sub><sup>3</sup>, R<sub>d</sub><sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, A, E, X and Y are each as  
defined above, or

30 39) reacting a compound of the formula :

35


- 318 -



10

or its reactive derivative at the carboxy group or a salt thereof with an amine to provide a compound of the formula :

15



25

or its salt, in the above formulas,  
 $R^1$ ,  $R^2$ ,  $R_d^3$ ,  $R^4$ ,  $R^5$ , A, E, X and Y are each as defined  
 above, and  
 $R_e^3$  is lower alkoxy which is substituted with carbamoyl  
 optionally substituted with lower alkyl.

30

6. A pharmaceutical composition comprising a compound of claim 1, as an active ingredient, in association with a pharmaceutically acceptable, substantially non-toxic carrier or excipient.

35

7. A compound of claim 1 for use as a medicament.

- 319 -

8. A method of therapeutic treatment and/or prevention of hypertension, heart failure, renal insufficiency, edema, ascites, vasopressin parasecretion syndrome, hepatocirrhosis, hyponatremia, hypokalemia, diabetic, circulation disorder, cerebrovascular disease, Meniere's syndrome or motion sickness which comprises administering an effective amount of a compound of claim 1 to human beings or animals.
9. Use of a compound of claim 1 for the manufacture of a medicament for treating and/or preventing hypertension, heart failure, renal insufficiency, edema, ascites, vasopressin parasecretion syndrome, hepatocirrhosis, hyponatremia, hypokalemia, diabetic, circulation disorder, cerebrovascular disease, Meniere's syndrome or motion sickness in human beings or animals.

# INTERNATIONAL SEARCH REPORT

Intern: 1/ Application No

PCT/JP 96/01533

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C237/42 C07C237/44 A61K31/165 A61K31/33 C07D295/18  
C07D295/20 C07D211/58 C07D211/46 C07D213/80 C07D209/48  
C07C271/16

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BOLL. CHIM. FARM. (1983), 122(4), 190-5 CODEN: BCFAAI; ISSN: 0006-6648, 1983, XP000601221 PLESCIA, S. ET AL: "A new pyrazolo[4,3-c][1,5]benzodiazocine derivative" see page 194; example 4E ---	1,2,5
A	EP.A.0 620 216 (FUJISAWA PHARMACEUTICAL CO) 19 October 1994 cited in the application see claims -----	1-9



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

12 September 1996

Date of mailing of the international search report

20.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Pauwels, G

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP 96/01533

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claim 8 is directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 96/01533

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0620216	19-10-94	AU-A- 5932294	20-10-94
		CA-A- 2121112	14-10-94
		CN-A- 1098406	08-02-95
		HU-A- 70197	28-09-95
		JP-A- 7002800	06-01-95
		US-A- 5521170	28-05-96
		ZA-A- 9402325	16-02-95
-----			